



2017

H-Workload Models and Applications: the First
International Symposium on Human Mental
Workload: Models and Applications

2017

A System to Monitor Cognitive Workload in Naturalistic High-Motion Environments

Bethany K. Bracken

Charles River Analytics, bbracken@cra.com

Seth Elkin-Frankston

Charles River Analytics, selkinfrankston@cra.com

Noa Palmon

Charles River Analytics, npalmon@cra.com

Michael Farry

Charles River Analytics, mfarry@cra.com

Blaise de B Frederick

McLean Hospital/Harvard Medical School, bbfrederick@mclean.harvard.edu

Follow this and additional works at: <http://arrow.dit.ie/hwork17>

 Part of the [Artificial Intelligence and Robotics Commons](#), [Behavior and Behavior Mechanisms Commons](#), [Graphics and Human Computer Interfaces Commons](#), and the [Psychological Phenomena and Processes Commons](#)

Recommended Citation

Bracken, B., Elkin-Frankston, S., Palmon, N., Farry, M. & De B Frederick, B. (2017). A system to monitor cognitive workload in naturalistic, high-motion environments. *H-Workload 2017: The first international symposium on human mental workload*, Dublin Institute of Technology, Dublin, Ireland, June 28-30. doi:10.21427/D7Q33W

This Conference Paper is brought to you for free and open access by the H-Workload Models and Applications: the First International Symposium on Human Mental Workload: Models and Applications at ARROW@DIT. It has been accepted for inclusion in 2017 by an authorized administrator of ARROW@DIT. For more information, please contact yvonne.desmond@dit.ie, arrow.admin@dit.ie, brian.widdis@dit.ie.



A System to Monitor Cognitive Workload in Naturalistic, High-Motion Environments

Bethany K. Bracken, PhD¹; Seth Elkin-Frankston, PhD¹; Noa Palmon, BS¹; Michael Farry, MEng¹; Blaise deB Frederick, PhD²

¹Charles River Analytics, Cambridge, MA; ²McLean Hospital/Harvard Medical School, Belmont, MA

{bbracken, selkinfrankston, npalmon, mfarry}@cra.com; bbfrederick@mclean.harvard.edu

Abstract. Across many careers, individuals face alternating periods of high and low attention and cognitive workload can impair cognitive function and undermine job performance. We have designed and are developing an unobtrusive system to Monitor, Extract, and Decode Indicators of Cognitive Workload (MEDIC) in naturalistic, high-motion environments. MEDIC is designed to warn individuals, teammates, or supervisors when steps should be taken to augment cognitive readiness. We first designed and manufactured a forehead sensor device that includes a custom fNIRS sensor and a three-axis accelerometer designed to be mounted on the inside of a baseball cap or headband, or standard issue gear such as a helmet or surgeon's cap. Because the conditions under which MEDIC is designed to operate are more strenuous than typical research efforts assessing cognitive workload, motion artifacts in our data were a persistent issue. Results show wavelet-based filtering improved data quality to salvage data from even the highest-motion conditions. MARA spline motion correction did not further improve data quality. Our testing shows that each of the methods is extremely effective in reducing the effects of motion transients present in the data. In combination, they are able to almost completely remove the transients in the signal while preserving cardiac and low frequency information in the signal which was previously unrecoverable. This has substantially improved the stability of the physiological measures produced by the sensors in high noise conditions.

1 Introduction

Across many careers, individuals face alternating periods of high and low attention and cognitive workload, which can result in impaired cognitive functioning and can be detrimental to job performance. For example, some professions (e.g., first-responders, doctors and nurses working in an emergency room, pilots) require long periods of low workload (boredom), followed by sudden, high-tempo operations during which they may be required to respond to an emergency and perform at peak cognitive levels. Conversely, other professions (e.g., air traffic controllers, market investors in financial industries, analysts) require long periods of high workload and multi-tasking during which added tasks may result in cognitive overload which can lead to mistakes. It is relatively simple to assess cognitive workload using neurophysiological sensors (e.g., functional near-infrared spectroscopy; fNIRS) when individuals are seated. When cognitive workload increases, there is a corresponding increase in prefrontal blood flow that correlates with increased task engagement. Once the task becomes too difficult, there is a decrease in blood flow that correlates with disengagement from the task and decreased performance [1-4]. However, methods to assess

cognitive workload during normal activities are only recently emerging [5; 6]. Even in these studies, “motion” includes nothing more strenuous than walking [7]. To address this gap, we have designed and are building an unobtrusive system to Monitor, Extract, and Decode Indicators of Cognitive Workload (MEDIC) in realistic, sometimes high-motion environments. Such a system could warn individuals, teammates, or supervisors when steps should be taken to augment cognitive readiness.

2 Method

We first designed a forehead sensor device that includes a custom fNIRS sensor and a three-axis accelerometer designed to be integrated into a baseball cap or headband, or standard issue gear such as a helmet or surgeon’s cap. This sensor is more portable and less obtrusive than most commercially-available sensors. Fig. 1 shows our fNIRS sensor device sensor alone (top left), mounted inside a helmet (top right), being worn during a jump roping task (bottom left), and being worn during a medical training simulation (bottom right).




Fig. 1. Custom fNIRS sensor alone (top left), mounted inside a helmet (top right), worn during jump roping (bottom left), and worn during a medical training simulation (bottom right)

The first full evaluation of this sensor suite included nineteen teams of three undergraduates completing physical and cognitive challenges including: (1) baseline (sitting quietly); (2) word list memorization; (3) balance board; (4) 20 questions; (5) puzzle; (6) hot potato; (7) logic problems; (8) moving boxes; (9) word list recall; and (10) team jump rope. We chose these tasks because the effect on physical and cogni-

tive workload is well validated in the literature, giving us the ability to predict the cognitive and physical workload that should result. A subset of tasks increase only cognitive workload (e.g., word list memorization, 20 questions), a subset of tasks increase only physical workload (e.g., moving boxes), and a subset of tasks increase both cognitive and physical workload (e.g., balance board). In this way, we can evaluate the ability of MEDIC to accurately model changes in cognitive workload during physical activity.

Word list memorization requires the team to remember as many words as possible, either as individuals or as a group. The limits of individual short-term memory are widely documented starting with Braine [8]. At the team level, the concept of transactive memory refers to remembering what is distributed across team members [9]. Balance board requires the team members coordinate rolling a ball edge to edge on a large, flat, weighted board without dropping it for a specified amount of time. For twenty questions, each team member is assigned an object and they need to ask yes-or-no questions (up to 20) of the experimenter to identify the object. This task has been used at the individual level to examine problem solving across the life span [10]. A variant of twenty questions has been used by Shockley at the team level by having a pair compare cartoon pictures [11]. For the puzzle task, we used standard cardboard or plastic puzzles of varying difficulty, giving teams up to two minutes to cooperate on constructing the puzzle. Various tasks have been used to measure individual spatial cognition such as the mental rotation task [12]. A similar task has been used with school-aged children to test interactions in single-sex and mixed-sex dyads during interactions involving Science Technology Engineering and Mathematics (STEM) learning [13]. For hot potato, team members must each maintain balance on a BOSU ball while passing weighted (medicine) balls from one team member to the next. Successful teams will coordinate their timing to maintain stability during the throw and catch while completing the circuit as quickly as possible. Increased postural sway of individual team members being correlated with better team performance has been well-documented across a range of conditions in the motor-coordination literature [11; 14; 15]. For logic problems, individuals were given logic problems (e.g., <http://www.brainbashers.com/logic.asp>). Clues were given in pieces, and teams were allowed to use a large poster board and markers to cooperate to solve the problems. Individual reasoning has been studied by Braine and colleagues [16], and at the team level, research has tied team-based learning to improved critical thinking [17]. For moving boxes, teams must split into dyads to lift and move boxes of variable weights and sizes to construct a wall. Dyads must coordinate with each other to take turns. The size weight illusion—the perception that larger objects are perceived to be lighter than smaller objects of the same mass—is persistent across both individual and team lifting situations. Boxes that are especially large or wide might be incorrectly perceived as light enough to carry safely [18-20]. For team jump rope, team members must jump synchronously to complete a specified number of consecutive jumps. They must coordinate both positioning and jump speed to correctly complete the task. Similar tasks have been used previously to study team coordination [21].

3 Result

Because these conditions are more strenuous than most research assessing cognitive workload, motion artifacts in our data were larger than normal. All wearable NIRS sensors are susceptible to artifacts arising from subject motion; as the subject moves, the sensor can lose contact with the skin or shift position, leading to large transients in the optical transmission, or even baseline shifts. In a typical setting, this is a non-issue because participants or clinical patients are relatively motionless. This is especially problematic when subjects are free to move naturally. These transient or step discontinuities cannot be effectively removed with conventional linear filters – the broad frequency content of the transients mean the signal is not significantly attenuated regardless of the selected frequency range. Moreover, linear filters tend to “ring” in response to transients, leading to contamination of the signal over a wide time period. We therefore have turned to nonlinear methods to remove motion related artifacts. Two of the most effective methods for motion artifact removal in NIRS signals are wavelet transient removal filters applied to the raw optical time course [22] and [23] movement artifact removal algorithm (MARA), which is based on moving standard deviation and spline interpolation applied to the calculated hemodynamic time courses.

There has been significant work comparing techniques for rejecting motion artifacts [4; 24; 25], however, within the large transients, the band-pass filtered raw data was of reasonable quality (i.e., the shape of the raw data was preserved). We needed a technique that would allow us to salvage the data during the high motion event rather than excluding it. We evaluated several standard techniques including MARA spline motion correction [23], wavelet-based filtering [22] and combined MARA and wavelet filtering. Wavelet-based filtering improved data quality to allow salvaging data from even the highest-motion conditions. MARA spline motion correction alone did not further improve data quality. Fig. 2 shows the entire time series of data (left) and a zoomed in window of 20 seconds of data (right). The black dashed line in the figures on the left indicate the data included in the 20 second zoom. The left column shows how motion artifacts are removed on the entire series, while the right column shows how artifacts are removed in detail.




Fig. 2. An entire time series of raw infrared (IR) reflectance data (left) and a zoomed in window of 20 seconds of data (right). The black dashed line in the figures on the left indicate the data included in the 20 second zoom.

4 Conclusion

Our testing shows that each of the methods is extremely effective in reducing the effects of motion transients present in the data. In combination, they are able to almost completely remove the transients in the signal while preserving cardiac and low frequency information in the signal which was previously unrecoverable. This has substantially improved the stability of the physiological measures produced by the sensors in high noise conditions. We are currently analyzing our time-series data to determine whether information on oxygenated blood volume alone is enough to assess cognitive workload as it is with computer-based tasks [1; 3], or whether the effect of physical activity requires the use of additional variables to assess cognitive workload. As we are building up our data processing and modeling pipeline, we are also collecting data for our next evaluation of MEDIC at Vanderbilt Medical School (Nashville, TN, USA) where we will assess medical students, residents, and faculty members during high-fidelity training simulations.

5 Acknowledgements

This work was supported by the U.S. Army Medical Research and Materiel Command under Contract No. W81XWH-14-C-0018. The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision unless so designated by other documentation. In the conduct of research where humans are the subjects, the investigators adhered to the policies regarding the protection of human subjects as prescribed by Code of Federal Regulations (CFR) Title 45, Volume 1, Part 46; Title 32, Chapter 1, Part 219; and Title 21, Chapter 1, Part 50 (Protection of Human Subjects).

References

1. Bunce, S. C., Izzetoglu, K., Ayaz, H., Shewokis, P., Izzetoglu, M., Pourrezaei, K., and Onaral, B. (2011). *Implementation of fNIRS for monitoring levels of expertise and mental workload*.: Springer.
2. Ayaz, H., Cakir, M. P., Izzetoglu, K., Curtin, A., Shewokis, P. A., Bunce, S. C., and Onaral, B. (2012). Monitoring expertise development during simulated UAV piloting tasks using optical brain imaging. *Aerospace Conference, 2012 IEEE*, 1-11.
3. Ayaz, H., Shewokis, P. A., Bunce, S., Izzetoglu, K., Willems, B., and Onaral, B. (2012). Optical brain monitoring for operator training and mental workload assessment. *Neuroimage*, 59, 36-47.
4. Ayaz, H., Izzetoglu, M., Shewokis, P. A., and Onaral, B. (2010). Sliding-window motion artifact rejection for functional near-infrared spectroscopy. *Engineering in Medicine and Biology Society (EMBC), 2010 Annual International Conference of the IEEE*, 6567-6570.
5. Izzetoglu, K., Ayaz, H., Merzagora, A., Izzetoglu, M., Shewokis, P. A., Bunce, S. C., Pourrezaei, K., Rosen, A., and Onaral, B. (2011). The evolution of field deployable fNIR spectroscopy from bench to clinical settings. *Journal of Innovative Optical Health Sciences*, 4, 239-250.
6. McKendrick, R., Parasuraman, R., Murtza, R., Formwalt, A., Baccus, W., Paczynski, M., and Ayaz, H. (2016). Into the Wild: Neuroergonomic Differentiation of Hand-Held and Augmented Reality Wearable Displays during Outdoor Navigation with Functional Near Infrared Spectroscopy. *Frontiers in human neuroscience*, 10.
7. McKendrick, R., Mehta, R., Ayaz, H., Scheldrup, M., and Parasuraman, R. (2017). Prefrontal Hemodynamics of Physical Activity and Environmental Complexity During Cognitive Work. *Human Factors*, 59, 147-162.
8. Miller, G. (1956). The magical number seven, plus or minus two: Some limits on our capacity for processing information. *Psychological Review*, 63, 81-97.
9. Wegner, D. M. (1987). Transactive memory: A contemporary analysis of the group mind. *Theories of group behavior*, 185-208.
10. Denney, N. W. (1985). A review of life span research with the twenty questions task: A study of problem-solving ability. *The International Journal of Aging and Human Development*, 21, 161-173.
11. Shockley, K., Santana, M.-V., and Fowler, C. A. (2003). Mutual interpersonal postural constraints are involved in cooperative conversation. *Journal of Experimental Psychology: Human Perception and Performance*, 29, 326.
12. Shepard, R. N. and Metzler, J. (1971). Mental rotation of three-dimensional objects.
13. Didonato, M., Martin, C., and England, D. (2013). Gendered interactions and their consequences: A dynamical perspective. *Gender and development (Current issues in developmental psychology)*.
14. Collins, J. J. and De Luca, C. J. (1995). The effects of visual input on open-loop and closed-loop postural control mechanisms. *Experimental Brain Research*, 103, 151-163.
15. Kim, S. H. and Chung, M. K. (1995). Rapid Communication Effects of posture, weight and frequency on trunk muscular activity and fatigue during repetitive lifting tasks. *Ergonomics*, 38, 853-863.
16. Braine, M. D. (1990). The "natural logic" approach to reasoning. *Reasoning, necessity, and logic: Developmental perspectives*, 133-157.

17. McNerney, M. J. and Fink, L. D. (2003). Team-based learning enhances long-term retention and critical thinking in an undergraduate microbial physiology course. *Microbiology Education*, 4, 3.
18. Amazeen, E. L. (1999). Perceptual independence of size and weight by dynamic touch. *Journal of experimental psychology: human perception and performance*, 25, 102.
19. Naylor, Y. K. and Amazeen, E. L. (2004). The size-weight illusion in team lifting. *Human Factors: The Journal of the Human Factors and Ergonomics Society*, 46, 349-356.
20. Amazeen, E. L. (2013). Box Shape Influences the Size-Weight Illusion During Individual and Team Lifting. *Human Factors: The Journal of the Human Factors and Ergonomics Society*, 0018720813497980.
21. Chronic pain patients are impaired on an emotional decision-making task
22. Vesper, C., van der Wel, Robrecht PRD, Knoblich, G., and Sebanz, N. (2013). Are you ready to jump? Predictive mechanisms in interpersonal coordination. *Journal of Experimental Psychology: Human Perception and Performance*, 39, 48.
23. Molavi, B. and Dumont, G. A. (2012). Wavelet-based motion artifact removal for functional near-infrared spectroscopy. *Physiological measurement*, 33, 259.
24. Scholkmann, F., Spichtig, S., Muehlemann, T., and Wolf, M. (2010). How to detect and reduce movement artifacts in near-infrared imaging using moving standard deviation and spline interpolation. *Physiological measurement*, 31, 649.
25. Cooper, R., Selb, J., Gagnon, L., Phillip, D., Schytz, H. W., Iversen, H. K., Ashina, M., and Boas, D. A. (2012). A systematic comparison of motion artifact correction techniques for functional near-infrared spectroscopy. *Frontiers in neuroscience*, 6, 147.
26. Tak, S. and Ye, J. C. (2014). Statistical analysis of fNIRS data: a comprehensive review. *Neuroimage*, 85, 72-91.