



2011-01-01

Estimation of Effective Lens Position Using a Method Independent of Preoperative Keratometry Readings

I. Dooley

Waterford Regional Hospital

S. Charalampidou

Waterford Regional Hospital

John Nolan

Waterford Institute of Technology

James Loughman

Dublin Institute of Technology, james.loughman@dit.ie

Molly Breen

Institute of Eye Surgery, Whitfield Clinic, Waterford

See next page for additional authors

Follow this and additional works at: <http://arrow.dit.ie/optomart>



Part of the [Optometry Commons](#)

Recommended Citation

Dooley, I., Charalampidou, S., Nolan, J., Loughman, J., Molloy, L., Beatty, S.: Estimation of Effective Lens Position Using a Method Independent of Preoperative Keratometry Readings. *Journal of Cataract and Refractive Surgery*, Vol.37, no.3, pp.506-12. 2011. doi:10.1016/j.jcrs.2010.09.027.

This Article is brought to you for free and open access by the Department of Optometry at ARROW@DIT. It has been accepted for inclusion in Articles by an authorized administrator of ARROW@DIT. For more information, please contact yvonne.desmond@dit.ie, arrow.admin@dit.ie, brian.widdis@dit.ie.



Authors

I. Dooley, S. Charalampidou, John Nolan, James Loughman, Molly Breen, and Stephen Beatty

Manuscript Number:

Title: Estimation of effective lens position using a method independent of preoperative keratometry readings

Article Type: Full Length Article

Section/Category: Cataract

Keywords: Effective lens position; Scheimpflug Pentacam; Holladay; SRK/T; phacoemulsification cataract surgery.

Corresponding Author: Mr Ian John Dooley, MRCOphth., MSc.

Corresponding Author's Institution: Waterford Regional Hospital

First Author: Ian John Dooley, MRCOphth., MSc.

Order of Authors: Ian John Dooley, MRCOphth., MSc.; Sonia Charalampidou, MRCOphth; John Nolan, PhD; James Loughman, PhD; Laura Molloy, BA; Stephen Beatty, FRCOphth, MD

Abstract: Purpose

To evaluate the validity of a keratometry-independent method of estimating effective lens position (ELP) prior to phacoemulsification cataract surgery.

Setting

Institute of Eye Surgery, Whitfield Clinic, Waterford, Ireland

Methods

The anterior chamber diameter (AGm) and corneal height (Hm) in 95 eyes (95 patients) scheduled for cataract surgery were measured with a rotating Scheimpflug camera. Hm and AGm were used to estimate the ELP in a keratometry-independent method (using the SRK/T [ELPrs] and Holladay 1 [ELPrh] formulae).

Results

The mean (\pm SD) ELP was calculated using traditional (mean ELPs \pm SD: 5.59 ± 0.52 mm and mean ELPh \pm SD: 5.63 ± 0.42 mm) and keratometry-independent (mean ELPrs \pm SD: 5.55 ± 0.42 mm and mean ELPrh \pm SD: 5.60 ± 0.36 mm) methods. Agreement between ELPs and ELPrs and between ELPh and ELPrh were represented by Bland Altman plots, with mean differences (\pm 1.96 SD; range; student's t-test) of 0.06 mm (\pm 0.65 mm; -0.59 to +0.71 mm; $p = 0.08$) in association with ELPrs and -0.04 mm (\pm 0.39 mm; -0.43 to +0.35 mm; $p = 0.08$) in association with ELPrh. The mean absolute error (MAE) for ELPs versus ELPrs estimation and for ELPh versus ELPrh estimation were 0.242 ± 0.222 mm (range: 0.001 to 1.272mm) and 0.152 ± 0.137 mm (range: 0.001 to 0.814 mm), respectively.

Conclusion

This novel study confirms that this keratometry-independent ELP estimation method is comparable to traditional keratometry-dependent methods and may be useful in post-refractive surgery patients.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Estimation of effective lens position using a method independent of preoperative keratometry readings

Authors

Dr. Ian Dooley, MRCOphth^{1,2}

Dr. Sofia Charalampidou, MRCPI, MRCOphth^{1,2}

Dr. John Nolan, PhD⁴

Dr. James Loughman FAOI, PhD³

Ms. Laura Molloy BA²

Mr. Stephen Beatty, FRCOphth, MD^{1,2,4}

Affiliations

1. Department of Ophthalmology, Waterford Regional Hospital, Dunmore Road, Waterford, Ireland

2. Institute of Eye Surgery, Suite 14, Whitfield Clinic, Butlerstown North, Cork Road, Waterford, Ireland

3. Department of Optometry, School of Physics Dublin Institute of Technology, Kevin Street, Dublin 8, Ireland.

4. Waterford Institute of Technology, Cork Road, Waterford, Ireland.

1
2
3 **Running title**
4
5

6
7 Utilising Scheimpflug imaging to estimate effective lens position in a keratometry-
8
9 independent fashion.
10

11
12
13
14
15
16 **Presentations**
17

18
19 This work has been presented at the Irish College of Ophthalmologist's annual
20
21 meeting May 2010.
22
23

24
25
26
27
28 **Funding**
29

30
31
32 No author received any funding for this study.
33
34
35
36
37

38
39 **Declaration**
40

41
42
43 The authors declare that they have no conflict of interest.
44
45
46
47

48
49 **Keywords**
50

51
52
53 Effective lens position; Scheimpflug Pentacam; Holladay; SRK/T;
54
55 phacoemulsification cataract surgery.
56
57
58
59
60
61

Correspondence

Corresponding Author: Dr. Ian Dooley

Address: Department of Ophthalmology, Mater Misericordiae University Hospital,
Eccles street, Dublin, Ireland; Institute of Eye Surgery, Suite 15, Whitfield Clinic,
Cork Road, Waterford, Ireland.

Email: iandooley@eustace.net

Abstract

Purpose

To evaluate the validity of a keratometry-independent method of estimating effective lens position (ELP) prior to phacoemulsification cataract surgery.

Setting

Institute of Eye Surgery, Whitfield Clinic, Waterford, Ireland

Methods

The anterior chamber diameter (AG_m) and corneal height (H_m) in 95 eyes (95 patients) scheduled for cataract surgery were measured with a rotating Scheimpflug camera. H_m and AG_m were used to estimate the ELP in a keratometry-independent method (using the SRK/T [ELP_{rs}] and Holladay 1 [ELP_{rh}] formulae).

Results

The mean (\pm SD) ELP was calculated using traditional (mean $ELP_s \pm$ SD: 5.59 ± 0.52 mm and mean $ELP_h \pm$ SD: 5.63 ± 0.42 mm) and keratometry-independent (mean $ELP_{rs} \pm$ SD: 5.55 ± 0.42 mm and mean $ELP_{rh} \pm$ SD: 5.60 ± 0.36 mm) methods. Agreement between ELP_s and ELP_{rs} and between ELP_h and ELP_{rh} were represented by Bland Altman plots, with mean differences (\pm 1.96 SD; range; student's t-test) of 0.06 mm (\pm 0.65 mm; -0.59 to +0.71 mm; $p = 0.08$) in association with ELP_{rs} and -0.04 mm (\pm 0.39 mm; -0.43 to +0.35 mm; $p = 0.08$) in association with ELP_{rh} . The mean absolute error (MAE) for ELP_s versus ELP_{rs} estimation and for ELP_h versus

1 ELP_{rh} estimation were 0.242 ± 0.222 mm (range: 0.001 to 1.272mm) and $0.152 \pm$
2
3 0.137 mm (range: 0.001 to 0.814 mm), respectively.
4
5
6
7

8 **Conclusion**

9
10
11
12 This novel study confirms that this keratometry-independent ELP estimation
13
14 method is comparable to traditional keratometry-dependent methods and may be
15
16 useful in post-refractive surgery patients.
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Introduction

As a result of improved predictability of refractive outcomes following cataract surgery, patients' expectations are high, and attaining precise postoperative refraction within ± 0.50 D of target is a realistic goal of the conscientious cataract surgeon. The accuracy of predicting the necessary power for an intraocular lens (IOL) is dependent upon the accuracy of several preoperative measurements.^{1 2} These include, depending upon the formula used, some or all of the following: central corneal refractive power (keratometry readings); axial length (biometry); horizontal corneal diameter (horizontal white-to-white); anterior chamber depth (ACD); lenticular thickness; preoperative refraction; and the age of the patient.¹⁻³

Definitions of ACD vary according to context and this should be acknowledged in any discussion of ACD. The clinical definition of ACD in the normal phakic eye is straightforward, and refers to the distance from the cornea to the anterior surface of the lens. Anatomically, ACD refers to the distance between the posterior surface of the cornea, but in an optical context (such as when discussing ACD in an IOL power formula) the distance is normally measured from the anterior surface of the cornea and includes the corneal thickness. This is justified, in part, by the position of the second principal plane of the cornea, which is close to the anterior surface (actually about 0.05 mm in front of the cornea).³

However, the end-point of the ACD distance is much more complex. Many formulae do not use the anterior surface of the IOL as the reference point, but rather the 'effective lens position' (ELP), defined as the effective distance from the anterior

1 surface of the cornea to the lens plane as if the lens was of infinite thinness. The ELP
2 may be back-calculated as the effective ACD 'predicting' the actual postoperative
3 refraction on a given dataset. Hence, the ELP is formula-dependent and need not
4 reflect the true ACD in the anatomical sense. This is the case for an ACD defined by
5 the manufacturer on an IOL container along with the A-constant. The ACD in this
6 context is most often based on the Binkhorst formula and cannot be taken to reflect
7 the true postoperative lens position of the pseudophakic eye.³ For the sake of clarity
8 we will, henceforth, refer to the postoperative ACD as the postoperative IOL position.
9

10
11
12
13
14
15
16
17
18
19
20 Models based on statistically analysed relationships between some or all of
21 the previously mentioned preoperative measurements of the eye and the postoperative
22 IOL position have been used to predict the ELP in the preoperative setting. Thus,
23 estimation of ELP remains an empirical component to all ocular biometric formulae
24 predicting refractive outcomes following cataract surgery, and different models for
25 doing this are important determinants of the accuracy of different formulae for
26 predicting refractive outcomes following cataract surgery.⁴ The predictability of ELP
27 has improved over recent years, attributable to enhancements in the formulae used
28 and the accuracy of the preoperative measurements of ocular biometric variables.
29
30
31
32
33
34
35
36
37
38
39
40
41
42

43 In 1975, Fyodorov derived an equation,⁵ based on an individual eye's
44 keratometry and axial length, to estimate ELP. Indeed, it is well known that in eyes
45 that have undergone corneal refractive surgery the pre-refractive surgery keratometry
46 (pre-op Ks) may be unknown, rendering the predictability of the ELP problematic,
47 thereby contributing to poor predictability of refractive outcomes following cataract
48 surgery in such eyes.⁶
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Ho et al, in 2008, described a regression-analysis derived form of the Fyodorov equation⁵ to calculate a theoretical corneal radius (R_t) which can be used to estimate the ELP, independently of preoperative keratometry readings.⁶

The Pentacam[®] (Oculus Optikgeräte GmbH, Wetzlar, Germany) is a rotating Scheimpflug camera designed to image the anterior segment. It provides topographic maps of the anterior and posterior corneal surfaces, pachymetry maps, and biometric measurements of the anterior segment.⁶⁻⁸

In this way, the Pentacam-measured anterior chamber diameter (measured from anterior chamber angle to anterior chamber angle, AG_m) and corneal height (measured from the internal cornea to the line connecting the anterior chamber angles, H_m) can be used to calculate the theoretical corneal radius (R_t).⁶ Ho et al reported that there was good agreement with this keratometry-independent ELP estimate when compared with traditional keratometry-dependent ELP estimation methods, using the Holladay 1 and SRK/T formulae.⁶ However, postoperative IOL position was not measured in the study by Ho et al, and therefore their conclusions were based solely upon estimation errors pre-cataract surgery.⁶

The keratometry-independent ELP methods, described by Ho et al,⁶ could represent an important advance in cataract surgery for patients who have had previous corneal refractive surgery. We will henceforth refer to this method as the keratometry-independent ELP estimation. We performed a study to compare keratometry-dependent ELP estimation methods versus keratometry-independent ELP estimation methods.

Methods

1
2
3
4 We prospectively studied 95 cataractous eyes of 95 consecutive patients with
5
6 no visually consequential ocular co-morbidity, all of whom underwent uneventful
7
8 phacoemulsification cataract surgery between November 2008 and May 2009. Eyes
9
10 with a history of trauma or ocular surgery were excluded. This study was approved by
11
12 the South East Regional Ethics Committee, and the study protocol adhered to the
13
14 tenets of the Declaration of Helsinki. Valid and informed consent was secured from
15
16 each volunteer prior to enrolment. We certify that all applicable institutional and
17
18 governmental regulations concerning the ethical use of human volunteers were
19
20 followed during this research.
21
22
23
24
25
26

27 All patients underwent a complete ophthalmic examination including,
28
29 automatic keratometry (IOLMaster® version 5 Carl Zeiss Ltd., Welwyn Garden City,
30
31 UK), ocular biometry (IOLMaster® version 5 Carl Zeiss Ltd., Welwyn Garden City,
32
33 UK), and a Pentacam scan.
34
35
36

37 In this study, the Pentacam system was set to acquire 25 images per scan at the
38
39 automatic release mode (images captured automatically). To obtain the Pentacam-
40
41 measured H_m , the Scheimpflug image in the horizontal meridian was displayed. The
42
43 software (version 1.16) showed the locations of the anterior chamber angles. A line
44
45 connecting the 2 points of the anterior chamber angles was drawn. Then, a line was
46
47 drawn from the anterior corneal vertex, which intersected and was perpendicular to
48
49 the line connecting the anterior chamber angles. The distance from the posterior
50
51 corneal surface to the intersection point was termed H_m . The distance between the 2
52
53 anterior chamber angle points was termed AG_m (the measured anterior chamber
54
55 diameter from angle to angle, Figure 1).⁶
56
57
58
59
60
61
62
63
64
65

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

The Appendix outlines the equations used by Ho et al to estimate ELP using 4 different methods (Equations 2 – 5).⁶ Equation 1 is a formula for calculating the theoretical corneal radius (R_{rt}), which is a substitute for keratometry in subsequent keratometry-independent formulae (Equations 3 and 5).⁶ Equation 2 describes a formula that estimates ELP_s , which is the ELP estimate obtained when using the SRK/T formula with keratometry in the traditional fashion. Equation 3 describes a formula that estimates ELP_{rs} , which is the ELP estimate obtained when using the SRK/T formula with the R_{rt} used instead of keratometry. Equation 4 describes a formula that estimates ELP_h , which is the ELP estimate obtained when using the Holladay 1 formula with keratometry in the traditional fashion. Equation 5 describes a formula that estimates ELP_{rh} , which is the ELP estimate obtained when using the Holladay 1 formula with the R_{rt} used instead of keratometry. The ELP was calculated for all study eyes, by the SRK/T and Holladay 1 formula using keratometry-dependent (Equations 2 and 4) and keratometry-independent ELP estimation methods (Equations 3 and 5), respectively.

Surgical procedures

41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

In each study eye, surgical procedures were performed under topical anaesthesia (proxymetacaine hydrochloride 0.5%, Minims, Chauvin Pharmaceuticals, Surrey, UK). All operations were performed through a 2.75 mm superior clear corneal incision. A continuous curvilinear capsulorrhexis was completed following viscoelastic injection (Healon®, Advanced Medical Optics Inc, Santa Ana, California, US), hydrodissection was performed using balanced salt solution (BSS, Alcon laboratories, Inc., UK). The irrigation solution contained 16% gentamicin sulphate (80mg Genticin® Roche, Welwyn Garden City, UK in 500ml BSS PLUS® surgical

1 solution, Alcon laboratories, Inc., UK). After removal of the nucleus using torsional
2 phaco-technology (Infiniti®, Alcon laboratories, Inc., UK), irrigation and aspiration
3 of soft lens matter was performed; then a foldable, posterior chamber intraocular lens
4 (Tecnis® ZA9003, Advanced Medical Optics Inc, Santa Ana, California, US; A-
5 Constant 119.1; ACD_{const} 5.6; Surgeon Factor 1.85) was implanted in the capsular bag
6 with an introducer (AMO EmeraldT Series Unfolder and Cartridge®, Advanced
7 Medical Optics Inc, Santa Ana, California, US). Stromal hydration was then
8 performed to achieve wound integrity; a 10/0 Nylon suture was occasionally placed in
9 the corneal wound when wound integrity was deemed inadequate by the surgeon.
10 Intracameral cefuroxime (1mg Zinacef™, GlaxoSmithKline, UK, in 0.1ml of sterile
11 water for injection, B. Braun Medical Inc., PA, US) was administered via the
12 paracentesis. Then a single drop of 1% apraclonidine (Iopidine® 1%, Alcon
13 laboratories, Inc., UK) and an aliquot of 1% fucidic acid ointment (Fucithalmic® Leo
14 Pharmaceuticals, Ontario, Canada) was administered to the corneal surface.
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33

34 35 **Postoperative examination**

36
37
38
39 The patient was examined two weeks postoperatively, and in cases where a
40 corneal suture was *in-situ*, it was removed at this visit. The patient was then re-
41 examined six weeks postoperatively, when postoperative IOL position was calculated,
42 using the inbuilt calipers on the Pentacam screen (Figure 1); this was done because of
43 the possible failure of the Pentacam to automatically identify the anterior surface of
44 the IOL⁹.
45
46
47
48
49
50
51
52
53
54

55 **Statistical analysis**

56
57
58 Data were analysed by an Aabel software package (version 3.0.3, GigaWiz
59
60
61
62
63
64
65

1 Ltd. Co., Tulsa, OK, USA). Agreement between the respective keratometry-
2 dependent and keratometry independent methods of ELP estimation is represented by
3 Bland-Altman plots, and expressed in terms of mean bias \pm 1.96 standard deviations
4 (SD). Correlations were calculated using Pearson's correlation coefficient (r) and a p-
5 value $<$ 0.05 was taken as statistically significant; the r^2 value is also documented for
6 completeness.
7
8
9
10
11
12
13
14
15

16 We calculated the differences (arithmetic and absolute) between the ELP
17 estimates using traditional keratometry-dependent and their respective and novel
18 keratometry-independent estimation methods. This yielded the mean arithmetic
19 estimation error (ME) and mean absolute estimation error (MAE). When using the
20 SRK/T formula to calculate the ELP_s and the ELP_{rs} , the $ME_s = \text{mean } ELP_{rs} - \text{mean}$
21 ELP_s and the $MAE_s = \text{mean absolute } ELP_{rs} - \text{mean absolute } ELP_s$. When using the
22 Holladay 1 formula to calculate the ELP_h and the ELP_{rh} , the $ME_h = \text{mean } ELP_{rh} -$
23 $\text{mean } ELP_h$ and the $MAE_h = \text{mean absolute } ELP_{rh} - \text{mean absolute } ELP_h$.
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39

40 Results

41
42
43
44
45
46
47

48 The biometric data in 95 unoperated eyes (Table 1) were used to estimate the
49 ELP using keratometry-dependent (ELP_s and ELP_h) and keratometry-independent
50 methods (ELP_{rs} and ELP_{rh}). In our study, the mean ELP_s (\pm SD) calculated using
51 preoperative keratometry values in the SRK/T formula (Equation 2) was 5.59 ± 0.52
52 mm, and was significantly and positively correlated with keratometry-independent
53
54
55
56
57
58
59
60
61
62
63
64
65

1 ELP_{rs} (5.55 ± 0.42 mm) ($r = 0.781$, $r^2 = 66.1\%$, $P < 0.001$, Figure 2), when the R_{rt}
2 values were used in the SRK/T formula (Equation 3). Agreement between ELP_s and
3 ELP_{rs} is represented by the Bland Altman plot in Figure 3, with a mean difference (\pm
4 1.96 SD; range) of $0.06 (\pm 0.65; -0.59 \text{ to } +0.71)$ mm in association with ELP_{rs}. The
5 ME_s for ELP_s estimation versus ELP_{rs} estimation was 0.061 ± 0.241 mm (range: -
6 0.589 to 1.272 mm) and the MAE_s for ELP_s estimation versus ELP_{rs} estimation was
7 0.242 ± 0.222 mm (range: 0.001 to 1.272 mm). Agreement between postoperative
8 IOL position and ELP_{rs} is represented by the Bland Altman plot in Figure 4, with a
9 mean difference (± 1.96 SD; range) of $1.22 (\pm 0.82; +0.40 \text{ to } +2.04)$ mm in
10 association with ELP_{rs}.
11
12
13
14
15
16
17
18
19
20
21
22
23

24 The mean ELP_h (\pm SD) calculated using preoperative keratometry readings in
25 the Holladay 1 formula (Equation 4) was 5.63 ± 0.42 mm, and was significantly and
26 positively correlated with keratometry-independent ELP_{rh} (5.60 ± 0.36 mm) ($r =$
27 0.874 , $r^2 = 76.4\%$, $P < 0.001$, Figure 5), when the preoperative R_{rt} values were used in
28 the Holladay 1 formula (Equation 5). Agreement between ELP_h and ELP_{rh} is
29 represented by the Bland Altman plot in Figure 6, with a mean difference (± 1.96 SD;
30 range) of $-0.04 (\pm 0.39; -0.43 \text{ to } +0.35)$ mm in association with ELP_{rh}. The mean
31 arithmetic estimation error (ME_h = mean ELP_{rh} - mean ELP_h) was -0.037 ± 0.203 mm
32 (range: -0.814 to 0.417 mm) and the mean absolute error (MAE_h = mean absolute
33 ELP_{rh} - mean absolute ELP_h) was 0.152 ± 0.137 mm (range: 0.001 to 0.814 mm).
34 Agreement between the postoperative IOL position and ELP_{rh} is represented by the
35 Bland Altman plot in Figure 7, with a mean difference (± 1.96 SD; range) of $1.27 (\pm$
36 $0.73; +0.54 \text{ to } +2.00)$ mm in association with ELP_{rh}.
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58

59 Discussion

1
2
3
4 The relationship between ELP calculated using keratometry-dependent and
5
6 keratometry-independent methods is positive and significant, and agreement between
7
8 the two techniques is good, and we could expect clinically comparable refractive
9
10 outcomes following cataract surgery using keratometry-independent and
11
12 keratometry-dependent methods of ELP estimation.
13
14
15

16
17 The ME_s and ME_h were slightly greater in our study (0.061 ± 0.241 mm and -
18
19 0.037 ± 0.203 mm, respectively) than those reported by Ho et al (-0.011 ± 0.263 mm
20
21 and -0.0004 ± 0.167 mm, respectively)⁶, but are broadly comparable to those reported
22
23 in the former study, that generated equations 1, 3 and 5 by regression analysis from
24
25 the data in their own study (thus explaining the better agreement that they report). It is
26
27 interesting to note that, in our study, when the SRK/T formulae employs keratometry-
28
29 independent methods to calculate the ELP (ELP_{rs}), it exhibited a slight tendency
30
31 towards overestimation (and consequential hyperopic shift) relative to keratometry-
32
33 dependent ELP estimation methods (ELP_s), in contrast to slight underestimation in the
34
35 Ho study⁶. However, ELP_{rh} yielded a slight underestimation (and consequential
36
37 myopic shift) in both studies⁶. The mean absolute errors of ELP estimation reported in
38
39 our study (ELP_{rs} : 0.242 mm; ELP_{rh} : 0.152 mm) are less than half the distance between
40
41 a sulcus-positioned IOL and an IOL implanted in the capsular bag (approximately
42
43 0.75 mm)¹⁰ which results in a 1.05 D difference in the IOL plane,¹⁰ equivalent to 0.78
44
45 D in the spectacle plane.¹⁰ This is consistent with the findings of Olsen et al, who
46
47 reported that a 0.1mm difference in postoperative IOL position corresponded to a
48
49 0.14D change in power in the IOL plane.¹¹
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

In our study, the approximate mean and maximum discrepancies in the IOL plane were 0.34 D and 1.78 D for ELP_{rs} (mean discrepancy = $MAE_s \times 1.4$; maximum discrepancy = maximum absolute error of $ELP_s \times 1.4$). In terms of the discrepancy in the spectacle plane (discrepancy in IOL plane $\times 0.743$),¹⁰ this would be represented by estimated mean and maximum discrepancies of 0.25 D and 1.32 D, respectively. Similarly the approximate mean and maximum discrepancies in the IOL plane were 0.21 D and 1.14 D for ELP_{rh} (mean discrepancy = $MAE_h \times 1.4$; maximum discrepancy = maximum absolute error of $ELP_h \times 1.4$)¹¹. In terms of the discrepancy in the spectacle plane, this would be represented by estimated mean and maximum discrepancies of 0.16 D and 0.84 D, respectively.

28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

With regard to the ELP_{rs} , the calculated discrepancy in the spectacle plane between keratometry-dependent and keratometry-independent estimation methods was: < 0.5 D in 90.5% of cases; < 1.0 D in 97.9% of cases; < 2.0 D in 100% of cases. With regard to the ELP_{rh} , the calculated discrepancy in the spectacle plane between keratometry-dependent and keratometry-independent estimation methods was: < 0.5 D in 96.8% of cases; < 1.0 D in 100% of cases.

43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

In our study, the mean postoperative manifest refraction spherical equivalent (MRSE) prediction error was 0.51 ± 0.45 D (range -1.95 to +1.16 D; where 66.3 % were < 0.5 D; 87.4 % were < 1.0 D; 100% were < 2.0 D). These results compare favourably with other studies looking at the mean postoperative MRSE prediction error following phacoemulsification cataract surgery, in which the mean postoperative MRSE prediction error is reported as follows: 45.5% to 92.0% of cases were < 0.5 D¹²⁻²¹ and 41% to 100% of cases were < 1.0 D.¹²⁻²¹ In our study, the calculated discrepancies in

1 the spectacle plane between keratometry-dependent and keratometry-independent
2 estimation methods using either SRK/T or Holladay 1, are less than our mean
3
4 postoperative MRSE prediction errors and also compare favourably with typically
5
6 reported mean postoperative MRSE prediction errors in the literature.¹²⁻²¹ These
7
8 calculated discrepancies also compare favourably with studies of patients
9
10 undergoing phacoemulsification cataract surgery, who have previously
11
12 undergone laser in-situ keratomileusis (LASIK), where it is reported that
13
14 between 13% and 67% of patients had a reported mean postoperative MRSE within
15
16 0.5 D of target, that between 25% and 100% had a reported mean postoperative mean
17
18 postoperative MRSE within 1.0 D of target, and that between 79% and 100% had a
19
20 reported mean postoperative mean postoperative MRSE within 2.0 D of target .²²⁻²⁴
21
22
23
24
25
26
27
28

29 Possible sources of the slightly differing results between our study and that of
30
31 Ho et al, in terms of ELP, may result from differences in terms of age and axial length
32
33 between the two sample populations. Our sample had a mean age (\pm SD) of 70.2 (\pm
34
35 10.7) years, over double that of Ho et al's sample at 34.4 (\pm 16.1) years;⁶ given that
36
37 ELP is positively correlated with age (ELP increases with age)⁴, this may account for
38
39 the slight tendency towards overestimation with the SRK/T keratometry-independent
40
41 ELP estimation method in our study, when compared with that of Ho et al.⁶ However,
42
43 this relationship between ELP and age fails to explain the observed discrepancy
44
45 between the slight underestimation seen when using the Holladay 1 keratometry-
46
47 independent ELP estimation method in our study, when compared with that of Ho et
48
49 al.⁶
50
51
52
53
54
55
56

57 While our sample was less myopic, with a mean spherical equivalent of -0.89
58
59 (\pm 4.54) D, compared with that of Ho et al's of -5.84 (\pm 3.92) D,⁶ and this is reflected
60
61
62
63
64
65

1 further in the differing axial lengths of the two studies [mean \pm SD: current study:
2 23.88 (\pm 1.80) mm; Ho et al: 25.73 (\pm 1.59) mm],⁶ it has been shown that estimates of
3 ELP are independent of preoperative refractive error,^{3 25} but correlate positively with
4 axial length.^{3 25} In a study looking at the accuracy of prediction of refractive outcomes
5 following cataract surgery using various formulae including SRK/T and Holladay 1,
6 Narvaez et al found a similar trend (which did not reach statistical significance),
7 reflected in varying refractive outcomes (range of MAE: 0.02 to 0.10 D) post-
8 phacoemulsification cataract surgery between groups of eyes of differing axial
9 lengths,¹⁸ and the observed discrepancy is similar in order of magnitude to the
10 discrepancy between our study and that of Ho et al.⁶ In spite of the differences in
11 population samples, however, it is noteworthy that the keratometry-independent ELP
12 method appears to be relatively robust, with comparable estimation errors (ME_s , ME_h ,
13 MAE_s and MAE_h) in both studies.

14 We compared this novel method of keratometry-independent ELP estimation
15 with the existing keratometry-dependent ELP methods, but also with the anatomic
16 postoperative IOL position, and this relationship has never been previously
17 investigated for the SRK/T and Holladay 1 formulae. We found that the ELP was
18 significantly greater than the anatomic postoperative IOL position by a mean
19 difference of 1.27 mm ($p < 0.001$; student's paired t-test). This is in agreement with
20 the theoretical model proposed by Holladay et al.²⁶ It is worth noting that, despite this
21 discrepancy, the ELP (ELP_{rs} and ELP_{rh}) was significantly and positively correlated
22 with the postoperative IOL position. Jin et al, in a study designed to predict
23 postoperative IOL position using formulae different to those in our study (Haigis
24 algorithm; $r = 0.6$; and Olsen 2 algorithm $r = 0.46$),²⁷ report similar levels of
25 correlation to those seen in our study (postoperative IOL position versus ELP_{rs} : $r =$

1 0.43 and postoperative IOL position versus ELP_{rh} : $r = 0.48$). Although this correlation
2 reached significance ($p < 0.01$ for both methods in our study), the observed
3 correlations between postoperative IOL position and ELP are simply not strong
4 enough in themselves for reliable prediction of postoperative IOL position.²⁷
5
6
7
8
9

10 Refractive laser surgery alters corneal thickness and keratometry, but does not
11 alter ACD or corneal height measured from the endothelial surface (H_m).²⁸ It follows,
12 therefore, that a method of ELP estimation, where the required biometric parameters
13 are unchanged by refractive laser surgery, such as the keratometry-independent ELP
14 estimation method described here, would be extremely useful in post-refractive
15 patients contemplating cataract surgery.
16
17
18
19
20
21
22
23
24
25

26 As there were only 11 post-refractive surgery patients undergoing cataract
27 surgery in Ho et al's paper,⁶ our results would support the view that this novel
28 keratometry-independent ELP estimation method appears to have the validity to
29 warrant formal testing, in the form of a concordance study with existing keratometry-
30 dependent methods^{6 29}, in the context of a large cohort of patients scheduled for
31 cataract surgery, but who have undergone prior refractive laser surgery.
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

References:

1. Holladay JT, Prager TC, Chandler TY, Musgrove KH, Lewis JW, Ruiz RS. A three-part system for refining intraocular lens power calculations. *J Cataract Refract Surg* 1988;14(1):17-24.
2. Holladay JT, Prager TC, Ruiz RS, Lewis JW, Rosenthal H. Improving the predictability of intraocular lens power calculations. *Arch Ophthalmol* 1986;104(4):539-41.
3. Olsen T. Prediction of the effective postoperative (intraocular lens) anterior chamber depth. *J Cataract Refract Surg* 2006;32(3):419-24.
4. Olsen T. Calculation of intraocular lens power: a review. *Acta Ophthalmol Scand* 2007;85(5):472-85.
5. Fyodorov SN, Galin MA, Linksz A. Calculation of the optical power of intraocular lenses. *Invest Ophthalmol* 1975;14(8):625-8.
6. Ho JD, Liou SW, Tsai RJ, Tsai CY. Estimation of the effective lens position using a rotating Scheimpflug camera. *J Cataract Refract Surg* 2008;34(12):2119-27.
7. Lackner B, Schmidinger G, Pieh S, Funovics MA, Skorpik C. Repeatability and reproducibility of central corneal thickness measurement with Pentacam, Orbscan, and ultrasound. *Optom Vis Sci* 2005;82(10):892-9.
8. Barkana Y, Gerber Y, Elbaz U, Schwartz S, Ken-Dror G, Avni I, et al. Central corneal thickness measurement with the Pentacam Scheimpflug system, optical low-coherence reflectometry pachymeter, and ultrasound pachymetry. *J Cataract Refract Surg* 2005;31(9):1729-35.
9. Su PF, Lo AY, Hu CY, Chang SW. Anterior chamber depth measurement in phakic and pseudophakic eyes. *Optom Vis Sci* 2008;85(12):1193-200.
10. Suto C, Hori S, Fukuyama E, Akura J. Adjusting intraocular lens power for sulcus fixation. *J Cataract Refract Surg* 2003;29(10):1913-7.
11. Olsen T. Sources of error in intraocular lens power calculation. *J Cataract Refract Surg* 1992;18(2):125-9.
12. Kugelberg M, Lundstrom M. Factors related to the degree of success in achieving target refraction in cataract surgery: Swedish National Cataract Register study. *J Cataract Refract Surg* 2008;34(11):1935-9.
13. Eleftheriadis H. IOLMaster biometry: refractive results of 100 consecutive cases. *Br J Ophthalmol* 2003;87(8):960-3.
14. Gale RP, Saldana M, Johnston RL, Zuberbuhler B, McKibbin M. Benchmark standards for refractive outcomes after NHS cataract surgery. *Eye (Lond)* 2009;23(1):149-52.
15. Haigis W. [IOL calculation for high ametropia]. *Ophthalmologe* 2008;105(11):999-1004.
16. MacLaren RE, Natkunarajah M, Riaz Y, Bourne RR, Restori M, Allan BD. Biometry and formula accuracy with intraocular lenses used for cataract surgery in extreme hyperopia. *Am J Ophthalmol* 2007;143(6):920-31.
17. MacLaren RE, Sagoo MS, Restori M, Allan BD. Biometry accuracy using zero- and negative-powered intraocular lenses. *J Cataract Refract Surg* 2005;31(2):280-90.

18. Narvaez J, Zimmerman G, Stulting RD, Chang DH. Accuracy of intraocular lens power prediction using the Hoffer Q, Holladay 1, Holladay 2, and SRK/T formulas. *J Cataract Refract Surg* 2006;32(12):2050-3.
19. Olsen T. Improved accuracy of intraocular lens power calculation with the Zeiss IOLMaster. *Acta Ophthalmol Scand* 2007;85(1):84-7.
20. Packer M, Fine IH, Hoffman RS, Coffman PG, Brown LK. Immersion A-scan compared with partial coherence interferometry: outcomes analysis. *J Cataract Refract Surg* 2002;28(2):239-42.
21. Rajan MS, Keilhorn I, Bell JA. Partial coherence laser interferometry vs conventional ultrasound biometry in intraocular lens power calculations. *Eye (Lond)* 2002;16(5):552-6.
22. Diehl JW, Yu F, Olson MD, Moral JN, Miller KM. Intraocular lens power adjustment nomogram after laser in situ keratomileusis. *J Cataract Refract Surg* 2009;35(9):1587-90.
23. Awwad ST, Kelley PS, Bowman RW, Cavanagh HD, McCulley JP. Corneal refractive power estimation and intraocular lens calculation after hyperopic LASIK. *Ophthalmology* 2009;116(3):393-400 e1.
24. Awwad ST, Manasseh C, Bowman RW, Cavanagh HD, Verity S, Mootha V, et al. Intraocular lens power calculation after myopic laser in situ keratomileusis: Estimating the corneal refractive power. *J Cataract Refract Surg* 2008;34(7):1070-6.
25. Holladay JT. Refractive power calculations for intraocular lenses in the phakic eye. *Am J Ophthalmol* 1993;116(1):63-6.
26. Holladay JT, Maverick KJ. Relationship of the actual thick intraocular lens optic to the thin lens equivalent. *Am J Ophthalmol* 1998;126(3):339-47.
27. Jin H, Rabsilber T, Ehmer A, Borkenstein AF, Limberger IJ, Guo H, et al. Comparison of ray-tracing method and thin-lens formula in intraocular lens power calculations. *J Cataract Refract Surg* 2009;35(4):650-62.
28. Rosa N, Lanza M, Capasso L, Lucci M, Polito B, Romano A. Anterior chamber depth measurement before and after photorefractive keratectomy: comparison between IOL master and Orbscan II. *Ophthalmology* 2006;113(6):962-9.
29. Aramberri J. Intraocular lens power calculation after corneal refractive surgery: double-K method. *J Cataract Refract Surg* 2003;29(11):2063-8.

Titles and legends to figures

1
2
3
4 Figure 1. Scheimpflug image from the Pentacam, showing the distance from the
5
6 posterior corneal surface of the vertex to the line connecting the anterior chamber
7
8 angles was termed H_m . The distance between the 2 anterior chamber angle points was
9
10 termed AG_m .
11
12

13
14 Figure 2. Scattergram of the effective lens position (ELP) estimate obtained
15
16 when using the SRK/T formula in a keratometry-dependent method (ELP_s) versus the
17
18 ELP estimate obtained when using the SRK/T formula with the theoretical corneal
19
20 radius (R_{rt}) in a keratometry-independent method (ELP_{rs}). Line of agreement is shown
21
22 ($y = 1x + 0$).
23
24
25
26

27
28 Figure 3. Bland Altman plot of the difference between the effective lens position
29
30 (ELP) estimate obtained when using the SRK/T formula in a keratometry-dependent
31
32 method (ELP_s) and the ELP estimate obtained when using the SRK/T formula with
33
34 the theoretical corneal radius (R_{rt}) in a keratometry-independent method (ELP_{rs}). Line
35
36 of mean difference (dotted line) and limits of ± 1.96 standard deviations (unbroken
37
38 line) are shown.
39
40
41
42

43
44 Figure 4 Bland Altman plot of the difference between the post-op intraocular
45
46 lens (IOL) position and the effective lens position estimate obtained when using the
47
48 SRK/T formula with the theoretical corneal radius (R_{rt}) used instead of keratometry
49
50 (ELP_{rs}). Line of mean difference (dotted line) and limits of ± 1.96 standard deviations
51
52 (unbroken line) are shown.
53
54
55

56
57 Figure 5 Scattergram of the effective lens position (ELP) estimate obtained
58
59 when using the Holladay 1 formula in a keratometry-dependent method (ELP_h) versus
60
61
62
63
64
65

1 the ELP estimate obtained when using the Holladay 1 formula with the theoretical
2 corneal radius (R_{rt}) in a keratometry-independent method (ELP_{rh}). Line of agreement
3
4 is shown ($y = 1x + 0$).
5
6

7
8 Figure 6 Bland Altman plot of the difference between the effective lens position
9 estimate obtained when using the Holladay 1 formula with keratometry in the
10 traditional fashion (ELP_h) and the effective lens position estimate obtained when
11 using the Holladay 1 formula with the theoretical corneal radius (R_{rt}) used instead of
12 keratometry (ELP_{rh}). Line of mean difference (dotted line) and limits of ± 1.96
13 standard deviations (unbroken line) are shown.
14
15
16
17
18
19
20
21
22

23 Figure 7 Bland Altman plot of the difference between the post-op intraocular
24 lens (IOL) position and the effective lens position estimate obtained when using the
25 Holladay 1 formula with the theoretical corneal radius (R_{rt}) used instead of
26 keratometry (ELP_{rh}). Line of mean difference (dotted line) and limits of ± 1.96
27 standard deviations (unbroken line) are shown.
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Effective lens position estimation utilising a keratometry-independent method, compares favourably with existing methods, and could potentially be utilised in post-refractive surgery intraocular lens calculation.

Figure 1
[Click here to download high resolution image](#)




Figure 2
[Click here to download high resolution image](#)




Figure 3

[Click here to download high resolution image](#)




Figure 4
[Click here to download high resolution image](#)




Figure 5
[Click here to download high resolution image](#)




Figure 6

[Click here to download high resolution image](#)




Figure 7

[Click here to download high resolution image](#)




Table 1. Characteristics of the 95 virgin eyes used to develop formulas to derive the effective lens position, (ELP) using keratometry-dependent and keratometry-independent methods. Mean values, standard deviations (SD) and ranges are shown. AG_m = Pentacam-measured anterior chamber diameter from angle to angle; H_m = Pentacam-measured corneal height; mm = millimeters.

Characteristic	Mean \pm SD	Range
Age (years)	70.2 \pm 10.7	38 to 88
Spherical equivalent (Dioptres)	-0.89 \pm 4.54	-20.38 to + 7.63
Anterior corneal radius (mm)	7.79 \pm 0.30	7.25 to 8.25
Axial length (mm)	23.88 \pm 1.80	20.66 to 30.50
H_m (mm)	3.49 \pm 0.37	2.34 to 4.41
AG_m (mm)	11.74 \pm 0.74	9.51 to 13.71

Appendix

[Click here to download Supplementary Material: Appendix ELP July 7th 1.doc](#)

Author Photograph

[Click here to download high resolution image](#)



To: Emanuel S. Rosen, FRCSE

Nick Mamalis, MD

Editors: JCRS

Dr. Ian Dooley

173 The Grange

Raheen

Limerick

Ireland.

086 8555993

iandooley@eustace.net

July 7, 2010

Dear Sirs,

I would appreciate your opinion regarding my original article, entitled:

“Estimation of effective lens position using a method independent of preoperative keratometry readings.” I would be most grateful, if you would please consider this work for publication in the highly esteemed Journal of Cataract and Refractive Surgery.

Sincerely,

Dr. Ian Dooley MRCOphth