



2011

Di-Tert Butyl Chlorosilane

Grainne Hargaden

Dublin Institute of Technology, grainne.hargaden@dit.ie

Timothy O'Sullivan

University College Cork, tim.osullivan@ucc.ie

Follow this and additional works at: <http://arrow.dit.ie/scschcpsart>

 Part of the [Chemistry Commons](#)

Recommended Citation

Hargaden, G. C. O'Sullivan, T. P. Di-Tert Butyl Chlorosilane. 2011.


This Article is brought to you for free and open access by the School of Chemical and Pharmaceutical Sciences at ARROW@DIT. It has been accepted for inclusion in Articles by an authorized administrator of ARROW@DIT. For more information, please contact yvonne.desmond@dit.ie, arrow.admin@dit.ie, brian.widdis@dit.ie.



This work is licensed under a [Creative Commons Attribution-NonCommercial-Share Alike 3.0 License](#)



Di-*tert*-butylchlorosilane



[56310-18-0]

C₈H₁₉ClSi

MW 178.775

Physical Data: bp 82-85 °C/45 mmHg; flash point 39 °C – closed cup; *d.* 0.880 g/cm³

Form: clear, colourless liquid


Preparative Methods: Can be prepared by the chlorination of di-*tert* butylchlorosilane, or by treatment of silicon tetrachloride with *t*-BuLi. {Dexheimer, 1975 #78} {Doyle, 1975 #77}

Purification: Distillation


Handling, Storage and Precautions: Moisture sensitive; avoid strong oxidizing agents and strong bases; flammable; causes burns; avoid inhalation of vapour or mist; use in a fume hood.

Intramolecular reducing agent


Considerable study of alkoxy silanes as intramolecular hydrogen transfer agents has been undertaken (eq X). Curran, 1995 #80} {Curran, 1995 #45} The di-*tert*-butyloxysilanes, prepared from di-*tert*-butylchlorosilane and the requisite alcohol, are especially effective and are generally more stable to column chromatography.



These alkoxy silanes are particularly useful for making five-membered rings *via* a 5-*endo*-trigonal radical cyclisation from a selenide precursor (eq X). {Clive, 1995 #48} {Clive, 1996 #42} It was observed that the (di-*tert*-butyl)silyl group may be difficult to cleave in the absence of an adjacent oxygen function.




This approach has been further developed to encompass polycyclic compounds. {Clive, 2001 #28} The reaction proceeded by a sequential 5-*exo*-diagonal cyclization, followed by 1,5-hydrogen transfer and then a final 5-*endo*-trigonal cyclization. This methodology has been applied to the synthesis of optically pure products when a single enantiomer of the chiral alcohol is used as the initial substrate. {Clive, 2001 #27} Additionally, alkyl iodides can be substituted in place of the selenide precursors. {Martinez-Grau, 1997 #38} {Martinez-Grau, 1995 #81}



The (di-*tert*-butyl)silyl group has also proved effective for the conversion of *cis*-2,5-disubstituted THF derivatives to the corresponding *trans*-2,5-disubstituted rings. {Donohoe, 2008 #9} Activation of the hydroxyl group followed by a 1,2-hydride shift


generates the oxonium ion at the C-2 position. The di-*tert*-butyloxysilane then delivers the hydride stereospecifically to form the *trans*-disubstituted product. This motif is found in many natural-product targets and a similar approach been successfully applied to the synthesis of (+)-Sylvaticin. {Donohoe, 2009 #5}



Silicon-Hydride

Di-*tert*-butylchlorosilane has been used in a number of synthetic sequences which involve a silicon-hydride transfer as the key step.

Treatment of a γ -iodoallylic alcohol with NaH and $t\text{-Bu}_2\text{SiHCl}$ afforded the corresponding silylated alcohol which was then exposed to UV irradiation in the presence of 10% hexabutylditin in a so-called UniMolecular Chain Transfer (UMCT) reaction of silicon hydrides to afford the silicon iodide (eq xx). {Martinez-Grau, 1997 #38}



Di-*tert*-butylchlorosilane has also found application in the Rhodium-catalyzed Si-H insertion of carbenoids, formed by the decomposition of α -diazoesters (eq xx). The chlorosilanes generated can be readily converted to alkoxy silanes by treatment with an alcohol and a base. In a study of a range of chlorosilanes by Landais, it was found that the bulky $t\text{-Bu}_2\text{SiClH}$ was found to be the most reactive in this process.


{Andrey, 1993 #54}

{Andrey, 1995 #46}


{Landais, 1997 #40}

Selective protecting group

Silyl groups have been widely employed as protecting groups for alcohols. Where a choice exists, typical silylation conditions lead to selective protection of the less-hindered hydroxy group. In contrast, di-*tert*-butylchlorosilane can be used for the one-pot silylation of the internal hydroxy group of a 1,2-alkanediol. {Tanino, 1998 #34} The observed selectivity arises from the kinetically controlled ring cleavage of the cyclic silyl ether intermediate where lithium complexes preferentially at the less hindered oxygen. Selectivity was noted to increase when *N,N,N',N'*-tetramethylethylenediamine (TMEDA) was present in the reaction mixture.





Silyl ether protecting groups are of interest in the synthesis of compounds containing vinyl ether groups, such as in the plasmalogens, where other protecting groups strategies invariably lead to decomposition. {Van, 2007 #12}



Protection of diols


Di-*tert*-butylchlorosilane has been used in the preparation of di-*tert*-butylsilyl ditriflate, a highly effective reagent for the protection of a wide range of 1,2-, 1,3- and 1,4-diols under mild conditions (X) {Corey, 1982 #69}



Preparation of alkynylsilanes


Alkynylsilanes are versatile synthetic intermediates which display interesting reactivity. {Molander, 2002 #23} {Mukherjee, 2009 #3} Such groups allow for the introduction of allene substituents at the C3 position of 2,3-epoxyalcohols. {Tanino, 2000 #30} The regioselectivity is dependent on the configuration of the epoxide moiety with *cis*-epoxides proceeding by 5-*exo* type cyclization while *trans*-epoxides undergo a 6-*endo*

cyclization. The allenylsilane intermediates are readily converted to the corresponding allenes with TBAF in 1-methyl-2-pyrrolidione (NMP).



Hydrosilylation of alkynes


Di-*tert*-butylchlorosilane has been utilised in the preparation of (*E*)-di-*tert*-butyl-(1-heptenyl)silanol by hydrosilylation of 1-heptyne and hydrolysis of the intermediate chlorosilanes (eq xx). {Denmark, 2006 #14}




Internal controlling groups

Di-*tert*-butylsilyl ethers have been used as a means of controlling the regioselectivity and stereoselectivity in certain reaction. The ability of silyl groups to stabilise carbocations at the β -position has been exploited in the diastereoselective synthesis of cyclic polyols. {Tanino, 1997 #36} The di-*tert*-butylsilyl ether group undergoes regio- and stereoselective migration in the presence of lithium and 4,4'-di-*t*-butylbiphenyl (DBB) followed by cyclisation under

basic conditions to form the allylsilane. Epoxidation of the double bond occurs primarily on the face opposite to the bulky *t*-butyl group. Ring-opening induced by SiO₂ leads to a β-silyl cationic intermediate and stereoselective introduction of the hydroxy group *via* neighbouring-group participation.




By linking reactive dienes and dienophiles with silaketal tether, the course of the intramolecular Diels-Alder reaction can be controlled with a high degree of stereoselectivity and with regiochemistry opposite to that predicted by bond polarization models. {Gillard, 1991 #61} The cyclisation proceeds in a ‘head-to-tail’ manner and the methyl group on the diene strongly favours an endo cyclisation due to steric factors imposed in the transition state. The di-*tert*-butylsilyl group was found more thermally stable than related alkylsilyls.




Applications in PET imaging

Di-*tert*-butylchlorosilane has been utilised in the synthesis of silicon-based building blocks for ¹⁸F-radiolabeling of peptides for application in PET imaging. {Mu, 2008 #8}

Nucleophilic substitution of di-*tert*-butylchlorosilane with {4-[2-(tetrahydro-2H-pyran-2-yloxy)ethyl]phenyl}lithium proceeded in 74% yield, the product of which was further modified to afford the ¹⁸F-radiolabeled compound. (eq. X)




The lithiated derivative of di-*tert*-butylchlorosilane has been utilized in the synthesis of 3'-silylated thymidine derivatives for application in PET imaging. (eq. X) {James, 2010 #2}




Formation of organometallic complexes

Di-*tert*-butylchlorosilane has been used in the preparation of a range of organometallic complexes, many through an oxidative addition process to Iridium, Manganese and Molybdenum. {Handwerker, 1992 #58} {Koloski, 1994 #53} {Zarate, 1995 #47} {Driess, 1996 #44}



They have also been utilised in the synthesis of aluminium amides (eq xx) and metallasiloxanes containing Si-O-Sn linkages (eq xx). {Choquette, 1992 #60} {Beckmann, 1998 #33}



References

Contributors