



2009-01-01

Transcriptional regulation of cholesterol 24-hydroxylase by histone deacetylase inhibitors.

Marjan Shafaati
Karolinska Institute

Riona O'Driscoll
Dublin Institute of Technology

Ingemar Bjorkhem
Karolinska Institute

Steve Meaney
Dublin Institute of Technology, steve.meaney@dit.ie

Follow this and additional works at: <http://arrow.dit.ie/scschbioart>

 Part of the [Biochemistry Commons](#), [Molecular Biology Commons](#), and the [Other Genetics and Genomics Commons](#)

Recommended Citation

Marjan Shafaati, Riona O'Driscoll, Ingemar Bjorkhem, Steve Meaney: Transcriptional regulation of cholesterol 24-hydroxylase by histone deacetylase inhibitors . *Biochemical and Biophysical Research Communications*, Volume 378, Issue 4, 23 January 2009, Pages 689-694

This Article is brought to you for free and open access by the School of Biological Sciences at ARROW@DIT. It has been accepted for inclusion in Articles by an authorized administrator of ARROW@DIT. For more information, please contact yvonne.desmond@dit.ie, arrow.admin@dit.ie.



This work is licensed under a [Creative Commons Attribution-NonCommercial-Share Alike 3.0 License](#)



Title:

Transcriptional regulation of cholesterol 24-hydroxylase by histone deacetylase inhibitors

Marjan Shafaati¹, Riona O'Driscoll, Ingemar Björkhem¹, Steve Meaney^{1,2*}

1 Division of Clinical Chemistry, Department of Laboratory Medicine, Karolinska University Hospital – Huddinge, Karolinska Institutet, Stockholm, Sweden.

2 School of Biological Sciences, Faculty of Science, Dublin Institute of Technology, Dublin 2, Ireland.

*Present address: School of Biological Sciences, Faculty of Science, Dublin Institute of Technology, Dublin 2, Ireland.

Corresponding Author:

Steve Meaney, PhD

School of Biological Sciences (Rm: 512),

Faculty of Science,

Dublin Institute of Technology,

Kevin Street,

Dublin 2,

Ireland.

Tel: +353-1-402(x)4643

E-mail: steve.meaney@dit.ie

Running Title: Epigenetics and Cholesterol

Abstract:

The mechanistic basis for the expression of tissue specific cholesterol elimination pathways is poorly understood. To gain additional insight into this phenomenon we considered it of interest to investigate if epigenetic mechanisms are involved in the regulation of the brain specific enzyme cholesterol 24-hydroxylase (*CYP46A1*), a key regulator of brain cholesterol elimination. We demonstrated a marked time-dependent derepression of the expression of *CYP46A1*, in response to treatment with the potent histone deacetylase (HDAC) inhibitor Trichostatin A. Additionally, the pattern of expression of the genes in the genomic region surrounding *CYP46A1* was found to be diametrically opposite in brain and liver. Intra-peritoneal injection of HDAC inhibitors in mice led to a significant derepression of hepatic *Cyp46a1* mRNA expression and tissue specific changes in *Hmgcr* and *Cyp39a1* mRNA expression. These results are discussed in the context of the phenomenology of tissue specific cholesterol balance.

Keywords: 24S-hydroxycholesterol, CYP46A1, CYP39A1, epigenetics, histone deacetylase inhibitor, oxysterol, brain cholesterol

Introduction:

Although all nucleated cells are capable of cholesterol synthesis, their ability to do so varies over several orders of magnitude and in the adult state most organisms have developed/reached a certain level of specialisation [1]. The rate of cholesterol synthesis and the cholesterol content of a given tissue are not necessarily directly related, e.g. the adult brain contains more than 25% of whole body cholesterol but turns over only 0.03% of its cholesterol content each day. This is much lower than the developing brain [1] where cholesterol rich myelin is being formed. Brain cholesterol turns over slowly, in marked contrast to that present in most other organs [1; 2]. The mechanistic basis for these differences is unknown, although they are probably a result of the functional isolation of brain and blood cholesterol [3; 4; 5].

According to current concepts the synthesis and elimination of cholesterol in the adult brain is compartmentalised – cells of astroglial lineage are believed to be responsible for the majority of cholesterol synthesis in the adult brain while they contribute relatively little to brain cholesterol elimination. Conversely the neuron specific cytochrome P450 cholesterol 24-hydroxylase (*CYP46A1*) is responsible for the elimination of about two thirds of the cholesterol synthesised by the brain [6; 7]. Recently it was shown that deletion of this gene led to the reduction of the rate of brain cholesterol synthesis and overall maintenance of the total level of brain cholesterol [8]. Due to the importance of this enzyme for brain cholesterol balance we have previously investigated its transcriptional regulation [9], with an expectation that a gene so involved in brain cholesterol balance would be subjected to complex mechanisms of regulation. Surprisingly we found the opposite – the expression of *CYP46A1* was ruggedly insensitive to mechanisms known to regulate the key genes in cholesterol synthesis and elimination, *HMGCR* and *CYP7A1* respectively. This is despite the fact that there are several examples in the literature indicating that *CYP46A1* can be induced under various

conditions [10; 11; 12]. Recently a study by Milagre *et al* indicated that members of the SP1 family of transcription factors may regulate *CYP46A1* [13].

We considered it of high importance to investigate if epigenetic mechanisms were involved in the control of the expression of *CYP46A1*. In this form of regulation, distinct patterns of gene expression are inherited by chromatin modifications, such as DNA and histone methylation, that do not involve changes in DNA [14]. Using a combination of *in vitro* and *in vivo* approaches we show that the expression of *CYP46A1*, as well as other cytochrome P450s involved in cholesterol homeostasis, are potently regulated by histone acetylation status.

Materials and Methods

Materials

Trichostatin A was obtained from Sigma or from Invivogen. Valproate was from Apoteket AB (Stockholm, Sweden). Adult brain total RNA was from Clontech. Adult liver RNA was a generous gift from Dr. Eva Ellis. Cell culture reagents were from Invitrogen. Primers for PCR were from Cybergene AB or assays on demand (Applied Biosystems, Carlsbad, CA).

Cell Culture and Treatment with Trichostatin A

Human SH-SY5Y neuroblastoma cells were cultured in DMEM containing 10% foetal calf serum, 1% Glutamax, 100 U/ml penicillin and 100 µg/ml streptomycin (all from Invitrogen, Carlsbad, CA) in a humidified incubator at 5% CO₂. Cells were seeded into 35 mm dishes and allowed to recover overnight before any addition of reagent. In initial time course experiments cells were exposed to TSA at a final concentration of 0.5 µM. After 0, 0.5, 1, 2, 4, 8 and 24 hours of incubation the medium was removed and cells were scraped into 1 mL of Trizol per dish (Invitrogen, Carlsbad, CA). Cell pellets in Trizol were frozen at -80°C until required. Subsequently cells were seeded into six well plates, allowed to recover and exposed to 0.5 µM TSA for 48 hours. Cells were harvested and stored as described above until required.

Animal Experiment I – Chronic Effects of Valproate Treatment

Six week old male C57/B6-J mice ($n = 5$) were injected intraperitoneally with 700 mg/kg Ergenyl (valproic acid) dissolved in physiological saline. Controls ($n = 5$) received an equal volume of vehicle. Injections were continued for five days and performed between 10 and 11 am. On day six the animals were stunned with CO₂ and killed by cervical dislocation. The

brain and liver were removed, weighed and snap frozen in liquid nitrogen. Organ samples were stored at -80°C until required for analysis.

Animal Experiment II – Acute Effects of Trichostatin A Treatment

C57/B6-J mice (as above) were injected intraperitoneally with increasing doses (0, 2, 5 or 10 mg/kg, respectively) of TSA dissolved in DMSO and diluted in physiological saline to a final DMSO concentration of 18%. The total volume injected was 250 µL. Two hours after injection mice were stunned with CO₂ and killed by cervical dislocation. The brain and liver were removed, weighed and snap frozen in liquid nitrogen. Organ samples were stored at -80°C until required for analysis.

Animal Experiment III – Chronic Effects of Trichostatin A Treatment

Male C57/B6-J mice ($n = 5$) were injected intraperitoneally with 2 mg/kg TSA as described above. Controls ($n = 5$) received an equal volume of vehicle. Injections were continued for nine days and performed between 7 and 8 am. On day nine, approximately three hours following the last injection, the mice were stunned with CO₂ and killed by cervical dislocation. Brain and liver samples were collected and stored as described above.

Gene Expression Analysis

Total RNA was purified from cell or tissue samples using Trizol (Invitrogen, Carlsbad, CA). cDNA was synthesised from 2 µg of total RNA using the High Capacity cDNA Reverse Transcription Kit (Applied Biosystems, Foster City, CA). Steady-state mRNA levels were estimated using either Taqman probes or SYBR green chemistries (primer sequences

available on request). All assays were run on an ABI Prism 7000 Sequence Detection System (Applied Biosystems; Foster City, CA). Estimation of relative gene expression was performed using the comparative threshold cycle method, using cyclophilin A and hypoxanthine-guanine phosphoribosyl transferase (*HPRT*) as endogenous controls for cell culture and tissue samples, respectively.

Long Range Chromatin Mapping

One microgram samples of human adult brain and liver total RNA were used as templates for first strand cDNA synthesis using Superscript III and random hexamers (both from Invitrogen, Carlsbad, CA). Real time quantitative PCR (RT-qPCR) was performed using the Power SYBR green kit and using *HPRT* as endogenous control (primers available on request). Expression levels were calculated according to the comparative threshold cycle method using liver as the reference tissue. The range of expression was calculated by solving the expression $2^{-\Delta\Delta Ct+S}$, where S is the sum of the squared standard deviations (SSD) for each gene in each tissue, which takes into account the variance in the expression of both *HPRT* and the target gene in each tissue.

Lipid Extraction and Analysis

Brain was essentially extracted according to Folch. Approximately 10 mg of brain tissue was added to 1 mL homogenisation buffer (5 mM EDTA, 50 µg/ml BHT in phosphate-buffered saline, pH 7.4) in a clean glass tube and the tissue was disrupted using a polytron homogeniser. Three millilitres of chloroform:methanol (2:1, v:v) were added to the homogenate and the samples were mixed by moderate vortexing. Samples were centrifuged at 5,000 g for five minutes to permit phase separation and the lower organic phase was removed to a new tube.

The aqueous phase was re-extracted five more times as above and the organic phases pooled. Lipids were extracted from liver by placing the entire liver directly into ten volumes of chloroform:methanol (2:1, v:v) for 48 hours. Extracts were dried under argon and re-dissolved in ethanol and stored at -20°C until required. Sterols were analysed by gas chromatography-mass spectrometry using deuterium labelled internal standards as previously described [15].

Statistics

Gene expression data is expressed as mean \pm range as described in Livak [16]. Sterol determinations are expressed as mean \pm standard error of the mean. Statistical comparisons were performed using the unpaired students T-test with the exception of the effects of histone deacetylase inhibitors on *Cyp46a1* expression. In accordance with our hypothesis of that this gene is derepressed by these agents we used the one-tailed Student's t-test. A *P* value of less than 0.05 was considered significant.

Ethical Aspects

All animal experiments received full approval from the local Animal Experimentation Ethics Committee.

Results:

Derepression of CYP46A1 Expression by Inhibition of Histone Deacetylases

The mechanisms underscoring the ectopic expression of *CYP46A1* have yet to be established. We thus investigated if *CYP46A1* expression was regulated by histone acetylation status. Treatment of SH-SY5Y neuroblastoma cells with TSA, a well know inhibitor and potent inhibitor of histone deacetylation, lead to a marked time dependent derepression of *CYP46A1* expression which essentially restored the expression of *CYP46A1* to that of neurons (Fig 1a). This observation was replicated in HepG2 hepatoma cells (results not shown). To further validate these findings we also performed RT-qPCR on SH-SY5Y cells treated with 0.5 μ M TSA for 48 hours and observed a highly significant 150-fold increase in *CYP46A1* expression ($P < 0.00001$) (Fig 1b). For reasons of comparison we also measured the expression levels of other genes known to be important for brain sterol homeostasis. There was no significant effect on the expression of either *HMGCR* or *CYP27A1*, while the expression of *CYP7B1* was decreased to about 10% of the control levels ($P < 0.0001$).

Long-range chromatin structure at CYP46A1

Previously published data indicated that the expression of *CYP46A1* is limited to the brain, with only trace amounts present in other tissues [17]. To determine if the expression of genes in the genomic neighbourhood of *CYP46A1* were coordinated in different tissues we examined the expression levels of all genes in a 1Mb region centred on *CYP46A1* (See Fig 2a). Using the brain and the liver as model organs we show that at this chromosomal region, *CYP46A1* and *DEGS2* were expressed at greater levels in the brain than in the liver (See Fig. 2b). This data is

consistent with previously presented data on these genes, as well as public gene expression datasets (e.g. the Allen Brain Atlas).

Effects of Valproate Treatment on Cholesterol Homeostasis

Given the potent effects of histone deacetylase inhibition on the mRNA levels of *CYP46A1* under *in vitro* conditions, we also considered it important to define if this finding could be replicated in an *in vivo* context. Initially we used the anti-epileptic and anti-anxiolytic valproic acid (VPA), which is well known to have histone deacetylase inhibitor activity. Intraperitoneal injection of a high dose of VPA (700mg/kg) for resulted in a modest induction of the mRNA expression of *Hmgcr* and *Cyp46a1* in the liver, and *Cyp46a1* in the brain (See Fig 3a).

However, treatment with VPA at this level lead to the death of three of the treated mice, two at day three and one at day four indicating that the observed changes in mRNA levels were likely influenced by the general toxic effects of VPA. A similar experiment with a lower dose of VPA (350 mg/kg body weight) led to a similar pattern of changes of mRNA with a significant increase in *Hmgcr* expression ($P < 0.05$) while there was an 1.5-fold increase in *Cyp46a1* ($P < 0.05$). There were no matching changes in hepatic or cerebral steady-state sterol levels (results not shown). To exclude the possibility that the capacity for downstream sterol metabolism (i.e. 7α -hydroxylation 24S-hydroxycholesterol) may have been altered by VPA treatment, we measured the mRNA levels of *Cyp39a1* in both the liver and brain. Surprising, we found a two-fold increase in expression of *Cyp39a1* in the brain ($P < 0.01$) and a 50% decrease in hepatic expression ($P < 0.002$) (Fig 3b).

Effects of TSA Treatment on Cholesterol Homeostasis

In light of the abovementioned toxic effects of high dose VPA treatment we decided to employ the potent histone deacetylase inhibitor TSA. We initially studied the dose dependent effects of TSA treatment in an acute model. IP injection of TSA to male mice at doses from 2mg/kg to 10mg/kg body weight resulted in dose dependent increases in the expression of brain *Cyp46a1* and *Cyp39a1*. This was recapitulated in the liver with increases in hepatic *Cyp46a1* and *Cyp7b1* expression and decreases in both *Hmgcr* and *Cyp39a1* levels (Fig 4a). Intriguingly, the response of female mice to the same regimen was markedly different (results not shown).

We then initiated a treated animals with 2mg/kg TSA for nine days. This resulted in a small-but-significant reduction in brain *Hmgcr* mRNA expression ($P < 0.05$). No significant change was observed in the cerebral expression of *Cyp46a1*, *Cyp39a1* or *Cyp7b1* (Fig 4b). The situation in the liver was similar to that observed following VPA treatment, with a 2.5-fold increase in hepatic *Cyp46a1* expression. In accordance with our hypothesis that inhibition of histone deacetylase increased the mRNA levels of *Cyp46a1*, this was significant following a one-tailed T-test ($P < 0.05$). Again, there were no differences in the brain or liver sterol levels (results not shown).

Discussion

Previously we have shown that the activation of a reporter construct containing firefly luciferase under the control of 2.6 Kb of the human *CYP46A1* promoter was refractory to various interventions known to potently regulate other genes involved in cholesterol regulation [9]. However, several groups have independently demonstrated the presence of ectopic expression of *CYP46A1* in a variety of different states, indicating a potential for regulation. The current study was initiated to determine if epigenetic mechanisms are involved in these phenomena.

The observation that TSA treatment potently derepressed *CYP46A1* expression supports our hypothesis that chromatin condensation was important for the transcriptional activation of *CYP46A1*. Examination of the relative expression of the genes in a 1Mb genomic region centered on *CYP46A1* revealed two areas with markedly higher expression in the brain compared with the liver – *CYP46A1* and *DEGS2*. Taken together these data provide support for the hypothesis that epigenetic mechanisms are involved in establishing the neuronal expression of *CYP46A1*. Inspection of the *CYP46A1* gene revealed two candidate sites for the neuron restrictive silencing factor (*NRSF*), a transcription factor known to be involved in neuron specific gene expression [18]. In pilot studies we attempted to abrogate any effects of *NRSF* on *CYP46A1* expression using a dominant negative *NRSF* mutant construct. However no significant effect on *CYP46A1* mRNA levels was observed (results not shown).

The overall pattern of changes in the mRNA expression of *Cyp46a1* was remarkably consistent under a variety of different treatment schemes, despite the difficulties in normalising gene expression data following treatment with such a potent gene regulator [19]. A further

complication is introduced by the pharmacokinetic profile of TSA – the plasma half life of this compound in mice is in the order of 7-9 minutes and available data indicates significant and rapid hepatic metabolism, potentially to less active compounds [20]. The short half life may be part of the reason for the differences between the acute and chronic TSA mediated regulation of selected genes e.g. *Cyp7b1*.

An intriguing finding of our investigations into the acute effects of TSA was the finding of gender specific responses to effects of the treatment. It has been established that hepatic expression of *Cyp7b1* is greater in male mice, whereas that of *Cyp39a1* is greater in females [21; 22]. This pattern appears to be potentiated by TSA treatment, at least under acute conditions, suggesting that epigenetic mechanisms may be a general regulator of sexually dimorphic cytochrome P450 expression.

A surprising finding was the detection of significant levels of *Cyp39a1* in the brain. This cytochrome P450, which is a 7α -hydroxylase highly specific for 24S-hydroxycholesterol, was originally cloned as liver specific species [21]. Subsequent investigations have identified *Cyp39a1* mRNA or protein in brain or the non-pigmented epithelium of the retina [23; 24]. To the best of our knowledge, this is the first demonstration of transcriptional regulation of *CYP39A1*. At the current time, the reason for the diametrically opposite regulation in brain and liver is unclear. However, this provides a plausible explanation for absence of changes in brain 24S-hydroxycholesterol - induction of the 24S-hydroxycholesterol 7α -hydroxylase within the brain may theoretically lead to intracerebral production of $7\alpha,24S$ -dihydroxycholesterol and mask any increase in 24S-hydroxycholesterol content.

It has previously been shown that epigenetic mechanisms are involved in the regulation of sterol

synthesis – Villagra *et al* demonstrated that lanosterol synthase is repressed by histone deacetylase-3 and presented data that some other genes in the cholesterol synthesis pathway were regulated by histone deacetylase inhibitor [25]. The current data supports the contention that epigenetic mechanisms are of importance for the regulation of the transcription of several cytochrome P450s involved in metabolism and elimination of cholesterol, particular in the context of the brain.

Acknowledgements

The research undertaken was supported by the Swedish Brain Foundation, Swedish Brain Power, the Thuring's Foundation the Stohnes Foundation, the Karolinska Institutet Research Fund and the Swedish Science Council. The initial part of this study was performed with support from Pfizer. We would also like to thank Dr. Elena Feinstein for useful discussions.

References

- [1] J.M. Dietschy, and S.D. Turley, Thematic review series: brain Lipids. Cholesterol metabolism in the central nervous system during early development and in the mature animal. *J Lipid Res* 45 (2004) 1375-97.
- [2] J.M. Dietschy, S.D. Turley, and D.K. Spady, Role of liver in the maintenance of cholesterol and low density lipoprotein homeostasis in different animal species, including humans. *J Lipid Res* 34 (1993) 1637-59.
- [3] S. Meaney, D. Lutjohann, U. Diczfalusy, and I. Bjorkhem, Formation of oxysterols from different pools of cholesterol as studied by stable isotope technique: cerebral origin of most circulating 24S-hydroxycholesterol in rats, but not in mice. *Biochim Biophys Acta* 1486 (2000) 293-8.
- [4] S. Meaney, M. Hassan, A. Sakinis, D. Lutjohann, K. von Bergmann, A. Wennmalm, U. Diczfalusy, and I. Bjorkhem, Evidence that the major oxysterols in human circulation originate from distinct pools of cholesterol: a stable isotope study. *J Lipid Res* 42 (2001) 70-8.
- [5] I. Bjorkhem, and S. Meaney, Brain cholesterol: long secret life behind a barrier. *Arterioscler Thromb Vasc Biol* 24 (2004) 806-15.
- [6] I. Bjorkhem, D. Lutjohann, O. Breuer, A. Sakinis, and A. Wennmalm, Importance of a novel oxidative mechanism for elimination of brain cholesterol. Turnover of cholesterol and 24(S)-hydroxycholesterol in rat brain as measured with $^{18}O_2$ techniques in vivo and in vitro. *J Biol Chem* 272 (1997) 30178-84.
- [7] C. Xie, E.G. Lund, S.D. Turley, D.W. Russell, and J.M. Dietschy, Quantitation of two pathways for cholesterol excretion from the brain in normal mice and mice with neurodegeneration. *J Lipid Res* 44 (2003) 1780-9.
- [8] E.G. Lund, C. Xie, T. Kotti, S.D. Turley, J.M. Dietschy, and D.W. Russell, Knockout of the cholesterol 24-hydroxylase gene in mice reveals a brain-specific mechanism of cholesterol turnover. *J Biol Chem* 278 (2003) 22980-8.
- [9] Y. Ohyama, S. Meaney, M. Heverin, L. Ekstrom, A. Brafman, M. Shafir, U. Andersson, M. Olin, G. Eggertsen, U. Diczfalusy, E. Feinstein, and I. Bjorkhem, Studies on the transcriptional regulation of cholesterol 24-hydroxylase (CYP46A1): marked insensitivity toward different regulatory axes. *J Biol Chem* 281 (2006) 3810-20.
- [10] C.E. Teunissen, S. Floris, M. Sonke, C.D. Dijkstra, H.E. De Vries, and D. Lutjohann, 24S-hydroxycholesterol in relation to disease manifestations of acute experimental autoimmune encephalomyelitis. *J Neurosci Res* 85 (2007) 1499-505.
- [11] C.M. Cartagena, F. Ahmed, M.P. Burns, A. Pajooohesh-Ganji, D.T. Pak, A.I. Faden, and G.W. Rebeck, Cortical Injury Increases Cholesterol 24S Hydroxylase (Cyp46) Levels in the Rat Brain. *J Neurotrauma* 25 (2008) 1087-98.
- [12] N. Bogdanovic, L. Bretillon, E.G. Lund, U. Diczfalusy, L. Lannfelt, B. Winblad, D.W. Russell, and I. Bjorkhem, On the turnover of brain cholesterol in patients with Alzheimer's disease. Abnormal induction of the cholesterol-catabolic enzyme CYP46 in glial cells. *Neurosci Lett* 314 (2001) 45-8.
- [13] I. Milagre, M.J. Nunes, M.J. Gama, R.F. Silva, J.M. Pascussi, M.C. Lechner, and E. Rodrigues, Transcriptional regulation of the human CYP46A1 brain-specific expression by Sp transcription factors. *J Neurochem* 106 (2008) 835-49.
- [14] S.H.L. J. Zlatanova, *Chromatin Structure and Dynamics: State-of-the-Art*, Elsevier, 2004.
- [15] M. Heverin, N. Bogdanovic, D. Lutjohann, T. Bayer, I. Pikuleva, L. Bretillon, U. Diczfalusy, B. Winblad, and I. Bjorkhem, Changes in the levels of cerebral and extracerebral sterols in the brain of patients with Alzheimer's disease. *J Lipid Res* 45 (2004) 186-93.

- [16] K.J. Livak, and T.D. Schmittgen, Analysis of relative gene expression data using real-time quantitative PCR and the 2(-Delta Delta C(T)) Method. *Methods* 25 (2001) 402-8.
- [17] E.G. Lund, J.M. Guileyardo, and D.W. Russell, cDNA cloning of cholesterol 24-hydroxylase, a mediator of cholesterol homeostasis in the brain. *Proc Natl Acad Sci U S A* 96 (1999) 7238-43.
- [18] N. Ballas, and G. Mandel, The many faces of REST oversee epigenetic programming of neuronal genes. *Curr Opin Neurobiol* 15 (2005) 500-6.
- [19] A. Mogal, and S.A. Abdulkadir, Effects of Histone Deacetylase Inhibitor (HDACi); Trichostatin-A (TSA) on the expression of housekeeping genes. *Mol Cell Probes* 20 (2006) 81-6.
- [20] L. Sanderson, G.W. Taylor, E.O. Aboagye, J.P. Alao, J.R. Latigo, R.C. Coombes, and D.M. Vigushin, Plasma pharmacokinetics and metabolism of the histone deacetylase inhibitor trichostatin a after intraperitoneal administration to mice. *Drug Metab Dispos* 32 (2004) 1132-8.
- [21] J. Li-Hawkins, E.G. Lund, A.D. Bronson, and D.W. Russell, Expression cloning of an oxysterol 7alpha-hydroxylase selective for 24-hydroxycholesterol. *J Biol Chem* 275 (2000) 16543-9.
- [22] H. Uppal, S.P. Saini, A. Moschetta, Y. Mu, J. Zhou, H. Gong, Y. Zhai, S. Ren, G.K. Michalopoulos, D.J. Mangelsdorf, and W. Xie, Activation of LXRs prevents bile acid toxicity and cholestasis in female mice. *Hepatology* 45 (2007) 422-32.
- [23] H. Ikeda, M. Ueda, M. Ikeda, H. Kobayashi, and Y. Honda, Oxysterol 7alpha-hydroxylase (CYP39A1) in the ciliary nonpigmented epithelium of bovine eye. *Lab Invest* 83 (2003) 349-55.
- [24] S. Steckelbroeck, M. Watzka, D. Lutjohann, P. Makiola, A. Nassen, V.H. Hans, H. Clusmann, A. Reissinger, M. Ludwig, L. Siekmann, and D. Klingmuller, Characterization of the dehydroepiandrosterone (DHEA) metabolism via oxysterol 7alpha-hydroxylase and 17-ketosteroid reductase activity in the human brain. *J Neurochem* 83 (2002) 713-26.
- [25] A. Villagra, N. Ulloa, X. Zhang, Z. Yuan, E. Sotomayor, and E. Seto, Histone deacetylase 3 down-regulates cholesterol synthesis through repression of lanosterol synthase gene expression. *J Biol Chem* 282 (2007) 35457-70.

Figure Legends

Figure 1

Derepression of cholesterol 24-hydroxylase by histone deacetylase inhibition. a) SH-SY5Y cells were treated with 0.5 μ M of trichostatin A for the indicated time periods and gene expression was evaluated by reverse transcription-PCR; b) SH-SY5Y cells were treated with 0.5 μ M of trichostatin A for 48 hours and gene expression was evaluated by quantitative reverse real-time PCR using. Data is expressed as mean \pm range.

Figure 2

Expression of genes in the genomic neighbourhood of cholesterol 24-hydroxylase is tissue specific. a) The genomic structure of a 1Mb region centred at 14q32.1. The orientation of the individual genes is indicated with an arrow and gene names are as follows: *BCL11B* - B-cell CLL/lymphoma 11B; *SETD3* – SET Domain Containing 3; *CCNK* – Cyclin K; *HHIPL1* – Hedgehog interacting protein like- 1; *CYP46A1* – Cholesterol 24-hydroxylase; *EML1* - Echinoderm microtubule associated protein like-1 ; *EVL* - Enah/Vasp-like; *DEGS2* - Degenerative spermatocyte homolog 2, lipid desaturase; *YY1* – YY-1 Transcription factor; b) Comparative gene expression of this region in human brain and liver. Gene expression in the liver is set to unity and a larger value indicates greater expression in the brain. Data is expressed as mean \pm range, where the range incorporates the variance for both the target gene and the endogenous control in both the liver and brain, respectively (see methods). Note that the two genes with marked overrepresentation of mRNA in the brain (i.e. *CYP46A1* and *DEGS2*) are involved in modification of lipids.

Figure 3

Effects of valproate treatment on hepatic and cerebral gene expression in the mouse. Following intraperitoneal injection of low or high doses of valproate, gene expression in the brain and liver was evaluated. a) Dose dependent effects on the expression of *Hmgcr* and *Cyp46a1*. Treatment with valproate led to dose dependent changes in the mRNA expression of *Hmgcr* and *Cyp46a1* in the liver, and *Cyp46a1* in the brain. b) Divergent regulation of *Cyp39a1* expression in brain and liver. Treatment with valproate lead to a two-fold increase in expression of *Cyp39a1* in the brain, while hepatic levels were reduced to approximately half of the control value (Fig 3b).

Figure 4

Effects of trichostatin A treatment on hepatic and cerebral gene expression in the mouse. Following acute and chronic trichostatin A treatment hepatic and cerebral gene expression were evaluated. a) Treatment with increasing doses of Trichostatin A leads to marked dose dependent changes in gene expression. Male mice were injected intraperitoneally with increasing doses of trichostatin A and killed after two hours. Hepatic mRNA levels of *Cyp46a1* and *Cyp7b1* were markedly increased while *Cyp39a1* and *Hmgcr* levels were suppressed. In the brain *Cyp46a1* was increased while *Cyp39a1* mRNA levels were again reduced. There were no consistent effects on the cerebral expression of either *Hmgcr* or *Cyp7b1*. b) Chronic treatment with 2 mg/kg of trichostatin A leads to modest changes in hepatic and gene expression. In consonance with previous results there was an increase in hepatic *Cyp46a1* mRNA levels was observed ($P < 0.05$) while no significant change was in the brain. Conversely, there was a small but significant reduction in brain *Hmgcr* mRNA levels ($P < 0.05$) while hepatic levels were unchanged. There were no significant changes in the mRNA levels of *Cyp39a1* or *Cyp7b1*.

Figure 1a




Figure 1b




Figure 2a





Figure 2b






Figure 3b




Figure 4a




Figure 4b

