



2011-03-01

Quantitative Assessment of the Shelf-Life of Ozonated Apple Juice

Sonal Patil

Dublin Institute of Technology

Vasilis Valdramidis

University College Dublin, vvaldram@gmail.com

Brijesh Tiwari

Manchester Metropolitan University

Patrick Cullen

Dublin Institute of Technology, picullen@dit.ie

Paula Bourke

Dublin Institute of Technology, paula.bourke@dit.ie

Follow this and additional works at: <http://arrow.dit.ie/schfsehart>

 Part of the [Food Biotechnology Commons](#), [Food Chemistry Commons](#), and the [Food Microbiology Commons](#)

Recommended Citation

Valdramidis V. et al. (2011). Quantitative assessment of the shelf-life of ozonated apple juice. *European Food Research and Technology*, 232(3), 469-477. DOI: 10.1007/s00217-010-1416-2

This Article is brought to you for free and open access by the School of Food Science and Environmental Health at ARROW@DIT. It has been accepted for inclusion in Articles by an authorized administrator of ARROW@DIT. For more information, please contact yvonne.desmond@dit.ie, arrow.admin@dit.ie, brian.widdis@dit.ie.



This work is licensed under a [Creative Commons Attribution-NonCommercial-Share Alike 3.0 License](#)



Antenna & High Frequency Research Centre

Articles

Dublin Institute of Technology

Year 2011

Quantitative assessment of the shelf-life
of ozonated apple juice

Sonal Patil

Vasilis Valdramidis
vvaldram@gmail.com

Brijesh K. Tiwari

PJ Cullen

Paula Bourke

— Use Licence —

Attribution-NonCommercial-ShareAlike 1.0

You are free:

- to copy, distribute, display, and perform the work
- to make derivative works

Under the following conditions:

- Attribution.
You must give the original author credit.
- Non-Commercial.
You may not use this work for commercial purposes.
- Share Alike.
If you alter, transform, or build upon this work, you may distribute the resulting work only under a license identical to this one.

For any reuse or distribution, you must make clear to others the license terms of this work. Any of these conditions can be waived if you get permission from the author.

Your fair use and other rights are in no way affected by the above.

This work is licensed under the Creative Commons Attribution-NonCommercial-ShareAlike License. To view a copy of this license, visit:

- URL (human-readable summary):
<http://creativecommons.org/licenses/by-nc-sa/1.0/>
 - URL (legal code):
<http://creativecommons.org/worldwide/uk/translated-license>
-

1 **Quantitative assessment of the shelf-life of ozonated apple juice**

2

3

4 S. Patil¹, V.P. Valdramidis^{2*}, B.K. Tiwari³, P.J. Cullen¹, P. Bourke¹

5

6 ¹School of Food Science and Environmental Health, Dublin Institute of Technology,

7 Cathal Brugha Street, Dublin 1, Ireland

8

9 ² Biosystems Engineering UCD School of Agriculture, Food Science and Veterinary Medicine University

10 College Dublin Belfield, Dublin 4, IRELAND

11

12 ³Department of Food & Tourism Management, Manchester Metropolitan University, Manchester, M14

13 6HR, UK

14

15

16

17

18

19

20

21

22

23

24

25 * Corresponding author. Tel: +353-876331281; E-mail: vvaldram@gmail.com

26

27

28 **Abstract**

29 Sterile apple juice inoculated with *S. cerevisiae* ATCC 9763 (10^3 CFU/mL) was
30 processed in a bubble column with gaseous ozone of flow rate of 0.12 L/min and
31 concentration of 33-40 $\mu\text{g/mL}$ for 8 min. The growth kinetics of *S. cerevisiae* as an
32 indicator of juice spoilage was monitored at 4, 8, 12 and 16 °C for up to 30 days. The
33 kinetics were quantitatively described by the primary model of Baranyi and Robert's and
34 the maximum specific growth rate was further modeled as a function of temperature by
35 the Ratkowsky type model. The developed model was successfully validated for the
36 microbial growth of control and ozonated samples during dynamic storage temperature of
37 periodic changes from 4 to 16 °C. Two more characteristic parameters were also
38 evaluated, the time of spoilage of the product under static temperature conditions and the
39 temperature quotient, Q_{10} . At lower static storage temperature (4 °C) no spoilage occurred
40 either for unprocessed or ozone processed apple juice. In the case of ozone processed
41 apple juice, the shelf life was increased when compared with the controls and the Q_{10} was
42 found to be 7.17, which appear much higher than that of the controls, indicating the
43 effectiveness of ozonation for the extension of shelf-life of apple juice.

44 **Keywords:** yeast, ozone, apple juice, shelf-life, dynamic modeling

45

46

47

48

49

50

51 **1. Introduction**

52 Acidic products such as fruit juices contain substantial amounts of fermentable sugars.
53 Spoilage of fruit and vegetable juices is primarily due to the proliferation of its natural
54 acid tolerant and osmophilic micro flora [1]. Yeasts, lactic acid bacteria and moulds may
55 account for the fermented taste, production of the carbon dioxide and the buttermilk off-
56 flavour production, as well as the spoilage of juices [2]. Yeasts predominate in spoilage
57 of acid food products as they have the ability to grow at low pH, high sugar concentration
58 and low water activity conditions and resist inactivation by heat processing which enables
59 them to survive or grow in fruit or fruit products [3, 4]. Fruit juices are generally rich in
60 simple carbohydrates and complex nitrogen sources, and hence are ideal substrates for
61 yeasts. More than 110 species of yeasts have been listed as associated with food and food
62 products, of which large proportions occur on fruits, and more than 40 are associated with
63 soft drinks [5]. The contamination of fruit juices with yeasts is normally indicative of
64 highly contaminated raw materials, failure in fruit juice pasteurization, in sanitation
65 practices or the presence of preservative resistant yeasts [6].
66 *Saccharomyces cerevisiae* is one of the most important yeasts causing spoilage of fruit
67 juices and soft drinks [5, 7-9] and can be considered as shelf-life indicator [10, 11].
68 Several authors reported that fruit juice concentrates, fruit pulps, packaged fruit juices
69 and soft drinks are particularly prone to fermentative spoilage with *S. cerevisiae*, *S.*
70 *bayanus* and to a lesser extent *S. pastorianus* [4, 12-18]. Therefore, numerous heat
71 inactivation studies have been conducted with *S. cerevisiae* because of its significance in
72 the spoilage of heat pasteurized fruit juices and carbonated beverages [8, 17, 19].
73 Fermentation of sugars such as glucose, fructose, and sucrose is the principal spoilage

74 reaction of *Saccharomyces* species. Growth of yeasts is usually accompanied by
75 formation of carbon dioxide and alcohol. Carbon dioxide gives the product a gassy,
76 frothy appearance and causes a packaged product to swell and explode. In addition, the
77 products develop a distinctive alcoholic, fermentative smell and taste [20]. Spoilage of
78 fruit juice makes it unacceptable for human consumption.

79 Heat treatment is the most widely used method for preservation of fruit and vegetable
80 juices due to its effectiveness in microbial inactivation [21] although it has certain
81 disadvantages for nutritional and organoleptic values [22, 23]. There is consumer demand
82 for a wider range of less heavily processed foods of improved quality with longer shelf-
83 life and negligible changes in the organoleptic and nutritional values. This has enhanced
84 interest in non-thermal technologies which could be effective on the inactivation of the
85 undesired microorganisms [24].

86 Alternatives to thermal pasteurization such as ozone treatment are under investigation for
87 potential application in fruit juice preservation. Apple juice (or apple cider in North
88 America) is one of these products which is consumed by people of all ages for its sensory
89 and nutritional qualities. The FDA's approval of ozone as a direct additive to food in
90 2001 triggered interest in ozone applications development, and industry guidelines for
91 apple juice and cider were published by the USFDA in 2004, which also highlighted gaps
92 in the scientific knowledge [25].

93 Ozone is a powerful antimicrobial agent due to its potential oxidizing capacity and it
94 appears to be active against bacteria, fungi, viruses, protozoa, as well as bacterial and
95 fungal spores [26, 27]. Ozone destroys microorganisms by progressive oxidation of vital
96 cellular components. Oxidation reactions are caused by either dissolved molecular ozone

97 or free radical species formed during auto-decomposition of ozone [28]. Activated
98 oxygen species resulting from ozone decomposition include singlet oxygen, hydroxyl
99 radical, superoxide anion (perhydroxyl radical at low pH) and hydrogen peroxide which
100 elicit potent cidal activity against a broad-spectrum of microorganisms [29].

101 The objective of this study was to investigate the effect of ozone as a non-thermal
102 treatment to extend the shelf life of an apple juice system. Modeling approaches that
103 describe the growth dynamics of *S. cerevisiae* in previously inoculated ozone processed
104 apple juice under static (isothermal) and dynamic storage temperature conditions are also
105 developed in order to quantitatively assess the effect of ozonation on the shelf life of the
106 product.

107 **2 Materials and Methods**

108 *2.1 Yeast strain and growth conditions*

109 *S. cerevisiae* ATCC 9763 was obtained from microbiology stock culture of the School of
110 Food Science and Environmental Health of the Dublin Institute of Technology, Dublin,
111 Ireland. This strain was maintained as frozen stock at -70 °C in the form of protective
112 beads, which were plated onto potato dextrose agar (PDA, Scharlau Chemie) and
113 incubated at 30 °C for 48 h to obtain single colonies before storage at 4 °C. Working
114 cultures were prepared by inoculating a single colony into malt extract broth (MEB,
115 Scharlau Chemie) and incubating at 30 °C for 24 h.

116 *2.2 Apple juice inoculation*

117 *S. cerevisiae* cells grown for 24 h were harvested by centrifugation (SIGMA 2K15,
118 Bench Top Refrigerated Ultracentrifuge, AGB scientific LTD) at 10,000 rpm for 10min
119 at 4 °C. The cell pellet was suspended in sterile phosphate buffered saline (PBS, Oxoid

120 LTD, UK) re-centrifuged twice as described above. Finally, after two washes with PBS,
121 the cell pellet was re-suspended in PBS and the yeast density was determined by
122 measuring absorbance at 550 nm using McFarland standard (BioMérieux, Marcy -
123 l'Etoile, France).

124 Sterile, commercially prepared apple juice was obtained from a local retailer. This juice
125 was chosen as a food system that could serve for performing controlled microbial
126 experiments (e.g., [10, 30, 31]). The inoculum was then diluted in the juice to obtain
127 approximately 10^6 CFU/mL. For each investigation, the cell concentration was further
128 diluted in apple juice to yield a final working concentration of 10^3 CFU/mL. The
129 inoculated apple juice with *S. cerevisiae* sample was then processed with ozone.

130 Soluble solids content of untreated apple juice was measured using a hand held
131 refractometer (Bellingham and Stanley Ltd., UK). One drop of the juice was placed on
132 the refractometer glass prism and soluble solid content was obtained as Brix. The
133 measured °Brix was 11 ± 0.001 . The pH of untreated product was measured using a pH
134 meter with a glass electrode (Orion Model, England) and was 3.23 ± 0.015 . Titratable
135 acidity was determined by titrating 20 mL of the untreated apple juice sample diluted in
136 80 mL distilled water with 0.1N NaOH using phenolphthalein as an indicator. The
137 volume of NaOH was converted to g malic acid per 100 mL of juice. The measured
138 titratable acidity was 0.45 ± 0.009 .

139 2.3 Ozone treatment

140 Ozone gas was generated using an ozone generator (Model OL80, Ozone services,
141 Canada, Fig. 1). Ozone was produced by a corona discharge generator. Pure oxygen was
142 supplied via an oxygen cylinder (Air Products Ltd., Dublin, Ireland) and the flow rate

143 was controlled using an oxygen flow regulator. Apple juice samples (90 mL) inoculated
144 with *S. cerevisiae* (10^3 CFU/mL) were processed in a 100 mL ozone bubble column with
145 a diameter of approximately 3.7 cm and height of around 21.7 cm. A previously
146 determined optimum flow rate of 0.12 L/min [32] with an ozone concentration of 33-40
147 $\mu\text{g/mL}$ was applied for each treatment for 8 min at ambient temperature (15-18°C) [33].
148 In that study quality (color, phenolic content) and microbial parameters (*E. coli* strains
149 ATCC 25922 and NCTC 12900) during ozone processing were assessed [33]. The ozone
150 concentration was recorded using an ozone analyzer. Excess ozone was destroyed by an
151 ozone destroyer unit. It should be mentioned that the apple juice contains large amount of
152 organic matter which does not permit measurement of dissolved ozone in the liquid phase
153 but also there was not any residual ozone effects as all ozone not targeting on the
154 microbial cells is consumed by the organic matter. All experiments were carried out in
155 duplicate.

156 2.4 Storage study

157 Storage studies were performed for the following three types of samples. Apple juice
158 inoculated with 10^3 CFU/mL served as an unprocessed control 1. The second sample was
159 the ozonated apple juice. Subsequently an unprocessed control 2 was prepared by
160 inoculating *S. cerevisiae* cells with an inoculum level of 10^1 CFU/mL in order to start
161 with a similar inoculum level that was attained after 8 min of ozone treatment.

162 2.4.1 Static storage temperature study (SST)

163 Unprocessed control samples of apple juice and ozone processed apple juice samples (45
164 mL each) were stored at constant temperatures of 4, 8, 12, and 16 °C respectively in
165 incubators (LMS cooled incubators, Lennox Laboratory Supplies, Dublin, Ireland) for a

166 period up to 30 days. Aliquots of unprocessed and processed samples were taken daily for
167 analysis.

168 *2.4.2 Dynamic storage temperature study (DST)*

169 For the DST study, unprocessed and processed apple juice samples were stored in an
170 incubator where the lowest and the highest temperatures were set to 4 and 16 °C. The
171 temperature was programmed to fluctuate according to a profile consisting of 4 °C for 12
172 h, followed by an increase of temperature from 4 to 16 °C and maintained at 16 °C for a
173 further 12 h. The actual temperature profiles were recorded every 10 min using a
174 temperature sensor connected to a data logger (Grant 1000 series Squirrel meter/data
175 logger, UK). This specific profile was chosen in order to create a scenario of temperature
176 abuse enhancing the microbial growth on which the developed modeling approaches
177 could be validated.

178 *2.5 Microbiological analysis*

179 Yeast populations were determined by plating onto PDA. Aliquots (1mL) were
180 withdrawn every day from ozone processed and unprocessed juice stored at each different
181 temperature, serially diluted in MRD and 0.1mL of appropriate dilutions were surface
182 plated on PDA in duplicate. Plates were incubated at 30 °C for 48 h and colony forming
183 units were counted. Results were reported as Log₁₀CFU/mL.

184 *2.6 Microbial modeling*

185 *2.6.1 Parameter identification under static conditions*

186 *S. cerevisiae* growth data in ozone processed apple juice stored under SST conditions
187 were fitted to the explicit version of the Baranyi, and Roberts [34] model (Eq.1-3).

188 Regression was performed by using the DMFit Excel add-in software, version
189 2.1(www.ifr.ac.uk/safety/DMFit). The model reads as follows

190

$$191 \quad N(t) = N(0) + \mu_{\max} A(t) - \ln \left(1 + \frac{e^{\mu_{\max} A(t)} - 1}{e^{(N_{\max} - N(0))}} \right) \quad (1)$$

$$192 \quad \text{with } A(t) = t + \frac{1}{\mu_{\max}} \ln \left(\frac{e^{(-\mu_{\max} t)} + q(0)}{1 + q(0)} \right) \quad (2)$$

193

$$194 \quad \text{and } \lambda = \ln \left(\frac{1 + \frac{1}{q(0)}}{\mu_{\max}} \right) \quad (3)$$

195 The kinetic parameters of maximum specific growth rate (μ_{\max}) (1/days), lag phase (λ)
196 (days), initial microbial population ($N(0)$) (Log₁₀CFU/mL) and maximum population
197 density (N_{\max}) (Log₁₀CFU/ml) have then been estimated. $q(0)$ (-) denotes the
198 concentration of substance critical to the microbial growth and is related to the
199 physiological state of the cells.

200 The maximum specific growth rates estimated under SST conditions were further
201 modeled as a function of storage temperature by using the Square root model [35, 36]:

202

$$203 \quad \mu_{\max} = b(T - T_{\min})^2 \quad (4)$$

204 where b is a constant, T is the storage temperature (°C), T_{\min} is the theoretical minimum
205 temperature for the growth of the organism. Eq. (4) has been used without the commonly
206 applied square root transformation of the μ_{\max} value. This required the performance of a
207 non-linear regression which is available from the DMFit software. A (geometric) mean

208 value for $h_0 = \lambda * \mu_{\max}$ for each of the experimental set-ups (Control 1, Control 2,
 209 Ozonated) was estimated from the individual growth curves, considering that the
 210 parameter is constant, independent of the storage temperature [34, 37, 38] and the fact
 211 that the resulting h_0 was derived from the 3 levels of temperatures (refer to results). $q(0)$
 212 is related to the parameter h_0 by the following equation [34]:

213

$$214 \quad q(0) = \frac{1 - e^{-h_0}}{e^{-h_0}} \quad (5)$$

215 2.6.2 Model validation under dynamic storage temperature (DST) conditions

216 The validation of the yeast growth model was performed under DST conditions based on
 217 the time temperature profile of apple juice samples during storage (control and ozone
 218 processed), in conjunction with the square root model Eq. (4). The predictions were
 219 performed with the differential equation of Baranyi and Roberts model (Eq. (6), (7)) in
 220 which the Runge-Kutta method (ode23s, Matlab, The Mathworks) was applied for the
 221 approximation of solutions of these ordinary differential equations:

$$222 \quad \frac{dN(t)}{dt} = \left(b(T(t) - T_{\min})^2 \right) \left(\frac{q(t)}{q(t) + 1} \right) \left(1 - \frac{N(t)}{N_{\max}} \right) N(t) \quad (6)$$

$$223 \quad \frac{dq(t)}{dt} = \left(b(T(t) - T_{\min})^2 \right) q(t) \quad (7)$$

224 The root mean squared error (Eq. 8) [39] was used for evaluating the model fitting while
 225 the accuracy and the bias factors presented by Baranyi et al. [40](Eq. 9, 10) were
 226 considered in order to assess the prediction capability of the developed model.

$$227 \quad \text{RMSE} = \sqrt{\frac{\sum_{i=1}^{n_t} (y_{\text{exp}i} - y_{\text{pre}})^2}{n_t - n_p}} \quad (8)$$

228 Where y_{expi} are experimental observations, y_{pre} are model predictions, n_t are number of
 229 data points and n_p are number of estimated model parameters.

$$230 \quad A_f = 10 \sqrt{\frac{\sum_{i=1}^n \left(\log_{10} \hat{N}_i - \log N_i \right)^2}{n}} \quad (9)$$

$$231 \quad B_f = 10 \frac{\sum_{i=1}^n \left(\log_{10} \hat{N}_i - \log_{10} N_i \right)}{n} \quad (10)$$

232 Where $\log_{10} \hat{N}_i$ is the predicted microbial load and n is the number of the experimental
 233 measurements.

234 *2.6.3 Calculation of the Q_{10} value*

235 The temperature quotient (Q_{10}) was also calculated from the information obtained in
 236 Section 2.6.1 (parameter identification under static conditions). Q_{10} shows the effect of
 237 temperature on the shelf-life and it is given as follows [41, 42]:

$$238 \quad Q_{10} = \frac{\text{shelf life at } T^{\circ}\text{C}}{\text{shelf life at } (T + 10^{\circ}\text{C})} \quad (11)$$

241 Observe that this parameter was developed for a zero order reaction when the influence
 242 of temperature on the reaction rate is described by using the Arrhenius relationship [43].
 243 Nevertheless this approach is proposed and applied for the current microbial kinetic study
 244 as an alternative method to assess the efficacy of the ozonated juice.

245 This Q_{10} value can be easily calculated by performing a regression between the ln shelf
 246 life (days) versus the temperature which yields a straight line. Consequently, $Q_{10} = \exp$
 247 $(10 \cdot k)$ with k the slope of the regression line. The estimation of the time of the shelf-life
 248 (t_s) was calculated considering that a microbial level $> 10^6$ CFU/mL resulted in a failure

249 (spoilage) of the product (see for similar examples in other products: Al-Kadamany, et al.
250 [44]). The shelf-life time, t_s , was obtained by solving Eq. (1-3) (*solve* command in
251 Matlab, The Mathworks) for the estimated parameters of the two controls and the
252 ozonated growth kinetics when $\log N(t_s) = 6 \log(\text{CFU/mL})$.

253 **3 Results**

254 The growth of *S. cerevisiae* in unprocessed and ozone processed apple juice was assessed
255 at SST conditions from 4 °C to 16 °C. Representative growth curves of the yeast
256 population are shown in Fig. 2. The initial inoculum of control 1 was similar to
257 previously reported levels of 10^3 CFU/mL [30, 45], while this level has also been
258 reported in sound apples [46]. Finally, Kisko et al. [47] recorded *ca.* 10^3 CFU/mL level of
259 *S. cerevisiae* in unprocessed apple juice. In the case of the unprocessed control samples 1
260 and 2 (i.e., initial inoculum level of 3.0 and 1.30 log CFU/mL, respectively) the lag phase
261 was not obvious when the juice was stored under high SST (12 °C and 16 °C) (Fig. 2a
262 and 2b). However, a typical growth pattern of *S. cerevisiae* was observed in the ozone
263 processed apple juice stored under SST of 12 °C and 16 °C, consisting of an initial lag
264 phase, an exponential growth phase followed by a stationary phase (Fig.2c).

265 The estimated kinetic parameters and statistical indices resulting from the regression of
266 the microbial data by the Baranyi and Roberts model are shown in Table 1. The values of
267 μ_{max} and λ varied according to the storage temperature. The μ_{max} of the unprocessed
268 control samples increased from 0.35 log CFU /day to 1.23 log CFU /day and for ozone
269 processed apple juice increased from 0.275 log CFU /day to 1.270 log CFU /day with
270 increase of the temperature from 8 to 16 °C. However, the lag phase for ozone processed
271 apple juice was decreased from 15.07 days at 8 °C to 2.84 days at 16 °C. For both

272 unprocessed and ozone processed apple juice, the maximum population density (N_{max})
273 was found to be unaffected when stored under high SST (12 °C and 16 °C). The effect of
274 storage temperature on μ_{max} was further modeled as a function of temperature by using
275 the secondary square root model. The estimated parameters of the model are shown in
276 Table 2. The model described satisfactorily the effect of temperature on the growth of *S.*
277 *cerevisiae*. The calculated value for the theoretical minimum temperature of growth in
278 ozone processed apple juice was 0.28 °C. The h_0 values obtained for the static
279 environments studied were 0.336, 0.671 and 3.417 for unprocessed control 1,
280 unprocessed control 2 and ozone processed apple juice samples, respectively.

281 The model developed under SST conditions was validated under DST conditions by
282 using a periodically changing temperature profile and performing predictions with Eq. (6)
283 and (7). As the maximum population density was independent of the applied storage
284 temperature it was fixed at 7.5 logs CFU/mL (average of N_{max} estimated during
285 isothermal conditions for which microbial stationary phase was reached). For the initial
286 concentration $N(0)$, a nominal value was taken from the measured plate count result, i.e.,
287 3.02 (for control 1), 1.32 (for control 2), 1.24 (for ozonated) log (CFU/mL). Finally, the
288 nominal values for $q(0)$ were 2.49, 1.05 and 0.03 for control 1, control 2 and ozonated
289 apple juice respectively, calculated using Eq.(5) and after estimation of the h_o from the
290 parameters derived under static environmental conditions. The comparison between the
291 predicted and observed growth of *S. cerevisiae* in unprocessed apple juice and ozone
292 processed apple juice samples are shown in Fig. 3. The performance of the model was
293 evaluated statistically by the calculation of the bias (B_f) and accuracy (A_f) factors.

294 Two more characteristic parameters were evaluated, the Q_{10} and the time of spoilage of
295 the product under SST conditions (Fig. 4). At the lowest SST (4 °C) no spoilage occurred
296 either for unprocessed or ozone processed apple juice. However, with the higher SST's
297 used product spoilage was observed in 9.45, 3.78, and 2.35 days for unprocessed control
298 1 at 8, 12 and 16 °C, respectively. For unprocessed control 2, the spoilage occurred after
299 15.08, 6.30 and 4.29 days at 8, 12 and 16 °C respectively. In the case of ozone processed
300 apple juice, the shelf life was increased when compared with both type of controls and
301 resulted in 34.26, 10.34 and 7.08 days at 8, 12 and 16 °C, respectively. Finally the Q_{10}
302 was found to be 7.17 in the case of ozonated juice. This was much higher than that
303 obtained for the controls, i.e., 5.68, 4.81, indicating the effectiveness of ozonation for
304 extension of the shelf-life of apple juice.

305 **4. Discussion**

306 The results of the present study showed that *S. cerevisiae* ATCC 9763 is able to grow in
307 apple juice stored within a temperature range of 8 to 16 °C. The Baranyi, and Roberts
308 model as well as the square root model described the growth of yeast populations in
309 unprocessed and ozone processed apple juice. Based on the static data, a new model was
310 developed that described the growth of *S. cerevisiae* population well in unprocessed and
311 ozone processed apple juice under dynamic conditions that simulated a storage
312 temperature abuse. At the lower SST's (4 and 8 °C), the longer lag phase indicates that
313 the yeast population needed longer time to adapt to the environment. However, at higher
314 storage temperatures this effect was not evident, indicating the ability of yeasts to grow at
315 these temperatures with a reduced or seemingly absent lag time. By comparison, in the
316 case of ozone processed apple juice stored at 8, 12 or 16 °C, the lag phase (λ) was

317 increased, indicating the effect of temperature and applied ozone stress on growth of *S.*
318 *cerevisiae* populations. Panagou et al. [10] reported a very short lag phase in different
319 pasteurized fruit juices even at the lowest storage temperatures, suggesting that
320 inoculated yeasts' adaptation time was unaffected by these temperatures (4, 8, 12 and 16
321 °C) . However, in this study a lag phase was observed for all ozone processed samples.
322 This could be due to the oxidizing action of the applied ozone treatment, which may exert
323 additional stress prior to allowing growth. Ozone has been reported to inactivate cytosolic
324 enzymes, with the most drastic inactivation for glyceraldehyde 3 phosphate
325 dehydrogenase and to lesser extent to other cytosolic enzymes. It also affects the quantity
326 of ATP and other nucleoside triphosphates, reducing to about 50% of its initial level [48].
327 The performance of the developed model was validated under dynamic conditions. Ross
328 et al. [49] reported that predictive models should ideally have an A_f and $B_f = 1.00$,
329 indicating a perfect model fit where the predicted and actual response values are equal
330 and satisfactory. B_f limits are more difficult to define because limits of acceptability are
331 related to the specific application of the model. Ranges of 0.6-3.99 have been reported for
332 the growth pathogen and spoilage microorganisms when compared with independent
333 published data [49]. The values of B_f and A_f indicated good agreement between observed
334 data and predicted data points. Nevertheless, in the case of Control 2 some discrepancy
335 was evident (Table 3). This could be attributed to the effect of the inoculum size on the
336 microbial adaptation phenomena. This observation may require further evaluation of the
337 inoculum size effects which could elucidate if different values of h_0 should be considered
338 for each of the performed microbial predictions.

339 Different technologies have been applied for inhibiting the growth of spoilage
340 microorganism in fruit juices. Patrignani et al. [30] evaluated the potential of high
341 pressure homogenization (HPH) for inactivation of *S. cerevisiae* 635 inoculated in apricot
342 and carrot juice and its shelf life extension. Four or more repeated passes at 100 MPa of
343 HPH to the apricot juice samples inoculated at a level of 3 log₁₀ CFU/mL showed that *S.*
344 *cerevisiae* population remained under the detection limit at least up to 144 h at 25 °C. For
345 carrot juice samples subjected to five or more repeated HPH passes, the *S. cerevisiae* cell
346 load was lower than 5 log₁₀ CFU/mL after 144 h at 25 °C. However, refrigerated storage
347 (4 °C) indicated satisfactory extension of shelf life of HPH processed juices. Qin et al.
348 [50] reported over 3 weeks extension of standard shelf life of pulsed electric field (PEF)
349 processed apple juice when stored at 4 °C and 25 °C. Ferrentino et al. [51] concluded that
350 high pressure carbon dioxide (HPCD) treatment proved to be a promising alternative
351 technique yielding juices with fresh-like characteristics and extension of shelf life with
352 safety. Suarez-Jacobo et al. [52] reported the efficacy of ultra high pressure
353 homogenization to develop fresh apple juice with an equivalent shelf life to pasteurized
354 apple juice with respect to the microbiological characteristics. Valdramidis et al. [53]
355 observed that no spoilage of apple juice was evident at storage temperatures of 4, 8 and
356 12 °C for 36 days after treatment with high hydrostatic pressure at 500 MPa and 550
357 MPa. From the present work it is evident that ozone is another non thermal technology
358 which can be employed for extending the shelf life of apple juice. The present results
359 proved an increase of the shelf-life of the ozonated product that varied between 2.79 to
360 24.81 days depending on the storage temperatures when compared with the control
361 samples.

362 Validation of the developed modeling approaches will be expanded based on the
363 suggestions by Pin et al. [54]. More specifically, kinetic data that come from competition
364 of inoculated *S. cerevisiae*, pathogenic microorganism with a naturally occurring
365 microflora of fresh apple juice will be incorporated in future model developments while
366 comparative studies between ozonated and other treated technologies will be applied.
367 This will permit the application of this model to apple juice products with different
368 properties. Further studies will focus on defining the failure (spoilage) of processed apple
369 juice based on the effect of ozone on additional to previously reported quality parameters
370 (e.g., color, phenolic content) [33] including volatiles responsible for flavor, odour and
371 sensory evaluation. Effect of the different inoculums levels on the microbial adaptation
372 phenomena will also be assessed to interpret possible modeling discrepancies.

373 **Acknowledgement**

374 Funding for this research was provided under the National Development Plan 2000-2006,
375 through the Food Institutional Research Measure, administered by the Department of
376 Agriculture, Fisheries & Food, Ireland.

377 **References**

- 378 1. Tahiri I, Makhlouf J, Paquin P, Fliss I (2006) Inactivation of food spoilage bacteria and
379 *Escherichia coli* O157:H7 in phosphate buffer and orange juice using dynamic high
380 pressure. Food Res Int 39:98–105
- 381 2. Tournas VH, Heeres J, Burgess L (2006) Moulds and yeasts in fruit salads and fruit
382 juices. Food Microbiol 23:684–688

- 383 3. Put HMC, De Jong J (1980) The heat resistance of selected yeast causing spoilage of
384 canned soft drinks and fruit products In: Skinner FA, Passmore SM, Davenport RR (eds.),
385 Biology and Activities of Yeasts (pp 181-214). Academic Press, London
- 386 4. Stratford M, Hofman PD, Cole MB (2000) Fruit juice, fruit drinks and soft drinks In:
387 Lund BM, Baird-Parker TC, Gould GW (eds.), The Microbiological Safety and Quality
388 of Food (pp 836-869). Aspen publishers, Gaithersberg, MD
- 389 5. Barnett JA, Payne RW, Yarrow D (2000) Yeast Characteristics and identification (3rd
390 ed.) Cambridge University Press, Cambridge
- 391 6. Lima Tribst AA, de Souza Sant'Ana A, de Massaguer PR (2009) Review:
392 Microbiological quality and safety of fruit juices- past, present and future perspectives.
393 Crit Rev Microbiol 35(4):310-339
- 394 7. Deak T, Beuchat LR (1996). Handbook of food spoilage yeasts. CRC press Boca
395 Raton FL
- 396 8. Fleet GH (1992) Spoilage yeasts. Crit rev Biotechnol 12:1-44
- 397 9. Pitt JI, Hocking AD (1997) Fungi and food spoilage, 2nd edn. Aspen Publishers,
398 Gaithersburg, MD
- 399 10. Panagou EZ, Karathanassi S, Le Marc Y, Nychas, GJE (2009) Development of a
400 product specific model for spoilage of pasteurized fruit juices by *Saccharomyces*
401 *cerevisiae* and validation under dynamic temperature conditions. Predictive Modelling in
402 Foods. September 8–12, Washington DC, USA (2009), (CDROM)
- 403 11. Valverde MT, Marín-Iniesta F, Calvo L (2010) Inactivation of *Saccharomyces*
404 *cerevisiae* in conference pear with high pressure carbon dioxide and effects on pear
405 quality. J Food Eng 98:421-428

- 406 12. Arias CR, Burns JK, Friedrich LM, Goodrich RH, Parish ME (2002) Yeast species
407 associated with orange juice: evaluation of identification methods. *Appl Environ*
408 *Microbiol* 68:1955-1961
- 409 13. Deak T, Beuchat LR (1993a) Yeasts associated with fruit juice concentrates. *J Food*
410 *Prot* 56: 777-782
- 411 14. Deak T, Beuchat LR (1993b) Comparison of SIM API 20C and ID32C systems for
412 identification of yeasts from fruit juice concentrates and beverages. *J Food Prot* 56:585-
413 592
- 414 15. Las Heras-Vazquez FJL, Mingorance-Cazorla L, Clemente-Jimenez, JF, Rodriguez-
415 Vico F (2003) Identification of yeast species from orange fruit and juice by RFLP and
416 sequence analysis of the 58S r RNA gene and two internal transcribed spacers. *FEMS*
417 *Yeast Res* 3:3-9
- 418 16. Sancho T, Gimenez-Jurado G, Malfeito-Ferreira M, Loureiro V (2000) Zymological
419 indicators: a new concept applied to the detection of potential spoilage yeast species
420 associated with fruit pulps and concentrates. *Food Microbiol* 17:613-624
- 421 17. Thomas S (1993) Yeasts as spoilage organisms in beverages In: Rose AH, Harrison
422 JS (eds.), *The Yeasts*, volume 5, 2nd edition, (pp 517-561). Academic press, London
- 423 18. Torok T, King AD (1991) Comparative study on the identification of foodborne
424 yeasts. *Appl Environ Microbiol* 57:1207-1212
- 425 19. Stratford M, James SA (2003) Non-alcoholic beverages and yeasts In: Boekhout T,
426 Robert V (eds.), *Yeasts in Food: Beneficial and Detrimental Aspects*, (pp 309-345).
427 Behr's-Verlag, Germany

- 428 20. Fleet GH (2006) *Saccharomyces* and related genera In: Clive de W Blackburn (ed),
429 Food Spoilage Microorganisms (pp. 306-335). CRC press, England
- 430 21. Tribst AAL, Franchi MA, Cristianini M (2008) Ultra-high pressure homogenization
431 treatment combined with lysozyme for controlling *Lactobacillus brevis* contamination in
432 model system. *Innov Food Sci Emerg Technol* 9:265–271
- 433 22. Pathanibul P, Taylor TM, Davidson PM, Harte F (2009) Inactivation of *Escherichia*
434 *coli* and *Listeria innocua* in apple and carrot juices using high pressure homogenization
435 and nisin. *Int J Food Microbiol* 129:316–320
- 436 23. Vachon JF, Kheadr EE, Giasson J, Paquin P, Fliss I (2002) Inactivation of foodborne
437 pathogens in milk using dynamic high pressure. *Journal of Food Prot* 65:345–352
- 438 24. Diels AMJ, Callewaert L, Wuytack EY, Masschalck B, Michiels CW (2005)
439 Inactivation of *Escherichia coli* by high-pressure homogenisation is influenced by fluid
440 viscosity but not by water activity and product composition. *Int J Food Microbiol*
441 101:281–291
- 442 25. US Food and Drug Administration (USFDA) (2004) FDA Guidance to Industry,
443 2004: Recommendations to Processors of Apple Juice or Cider on the Use of Ozone for
444 Pathogen Reduction Purposes. Available online:
445 [http://www.fda.gov/Food/GuidanceComplianceRegulatoryInformation/GuidanceDocume](http://www.fda.gov/Food/GuidanceComplianceRegulatoryInformation/GuidanceDocuments/Juice/ucm072524.htm)
446 [nts/Juice/ucm072524.htm](http://www.fda.gov/Food/GuidanceComplianceRegulatoryInformation/GuidanceDocuments/Juice/ucm072524.htm) (Accessed 5 July 2010)
- 447 26. Khadre MA, Yousef AE, Kim JG (2001) Microbiological aspects of ozone
448 applications in food: A review. *J Food Sci* 66(9):1241-1252

- 449 27. Cullen PJ, Valdramidis VP, Tiwari BK, Patil S, Bourke, P, O'Donnell CP (2010)
450 Ozone Processing For Food Preservation: An Overview on Fruit Juice Treatments.
451 Ozone: Sci Eng 32(3):166-179
- 452 28. Hunt NK, Marinas BJ (1997) Kinetics of *Escherichia coli* inactivation with ozone.
453 Water Res 31:1355-1362
- 454 29. Korycka-Dahl M, Richardson T (1978) Activated oxygen species and oxidation of
455 food constituents. Crit Rev Food Sci Nutr 10:209-241
- 456 30. Patrignani F, Vannini L, Kamdem SLS, Lanciotti R, Guerzoni ME (2009) Effect of
457 high pressure homogenization on *Saccharomyces cerevisiae* inactivation and physico-
458 chemical features in apricot and carrot juices. Int J Food Microbiol 136(1):26-31
- 459 31. Mosqueda-Melgar J, Raybaudi-Massilia RM, Martín-Belloso O (2008) Combination
460 of high-intensity pulsed electric fields with natural antimicrobials to inactivate pathogenic
461 microorganisms and extend the shelf-life of melon and watermelon juices. Food
462 Microbiology 25:479-491
- 463 32. Patil S, Cullen PJ, Kelly B, Frias JM, Bourke P (2009) Extrinsic control parameters
464 for ozone inactivation of *Escherichia coli* using ozone bubble column. J Appl Microbiol
465 107(3):830-837
- 466 33. Patil S, Torres B, Tiwari BK, Wijngaard HH, Bourke P, Cullen PJ, O'Donnell CP,
467 Valdramidis VP (2010) Safety and quality assessment during the ozonation of cloudy
468 apple juice. J Food Sci 75(7):M437-M443
- 469 34. Baranyi J, Roberts TA (1994) A dynamic approach to predicting bacterial growth in
470 food. Int J Food Microbiol 23:277-294

- 471 35. Ratkowsky DA, Olley J, McMeekin TA, Ball A (1982) Relationship between
472 temperature and growth-rate of bacterial cultures. J Bacteriol 149:1-5
- 473 36. Ratkowsky DA, Lowry RK, McMeekin TA, Stokes AN, Chandler RE (1983) Model
474 for bacterial cultures growth rate throughout the entire biokinetic temperature range. J
475 Bacteriol 154:1222-1226
- 476 37. Fu B, Taoukis PS, Labuza TP (1991) Predictive microbiology for monitoring
477 spoilage of dairy products with time-temperature indicators. J Food Sci 56:1209-1215
- 478 38. Le Marc Y, Valik L, Medved'ova A (2009) Modelling the effect of the starting
479 culture on the growth of *Staphylococcus aureus* in milk. Int J Food Microbiol 129:306-
480 311
- 481 39. Neter, Wasserman, Whitmore (1992) Applied statistics, 4th edn. (pp. 1-989). Prentice
482 hall, Englewood cliffs, NJ
- 483 40. Baranyi J, Pin C, Ross T (1999) Validating and comparing predictive models. Int J
484 Food Microbiol 48:159-166
- 485 41. Duyvesteyn WS, Shimoni E, Labuza TP (2001) Determination of end of shelf-life for
486 milk using Weibull hazard method. Lebensm Wiss Technol 34:143-148
- 487 42. Labuza TP (1982) Shelf-life dating of foods pp 54-58, 223, Food and Nutrition Press
488 Inc., Westport, Connecticut
- 489 43. Man D, Jones (2000) Shelf-life evaluation of foods In: Singh RP (ed) Scientific
490 principles of shelf-life evaluation Aspen Publication, Gaithersburg, Maryland
- 491 44. Al-kadamany E, Toufelli I, Khattar M., Abou-Jawdeh Y, Harakeh S, Haddad T
492 (2002) Determination of shelf life of concentrated yogurt (Labneh) produced by in-bag
493 straining of set yogurt using hazard analysis J Dairy Sci 85:1023-1030

- 494 45. Thomas LV, Ingrama RE, Yub S, Delves-Broughtona J (2004) Investigation of the
495 effectiveness of Ascopyrone P as a food preservative. *Int J Food Microbiol* 93:319–323
- 496 46. Brackett RE (1994) Microbiological spoilage and pathogens in minimally processed
497 refrigerated fruits and vegetables. In: Wiley RC (ed), *Minimally Processed Refrigerated*
498 *Fruits and Vegetables* (pp. 269-312). Chapman & Hall, New York, London
- 499 47. Kisko G, Sharp R, Roller S (2005) Chitosan inactivates spoilage yeasts but enhances
500 survival of *Escherichia coli* O157:H7 in apple juice. *J Appl Microbiol* 98: 872-880
- 501 48. Hinze H, Prakash D, Holzer H (1987) Effect of ozone on ATP, cytosolic enzymes and
502 permeability of *Saccharomyces cerevisiae*. *Arch Microbiol* 147:105-108
- 503 49. Ross T, Dalgaard P, Tienungoon S (2000) Predictive modeling of the growth and
504 survival of *Listeria* in fishery products. *Int J Food Microbiol* 62:231-245
- 505 50. Qin BL, Chang FJ, Barbosa-Canovas GV, Swanson BG (1995) Nonthermal
506 inactivation of *Saccharomyces cerevisiae* in apple juice using pulsed electric fields.
507 *Lebensmm Wiss Technol* 28:564-568
- 508 51. Ferrentino G, Bruno M, Ferrari G, Poletto M, Balaban MO (2009) Microbial
509 inactivation and shelf life of apple juice treated with high pressure carbon dioxide. *J Biol*
510 *Eng* 3:3
- 511 52. Suarez-Jacobo A, Gervilla R, Guamis B, Roig-Sagues AX, Saldo J (2009) Effect of
512 UHPH on indigenous microbiota of apple juice: a preliminary study of microbial shelf
513 life. *Int J Food Microbiol* 136(3):261-267
- 514 53. Valdramidis VP, Graham WD, Beattie A, Linton M, McKay A, Fearon AM,
515 Patterson M F (2009) Defining the stability interfaces of apple juice: implications on the

516 optimisation and design of high hydrostatic pressure treatment. *Innov Food Sci Emerg*
517 *Technol* 10(4): 396-404
518 54. Pin C, Sutherland JP, Baranyi J (1999) Validating predictive models of food spoilage
519 organisms. *J Appl Microbiol* 87:491-499
520
521
522