



2008

Clindamycin Resistant clone of *C. difficile* PCR ribotype 027 in Europe

Denise Drudy

Dublin Institute of Technology, denise.drudy@dit.ie

B. Goorhuis

Dennis Bakker

Lorraine Kyne

Renate Van der Berg

See next page for additional authors

Follow this and additional works at: <http://arrow.dit.ie/scschbioart>

Recommended Citation

Drudy, D. (2008) Clindamycin Resistant clone of *C. difficile* PCR ribotype 027 in Europe, *Emerg Infect Dis.*14(9), pp.1485-1487.
doi:0.3201/eid1409.071346

This Article is brought to you for free and open access by the School of Biological Sciences at ARROW@DIT. It has been accepted for inclusion in Articles by an authorized administrator of ARROW@DIT. For more information, please contact yvonne.desmond@dit.ie, arrow.admin@dit.ie, brian.widdis@dit.ie.



This work is licensed under a [Creative Commons Attribution-NonCommercial-Share Alike 3.0 License](https://creativecommons.org/licenses/by-nc-sa/3.0/)



Authors

Denise Drudy, B. Goorhuis, Dennis Bakker, Lorraine Kyne, Renate Van der Berg, Lynda Fenelon, and Seamus Fanning

Mayboun Heuangvongsy, Chanpheng Thammavong, Bouachanh Rasachack, Bounkong Syhavong, Nicholas J. White, Suriyasack Thongpaseuth, Anisone Changthongthip, Viengmone Davong, Olay Lattana, Manivanh Vongsouvath, Kai-amporn Keopaseuth, Sengmani Symanivong, Viengmala Sihalath, and Alatsany Chandara for participating in the study; and Pommek Dalalay and Somphone Phounsavath for support.

This study was supported by the Wellcome Trust–Mahosot Hospital–Oxford Tropical Medicine Research Collaboration, which was supported by the Wellcome Trust of Great Britain.

**Philippe Parola,
Stuart D. Blacksell,
Rattanaphone Phetsouvanh,
Simaly Phongmany,
Jean-Marc Rolain,
Nicholas P.J. Day,
Paul N. Newton,
and Didier Raoult**

Author affiliations: World Health Organization Collaborative Center for Rickettsial Diseases and Other Arthropod Borne Bacterial Diseases, Marseille, France (P. Parola, J.-M. Rolain, D. Raoult); Mahosot Hospital, Vientiane, Laos (S.D. Blacksell, R. Phetsouvanh, S. Phongmany, N.P.J. Day, P.N. Newton); University of Oxford, Oxford, United Kingdom (S.D. Blacksell, N.P.J. Day, P.N. Newton); and Mahidol University, Bangkok, Thailand (S.D. Blacksell, N.P.J. Day)

DOI: 10.3201/eid1409.071259

References

- Phongmany S, Rolain JM, Phetsouvanh R, Blacksell SD, Soukhaseum V, Rasachack B, et al. Rickettsial infections and fever, Vientiane, Laos. *Emerg Infect Dis*. 2006;12:256–62.
- Tamura A, Yamamoto N, Koyama S, Makisaka Y, Takahashi M, Urabe K, et al. Epidemiological survey of *Orientia tsutsugamushi* distribution in field rodents in Saitama Prefecture, Japan, and discovery of a new type. *Microbiol Immunol*. 2001;45:439–46.
- Mahajan SK, Rolain JM, Kashyap R, Bakshi D, Sharma V, Prasher BS, et al. Scrub typhus in Himalayas. *Emerg Infect Dis*. 2006;12:1590–2.
- Jiang J, Chan TC, Temenak JJ, Dasch GA, Ching WM, Richards AL. Development of a quantitative real-time polymerase chain reaction assay specific for *Orientia tsutsugamushi*. *Am J Trop Med Hyg*. 2004;70:351–6.
- Fournier PE, Siritantikorn S, Rolain JM, Suputtamongkol Y, Hoontrakul S, Charoenwat S, et al. Detection of new genotypes of *Orientia tsutsugamushi* infecting humans in Thailand. *Clin Microbiol Infect*. 2008;14:168–73.
- Blacksell SD, Luksameetanasan R, Kallabaheti T, Aukkanit N, Paris DH, McGready R, et al. Genetic typing of the 56-kDa type-specific antigen gene of contemporary *Orientia tsutsugamushi* isolates causing human scrub typhus at two sites in north-eastern and western Thailand. *FEMS Immunol Med Microbiol*. 2008;52:335–42.
- Qiang Y, Tamura A, Urakami H, Makisaka Y, Koyama S, Fukuhara M, et al. Phylogenetic characterization of *Orientia tsutsugamushi* isolated in Taiwan according to the sequence homologies of 56-kDa type-specific antigen genes. *Microbiol Immunol*. 2003;47:577–83.
- Kawamura A, Tanaka H. *Tsutsugamushi* disease: an overview. Tokyo: University of Tokyo Press; 1995.
- Seong SY, Kim MK, Lee SM, Odgerel Z, Choi MS, Kim IS, et al. Neutralization epitopes on the antigenic domain II of the *Orientia tsutsugamushi* 56-kDa protein revealed by monoclonal antibodies. *Vaccine*. 2000;19:2–9. DOI: 10.1016/S0264-410X(00)00167-5
- Suttinont C, Losuwanaluk K, Niwatayakul K, Hoontrakul S, Intaranongpai W, Silpasakorn S, et al. Causes of acute, undifferentiated, febrile illness in rural Thailand: results of a prospective observational study. *Ann Trop Med Parasitol*. 2006;100:363–70. DOI: 10.1179/136485906X112158

Address for correspondence: Didier Raoult, Unité des Rickettsies, Centre National de la Recherche Scientifique–Institut de Recherche pour le Développement, Unité Mixte de Recherche 6236, World Health Organization Collaborative Center for Rickettsioses and Other Arthropod Borne Bacterial Diseases, Faculté de Médecine, 27 Bd Jean Moulin, 13005 Marseille, France; email: didier.raoult@gmail.com

The opinions expressed by authors contributing to this journal do not necessarily reflect the opinions of the Centers for Disease Control and Prevention or the institutions with which the authors are affiliated.

Clindamycin-Resistant Clone of *Clostridium difficile* PCR Ribotype 027, Europe

To the Editor: Since 2003, outbreaks of *Clostridium difficile*–associated disease (CDAD) associated with the emergence of a hypervirulent strain have been reported worldwide (1,2; www.eurosurveillance.org/em/v12n06/1206-221.asp). This strain has been associated with increased disease severity and attributable mortality. Patients infected with *C. difficile* 027 fail to respond to metronidazole therapy (1). Several typing methods have been applied to further characterize *C. difficile* PCR ribotype-027, including pulsed-field gel electrophoresis (PFGE) (North American pulsed field type 1) and restriction enzyme analysis (REA) (BI). PFGE and REA are widely used in the United States; PCR ribotyping is more commonly used throughout Europe. More recently, 2 multiple-locus variable-number tandem-repeat analysis (MLVA) protocols have been applied to type *C. difficile*, and these proved more discriminatory compared to other methods (3,4). Furthermore, MLVA can subgroup geographically diverse 027 isolates (G. Killgore et al., unpub data) as well as 027 isolates that are common to 1 institution (5).

We reported a case of *C. difficile* PCR 027 in Ireland, where the isolate had an identical antibiogram profile compared with those strains reported across Europe (6,7) (i.e., resistant to fluoroquinolones and erythromycin, susceptible to clindamycin). We have subsequently identified *C. difficile* 027 in 6 more healthcare settings. To date >100 Irish *C. difficile* 027 isolates have been characterized by analysis of their antibiogram profiles, toxinotyping, and 16S–23S rDNA PCR ribotyping. All *C. difficile* 027 isolates were resistant to moxifloxacin, gatifloxacin,

ciprofloxacin (MIC >32 mg/L), and erythromycin (MIC >256 mg/L) but susceptible to metronidazole (MIC 0.25 mg/L) and vancomycin (MIC >0.5 mg/L). Clindamycin susceptibility varied between isolates from unrelated institutions. Isolates from 2 healthcare settings were susceptible to clindamycin (n = 11; MIC₉₀ 4 mg/L). However, clindamycin-resistant PCR 027 isolates (n = 96; MIC₉₀ >256 mg/L) were identified in the other 5 healthcare institutions. All clindamycin-resistant PCR 027 isolates were positive for the *ermB* gene, encoding the macrolide-lincosamide-streptogramin-B genotype.

A subset of clindamycin-sensitive and -resistant Irish 027 strains isolated throughout 2006 (n = 22) were further characterized by using a recently described MLVA protocol (3). Six clindamycin-susceptible isolates were selected from 2 healthcare settings. One hospital conducted active routine laboratory surveillance and molecular genotyping (n = 3). The second hospital submitted only random isolates (n = 3) for typing during a *C. difficile* outbreak. Sixteen clindamycin-resistant PCR 027 isolates were also included in the MLVA. Resistant isolates were selected from 5 healthcare settings. These included isolates from 2 *C. difficile* outbreaks with ongoing laboratory surveillance (n = 5, n = 6, respectively); a third hospital with ongoing laboratory surveillance (n = 3) and 2 hospitals that each submitted fecal samples from patients with severe cases of *C. difficile* disease (n = 1). The Stoke-Mandeville control strain R20291 was included for comparison.

MLVA determined that all strains within the clindamycin-resistant cluster were closely related and were single- or double-locus variants with a maximum 5 summed tandem-repeat difference (STRD). In contrast, the closest relationship between the clindamycin-resistant and the clindamycin-sensitive clusters was a triple-locus variant with an STRD of 17.

The nonrelated reference strain of the Stoke-Mandeville outbreak (R20291) differed considerably from all Irish isolates but was more related to the clindamycin-sensitive cluster than to the clindamycin-resistant cluster (Figure). We thus linked a defined genetic marker with the clindamycin-resistant phenotype in *C. difficile* PCR-027. MLVA could clearly differentiate clindamycin-resistant and -susceptible isolates from the same geographic region and subgrouped them into 2 distinct clusters (Figure).

Although high-level resistance to fluoroquinolone antimicrobial agents has been well documented in PCR 027 (1,6), resistance to clindamycin is rare. Subsequently, clindamycin has been considered as a “protective” antimicrobial agent for the development of CDAD in an epidemiologic survey in the Netherlands (8). Currently, resistance to this agent in NAP 1/PCR 027 has been restricted to the United States. McDonald and colleagues reported that 19 (79%) of 24 NAP 1 isolates were

classified as less susceptible (MIC 4 mg/L) or resistant (MIC 8 mg/L) to clindamycin when Clinical and Laboratory Standards Institute criteria were used (2). Unfortunately, MIC values were not reported, and the corresponding resistance genes were not investigated. In contrast, Canadian studies to date have not reported clindamycin resistance in this strain type. The MIC₉₀ of Canadian NAP 1 isolates for clindamycin was 4 mg/L (9,10). Although outbreaks and sporadic cases of PCR 027 have been identified in several European countries, to date no clindamycin-resistant clone has been reported.

Detection of clindamycin-resistant *C. difficile* PCR 027 strains is an important and worrying development. Resistance to this antimicrobial agent increases the risk for CDAD in patients, and its use may be an important factor contributing to the persistence and spread of PCR 027. A similar feature has already been observed when fluoroquinolones and cephalosporins are prescribed. Clindamycin-resistant PCR




Figure. Minimal spanning tree of 23 *Clostridium difficile* isolates. In the circles, the individual isolates are mentioned. The numbers between the circles represent the summed tandem repeat differences (STRDs) between multiple-locus variable-number tandem-repeat analysis types. Straight lines represent single-locus variants, dashed lines double-locus variants. Curved lines represent triple-locus variants. Two related clusters can be discriminated: the light gray cluster (isolates B1, B4, M246, B6, and M216) and the cluster within dotted lines (isolates V6-44, V6-142, V6-81, 1ML, C1, 4108, V6-35, V6-80, L1, 2191cc, C4, C8, 3ML, C44, C37, and 13ML). The isolates in the light gray cluster are sensitive to clindamycin; isolates in the cluster surrounded by dotted lines are resistant. Two isolates (M278 and R20291) did not belong to a cluster but were more related to the sensitive cluster than to the resistant cluster. Genetically related clusters were defined by an STRD ≤10.

027 probably reflects the emergence of a new clone because MLVA clearly differentiates between clindamycin-susceptible and -resistant isolates.

**Denise Drudy, Bram Goorhuis,
Dennis Bakker, Lorraine Kyne,
Renate van den Berg,
Lynda Fenelon,
Seamus Fanning,
and Edward J. Kuijper**

Author affiliations: University College Dublin, Dublin, Ireland (D. Drudy, L. Kyne, L. Fenelon, S. Fanning); Leiden University Medical Center, Leiden, the Netherlands (B. Goorhuis, D. Bakker, R. van den Berg, E.J. Kuijper); and European Centre for Disease Prevention and Control, Stockholm, Sweden (E.J. Kuijper)

DOI: 10.3201/eid1409.071346

References

- Kuijper EJ, Coignard B, Tull P. the ESCMID Study Group for *Clostridium difficile* (ESGCD)*; EU Member States and the European Centre for Disease Prevention and Control (ECDC). Emergence of *Clostridium difficile*-associated disease in North America and Europe. *Clin Microbiol Infect*. 2006;12:2–18. DOI: 10.1111/j.1469-0691.2006.01580.x
- McDonald LC, Killgore GE, Thompson A, Owens RC Jr, Kozakova SV, Sambol SP, et al. An epidemic, toxin gene-variant strain of *Clostridium difficile*. *N Engl J Med*. 2005;353:2433–41. DOI: 10.1056/NEJMoa051590
- van den Berg RJ, Schaap I, Templeton KE, Klaassen CH, Kuijper EJ. Typing and subtyping of *Clostridium difficile* isolates by using multiple-locus variable-number tandem-repeat analysis. *J Clin Microbiol*. 2007;45:1024–8. DOI: 10.1128/JCM.02023-06
- Marsh JW, O'Leary MM, Shutt KA, Pascualle AW, Johnson S, Gerding DN, et al. Multilocus variable-number tandem-repeat analysis for investigation of *Clostridium difficile* transmission in hospitals. *J Clin Microbiol*. 2006;44:2558–66. DOI: 10.1128/JCM.02364-05
- Fawley WN, Freeman J, Smith C, Harmanus C, van den Berg RJ, Kuijper EJ, et al. Use of highly discriminatory fingerprinting to analyze clusters of *Clostridium difficile* infection cases due to epidemic ribotype 027 strains. *J Clin Microbiol*. 2008;46:954–60. DOI: 10.1128/JCM.01764-07
- Long S, Fenelon L, Fitzgerald S, Nolan N, Burns K, Hannan M, et al. First isolation and report of clusters of *Clostridium difficile* PCR 027 cases in Ireland. *Eurosurveillance* 2007;12:E070426.3.
- Drudy D, Kyne L, O'Mahony R, Fanning S. *GyrA* mutations in fluoroquinolone-resistant *Clostridium difficile* PCR-027. *Emerg Infect Dis*. 2007;13:504–5.
- Goorhuis A, Van der Kooij T, Vaessen N, Dekker FW, Van den Berg R, Harmanus C, et al. Spread and epidemiology of *Clostridium difficile* polymerase chain reaction ribotype 027/toxinotype III in The Netherlands. *Clin Infect Dis*. 2007;45:695–703. DOI: 10.1086/520984
- Bourgault AM, Lamothe F, Loo VG, Poirier L; CDAD-CSI Study Group. In vitro susceptibility of *Clostridium difficile* clinical isolates from a multi-institutional outbreak in Southern Québec, Canada. *Antimicrob Agents Chemother*. 2006;50:3473–5. DOI: 10.1128/AAC.00479-06
- MacCannell DR, Louie TJ, Gregson DB, Laverdiere M, Labbe AC, Laing F, et al. Molecular analysis of *Clostridium difficile* PCR ribotype 027 isolates from Eastern and Western Canada. *J Clin Microbiol*. 2006;44:2147–52. DOI: 10.1128/JCM.02563-05

Address for correspondence: Denise Drudy, Centre for Food Safety, Veterinary Sciences Centre, University College Dublin, Belfield, Dublin 4, Ireland; email: denise.drudy@ucd.ie

Letters

Letters commenting on recent articles as well as letters reporting cases, outbreaks, or original research are welcome. Letters commenting on articles should contain no more than 300 words and 5 references; they are more likely to be published if submitted within 4 weeks of the original article's publication. Letters reporting cases, outbreaks, or original research should contain no more than 800 words and 10 references. They may have one Figure or Table and should not be divided into sections. All letters should contain material not previously published and include a word count.

Increasing Incidence of *Clostridium difficile*-associated Disease, Singapore

To the Editor: *Clostridium difficile*-associated disease (CDAD) has increased in incidence across North America and Europe (1). Recent reports document the emergence of an epidemic strain of *C. difficile*, NAP1/BI/027, associated with increased virulence (2,3). However, less information is available regarding CDAD epidemiology in Asia. We examined the incidence of *C. difficile* among hospitalized patients in Singapore from 2001 through 2006 and conducted a case-control study to evaluate risk factors for testing positive for *C. difficile* toxin (CDT) in our population.

Tan Tock Seng Hospital (TTSH) is a 1,200-bed, acute-care general hospital in Singapore that serves an urban population of 4 million. We calculated CDAD incidence using the number of patients testing positive for CDT per 10,000 patient days from 2001 through 2006. We used this calculation because CDT testing would have been ordered for clinical indications. CDT testing was performed by using the same ELISA (Premier Toxins A&B; Meridian Bioscience, Inc., Cincinnati, OH, USA) throughout the entire period of investigation.

Case-patients and controls were selected from patients hospitalized at TTSH from January 1 through December 31, 2004. Microbiology laboratory records were used to define 3 groups. Case-patients were defined as CDT-positive inpatients (group 1). Two sets of negative controls were defined: the first (group 2) consisted of patients who tested negative for CDT. However, because false-negatives could nullify differences between groups 1 and 2, we defined a second set of negative controls (group 3) from among 18,000 inpatients not tested for CDT.