



2007

# Development and assessment of a rapid method to detect *Escherichia coli* O26, O111 and O157 in retail minced beef.

Mary Murphy

*Health Service Executive-South Western area, Cherry Orchard Hospital, Ballyfermot, Dublin 10., Centre for Food Safety*

Anne Carroll

*Public Health Laboratory, Dublin.*

Paul Whyte

*Health Service Executive-South Western area, Cherry Orchard Hospital, Ballyfermot, Dublin 10., Centre for Food Safety.*

Micheál O'Mahoney

*Health Service Executive-South Western area, Cherry Orchard Hospital, Ballyfermot, Dublin 10., Centre for Food Safety*

Wayne Anderson

*University College Dublin, Ireland*

*See next page for additional authors*

Follow this and additional works at: <http://arrow.dit.ie/schfsehart>



Part of the [Food Science Commons](#)

## Recommended Citation

Murphy, M., Walsh, C. et al (2007) Development and assessment of a rapid method to detect *Escherichia coli* O26, O111 and O157 in retail minced beef. *International Journal of Hygiene and Environmental Health*, 210, 155-161, 2007.. doi: 10.1016/j.ijheh.2006.09.004

This Article is brought to you for free and open access by the School of Food Science and Environmental Health at ARROW@DIT. It has been accepted for inclusion in Articles by an authorized administrator of ARROW@DIT. For more information, please contact [yvonne.desmond@dit.ie](mailto:yvonne.desmond@dit.ie), [arrow.admin@dit.ie](mailto:arrow.admin@dit.ie), [brian.widdis@dit.ie](mailto:brian.widdis@dit.ie).



This work is licensed under a [Creative Commons Attribution-NonCommercial-Share Alike 3.0 License](#)



---

**Authors**

Mary Murphy, Anne Carroll, Paul Whyte, Micheál O'Mahoney, Wayne Anderson, Eleanor McNamara, Seamus Fanning, and Ciara Walsh

1 **Version 1**

2

3 **Development and assessment of a rapid method to detect *Escherichia***  
4 ***coli* O26, O111 and O157 in retail minced beef.**

5

6

7 Mary Murphy<sup>1, 2</sup>, Anne Carroll<sup>1</sup>, Paul Whyte<sup>2</sup>, Micheál O' Mahony<sup>2, 3</sup>, Wayne  
8 Anderson<sup>3</sup>, Eleanor McNamara<sup>1</sup> and Séamus Fanning<sup>2</sup>.

9

10

11

12

13 Public Health Laboratory<sup>1</sup>, Health Service Executive-South Western area,  
14 Cherry Orchard Hospital, Ballyfermot, Dublin 10., Centre for Food Safety<sup>2</sup>,  
15 School of Agriculture, Food Science and Veterinary Medicine, University  
16 College, Belfield, Dublin 4., Food Safety Authority of Ireland<sup>3</sup>, Abbey Court,  
17 Lower Abbey Street, Dublin 1, Ireland.

18

19

20

21

22 **Corresponding Author:** Professor Séamus Fanning,  
23 Centre for Food Safety,  
24 School of Agriculture, Food Science and  
25 Veterinary Medicine,  
26 University College,  
27 Belfield, Dublin 4,  
28 Ireland.

29 Tel. (+353-1) 716 6082,

30 Fax (+353-1) 716 6091.

31 [sfanning@ucd.ie](mailto:sfanning@ucd.ie)

32 **ABSTRACT**

33 A molecular-based detection method was developed to detect *Escherichia coli*  
34 O26, O111 and O157 in minced (ground) beef samples. This method consists of  
35 an initial overnight enrichment in modified tryptone soya broth (mTSB) and  
36 novobiocin prior to DNA extraction and subsequent serogrouping using a triplex  
37 PCR. This method has a low limit of detection and results are available within 24  
38 hours of receipt of samples. Once optimized, this rapid method was utilized to  
39 determine the prevalence of these *E. coli* serogroups in six hundred minced beef  
40 samples all of which were previously examined by immunomagnetic separation  
41 (IMS) and selective plating for *E. coli* O26 and O111. Using IMS, two *E. coli* O26  
42 isolates were detected. No *E. coli* O111 were recovered. The multiplex PCR  
43 technique described here did not detect *E. coli* O111 nor O157 in any of the  
44 samples, however six minced beef samples were positive for *E. coli* O26 using  
45 our method, only two of these were previously detected by IMS and culture.  
46 Application of molecular methods are useful to support culture-based approaches  
47 thereby further contributing to risk reduction along the food chain.

48

49

50

51

52

53 **Keywords:** non-O157 *E. coli*; *E. coli* O157, O26 and O111; serogroup-specific  
54 PCR; rapid detection in food.

55 **INTRODUCTION**

56 *Escherichia coli* O157 was initially implicated in human disease in 1982 (Riley et  
57 al., 1983) and since then the reported incidence of verocytotoxigenic *Escherichia*  
58 *coli* (VTEC) related illness has increased worldwide. In addition to *Escherichia*  
59 *coli* (*E. coli*) O157, other serogroups of *E. coli* such as O111 and O26 have  
60 emerged as important etiological agents of gastrointestinal disease in humans  
61 (Tarr and Neill, 1996). There is evidence of these infections in Ireland including  
62 an outbreak of *E. coli* O26 which was reported in a crèche in 1999 (McMaster et  
63 al., 2001). However, the true incidence and clinical significance of non-O157  
64 serogroups in Ireland still remains to be determined. Routine surveillance to  
65 detect these non-O157 *E. coli* in foods, animals and the environment need to be  
66 introduced so that public health measures can be implemented. Development of  
67 rapid diagnostics capable of detecting a wide range of *E. coli* serogroups  
68 associated with disease in humans is a priority.

69

70 In recent years, a great deal of work has been carried out to ascertain the most  
71 sensitive and specific methods for isolating and identifying *E. coli* O157. The  
72 current method of choice for the detection of *E. coli* O157 in foods is enrichment  
73 followed by concentration using Immunomagnetic separation (IMS), with  
74 subsequent plating onto the selective Cefixime-Tellurite Sorbitol MacConkey  
75 agar (CT-SMAC). This procedure was adapted to allow for the detection of *E.*  
76 *coli* O26 and O111 in foods with the use of serogroup-specific beads. However, a

77 major obstacle in detecting *E. coli* O26 and *E. coli* O111 by this method is the  
78 lack of suitable selective agars.

79

80 CT-SMAC is the selective agar of choice used to detect *E. coli* O157 from many  
81 food matrices as the majority of *E. coli* O157 are non-sorbitol fermenting.

82 However sorbitol fermenting *E. coli* O157 strains capable of causing haemolytic  
83 uremic syndrome (HUS) have been reported (Bettleheim et al., 2002; Karch and

84 Bielaszewska, 2001; Karch et al., 1993). While *E. coli* O26 and O111 have

85 demonstrated abilities to ferment both sorbitol and lactose (Batson et al., 2002;

86 Hiramatsu et al., 2002), non-sorbitol fermenting *E. coli* O26 and O111 have also

87 been isolated (Batson et al., 2002; Hiramatsu et al., 2002; Louie et al., 1998).

88 Batson et al. (2002) reported on the carbohydrate fermenting ability of forty-one

89 isolates of *E. coli* O26 and twenty-two isolates of *E. coli* O111. These authors

90 reported that all isolates could ferment lactose, 92.5% of the *E. coli* O26 isolates

91 and only 64.7% of the *E. coli* O111 fermented sorbitol.

92

93 As sugar fermenting characteristics are not consistent, it is difficult to choose (or

94 formulate) a selective agar for the detection of all of these serogroups in food

95 matrices containing high numbers of background microflora. The situation has

96 improved somewhat, with the introduction of Rhamnose MacConkey agar

97 (Hiramatsu et al., 2002) as a selective media for *E. coli* O26 and with the

98 development of other chromogenic agars. However, laboratories remain over-

99 reliant on the fermentation characteristics of these micro-organisms which could

100 result in pathogenic isolates going undetected. Taking this into consideration  
101 along with the fact that culture methods are time-consuming, a PCR based  
102 method could be valuable in providing additional information to identify specific  
103 *E. coli* serogroups.

104

105 In this paper, we report on the development of a triplex PCR method to detect *E.*  
106 *coli* O26, O111 and O157 in a single tube reaction following enrichment. Our  
107 method was applied retrospectively to analyze 600 minced beef samples for the  
108 three serogroups. This food matrix was previously investigated by microbiological  
109 methods (Murphy et al., 2005), and in this case failed to identify a number of  
110 positive samples. Application of our method is discussed as a convenient tool to  
111 support risk reduction measures along the food chain.

112

113

114

115

116

117

118

119

120

121

122

123 **MATERIALS & METHODS**

124 *Preparation of bacterial strains for inoculum.*

125 A single colony of *E. coli* O26 (NCTC 8781) or *E. coli* O111 (NCTC 9111) or  
126 *E. coli* O157 (NCTC 12900), were inoculated into three separate falcon test-  
127 tubes containing 4 ml of nutrient broth (Oxoid, Basingstoke, UK) and incubated at  
128 37°C for 4 h. Serial dilution of 4 h cultures were prepared in 9 ml Maximum  
129 Recovery Diluent (MRD; Oxoid) down to a dilution of 10<sup>-9</sup>. One hundred micro-  
130 liters of 10<sup>-5</sup>, 10<sup>-6</sup>, 10<sup>-7</sup>, 10<sup>-8</sup> and 10<sup>-9</sup> serial strain dilutions were plated directly  
131 onto nutrient agar (Oxoid) and incubated at 42°C for 16 h to determine the cell  
132 numbers in the initial inoculum. All inoculations were carried out in duplicate.

133

134 *Inoculation of minced beef samples to determine limit of detection.*

135 Minced beef was purchased at a local supermarket and stored at 4°C prior to  
136 inoculation. One ml of the 10<sup>-7</sup>, 10<sup>-8</sup> and 10<sup>-9</sup> serial dilutions of each strain were  
137 added to separate 25 g minced (ground) beef samples containing 225 ml of  
138 modified Tryptone Soya Broth (mTSB; Oxoid) with novobiocin (20 mg/L), (LabM  
139 Ltd., Lancashire, UK) and incubated at 42°C overnight (16 h). An uninoculated  
140 sample of minced beef was also included as a control.

141

142 *DNA Extraction*

143 DNA was extracted from the overnight (16 h) enrichments both immediately and  
144 after storage for approximately 7 days at -20°C, using the Dneasy tissue kit  
145 (Qiagen, Hilden, Germany) according to manufacturer's instructions with minor



146 modifications. Briefly, 25 µl of each sample were added to eppendorf tubes  
147 containing 180 µl of the kit tissue lysis buffer (ATL). Twenty micro-liters of  
148 proteinase K (supplied in the kit) was added to each tube and mixed. All samples  
149 were heated for approximately 135 min at 55°C and were mixed occasionally  
150 during incubation to maintain homogeneity. Following incubation, samples were  
151 vortexed for 15 seconds and the Protocol for 'Isolation of total DNA from animal  
152 tissues', was followed from step 3, of the Dneasy tissue kit instructions (Qiagen).

153

#### 154 *Triplex-PCR mediated serogrouping*

155 All spiked samples were examined for the presence of *E. coli* O26, O111 and  
156 O157 using the triplex PCR. This triplex reaction was designed by combining  
157 previously published primers by Paton and Paton (1998) for the *rfb* genes of *E.*  
158 *coli* O157 and O111 and primers published by Debroy et al. (2004) for *E. coli*  
159 O26 *wzx* gene. All primers were tested against several bacterial species other  
160 than *E. coli* and also against different serogroups of *E. coli* (Table 1).

161

#### 162 *Triplex-PCR*

163 Samples (5 µl) of each purified template DNA [approx. 100 ng] were amplified in  
164 50 µl reaction mixtures containing 5 µl of dNTP mixture (consisting of 3 mM each  
165 of dATP, dCTP, dGTP, and dTTP), 2 µl of forward and reverse primer mix (5  
166 pmol/µl each primer—see Table 2), 5 µl of 10 x buffer (750 mM Tris-HCL, pH 8.8,  
167 200 mM (NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub>, 0.1% (v/v) Tween 20), 5 mM MgCl<sub>2</sub> and 2.5 U Red Hot *Taq*  
168 DNA Polymerase (Abgene, Surrey, U.K.). Thermal cycling consisted of 36 cycles,

169 each of 95°C for 30 sec, 60°C for 60 sec and 72°C for 60 sec, and a final  
170 extension at 72°C for 10 min. Each amplification reaction included a positive  
171 control and a no-DNA template control. Amplified DNA products were resolved  
172 by 2% (w/v) agarose gel electrophoresis in 1 x Tris-Borate-EDTA (TBE) buffer  
173 containing 0.5 mg/ml ethidium bromide at 100 V and the results visualized and  
174 photographed using the Gel Doc 2000 system (Bio-Rad, Hercules, CA, U.S.A.).

175

176 *Purification and pooling of DNA templates.*

177 In a previous study, six hundred retail minced beef samples (25 g each), were  
178 examined by IMS and culture for the presence of *E. coli* O26 and O111 (samples  
179 were not examined for *E. coli* O157 (Murphy et al., 2005)). Two ml aliquots of  
180 each beef sample were also frozen at -70°C after an overnight enrichment (for 16  
181 h) in mTSB and novobiocin. Each frozen enrichment broth was thawed rapidly at  
182 50°C (Ternent et al., 2004) and 1 ml aliquots were pooled into lots containing five  
183 representative meat samples. DNA was purified from each pool using the DNA  
184 extraction protocol previously outlined and the recovered DNA pools were then  
185 subjected to the triplex PCR. Individual samples from any positive pool were  
186 subsequently tested separately, to identify the positive sample(s). Further work to  
187 determine the virulence status of all positive samples was undertaken using real-  
188 time PCR for *vt1*, *vt2*, *eae* and *hlyA*.

189

190

191

192 *'Real-Time' PCR for detection of vt1, vt2, eae and hlyA-encoding genes*  
193 Five micro-liters of purified template genomic DNA were added to the following  
194 25 µl PCR mix containing 1 X SYBR-1 green PCR master mix (Bio-Rad  
195 Laboratories) and 50 ng of each primer to target the *vt1*, *vt2* (Gannon et al.,  
196 1992), *eae* (Beebakhee et al., 1992) and *hlyA* genes (Paton and Paton, 1998).  
197 Samples were amplified in an I-Cycler (Bio-Rad, Hercules, CA, U.S.A), at 95°C  
198 for 2 min followed by 30 cycles each of 45 sec at 95°C, 45 sec at 58°C, and 60  
199 sec at 72°C and a final extension step of 72°C for 10 minutes. Amplification was  
200 monitored by the accumulation of SYBR-1 Green as the amplification reaction  
201 progressed and data captured was later analyzed to produce the corresponding  
202 melt curve.

203

## 204 **RESULTS**

205 In this study, a triplex PCR was developed to detect *E. coli* O26, O111 and O157  
206 in a single reaction. The specificity of the PCR was tested using a panel of  
207 micro-organisms (Table 1). This triplex PCR method was used in conjunction  
208 with the Dneasy animal tissue kit to detect *E. coli* O26, O111 and O157 from  
209 minced (ground) beef, after an initial 16 h enrichment. The limit of detection for  
210 the assay was determined to be 10 colony forming units (CFU) for *E. coli* O111,  
211 15 CFU for *E. coli* O26, <10 CFU for *E. coli* O157 in 25 g of minced beef (Fig.  
212 1a). The uninoculated minced beef sample was determined to be negative for *E.*  
213 *coli* O157, O26 and O111 by this method. Results were available in less than 24  
214 h. No difference in the limit of detection was observed between spiked broths

215 examined directly after enrichment and the enrichment broths that were frozen  
216 for a period before testing (data not shown). Enrichment time trials were also  
217 undertaken, that included times of 4 h, 6 h and overnight (16 h) enrichments.  
218 Data showed that an overnight enrichment (16 h) was required to obtain the low  
219 limits of detection described above (data not shown). These data correlated with  
220 a recent study by O'Hanlon et al. (2004) wherein these authors showed that a  
221 longer enrichment time allowed for the multiplication of slower growing  
222 serogroups, thereby reducing the possibility of false negative results being  
223 reported.

224

225 Compared to IMS and culture (Murphy et al., 2005) this molecular approach  
226 identified six *E. coli* O26 isolates, only two of which were previously detected. No  
227 *E. coli* O157 nor *E. coli* O111 were detected using this rapid method. The latter  
228 case correlates with the IMS and culture results where *E. coli* O111 was not  
229 detected (Murphy et al., 2005). All six *E. coli* O26 isolates were tested for the  
230 presence of the virulence factors *vt1*, *vt2*, *eae* and *hlyA* by real-time PCR. Table  
231 3 showed that isolate-4 (Fig. 1b, 1c) contained both *vt2* and *hlyA* and isolate-6  
232 contained the *eae* gene alone (Fig. 1c and Table 3). The remaining four isolates  
233 did not contain any virulence factors as determined by this method (Table 3).

234

235

236

237 **DISCUSSION**

238 A study, was recently carried out on the 'Prevalence and characterization of  
239 *Escherichia coli* O26 and O111 in retail minced beef in Ireland' (Murphy et al.,  
240 2005), which highlighted the need for an alternative method to detect these  
241 micro-organisms in food. IMS and culture is the technique routinely used  
242 worldwide for the detection and identification of *E. coli* O157 in food matrices.  
243 However, this technique is time-consuming and is sub-optimal when used for the  
244 detection of other *E. coli* serogroups such as *E. coli* O26 and O111, increasing  
245 the risk of reporting false negative results.

246

247 In our study, a triplex PCR was developed to detect the three *E. coli* serogroups  
248 in minced beef after an initial overnight enrichment step (16 h) followed by DNA  
249 purification. The presence of these micro-organisms in minced beef was  
250 determined within twenty-four hours, reducing the time required when compared  
251 to standard methods by up to four days. Speed of detection is a very important  
252 factor in outbreak scenarios and routine food surveillance. Our method could  
253 detect less than 15 CFU of all the serogroups tested, which is comparable to the  
254 limit of detection for the IMS and culture technique (Safarikova et al., 2001) and  
255 to data previously reported from studies using real-time PCR (O'Hanlon et al.,  
256 2004; Sharma et al., 2002). While, the molecular method described in this paper  
257 compliments the methodologies of O'Hanlon et al. (2004) and Sharma et al.  
258 (2002), a useful feature is its capability to detect all three serogroups in a single  
259 triplex reaction whether virulent or non-virulent. The additional real-time PCR

260 allows for further characterization of any positive samples for *vt1*, *vt2*, *eae* and  
261 *hlyA* genes.

262

263 Using the protocol described, serogrouping results are available within eight  
264 hours after enrichment therefore, if these broths were stored at 4°C employing  
265 suitable biosecurity measures, positive samples can be cultured, if required. A  
266 unique finding of our study relates to large longitudinal surveillance studies. In  
267 this case culture enrichments may be frozen and tested at a later stage in bulk or  
268 in pools, a point which would contribute positively to reducing the financial  
269 burden associated with these studies.

270

271 Analysis of the six hundred frozen enrichment broths demonstrated the ability of  
272 this method to detect target sequences in samples negative for IMS and culture.  
273 As these isolates were detected among high numbers of background micro-flora  
274 in enriched minced beef, it would be reasonable to assume that all these isolates  
275 were viable, an important point when applying PCR methods. As one of the *E.*  
276 *coli* O26 isolates detected using the rapid method contained *vt2* and *hlyA* genes  
277 and the other isolate contained the *eae* gene alone, these organisms could be of  
278 potential public health significance, should they be ingested by a vulnerable  
279 individual. This observation demonstrates the advantage of combining the  
280 approach described here along with conventional approaches, emphasizing the  
281 value of molecular methods.

282 Implementation of the method outlined in this study would be of value in food  
283 surveillance and public health laboratories. Results of analysis are available in  
284 less than twenty-four hours. It has a low limit of detection and it is less labour  
285 intensive compared to the more traditional methods applied. Our method could  
286 be extended to include a wider range of serogroups or other food-borne  
287 pathogens of public health significance.

288

289

## 290 **ACKNOWLEDGMENTS**

291 We wish to thank the laboratory and research staff in both the Public Health  
292 Laboratory, Cherry Orchard Hospital and the Centre for Food Safety, at  
293 U.C.D., for their valued assistance throughout the project. We also wish to  
294 acknowledge the financial support provided by the Food Safety Authority of  
295 Ireland (FSAI) for part-funding this project.

296

297

298

299

300

301

302

303

304

305 **References**

306 Batson, S.D., Murinda, S.E., Nguyen, L.T., Oliver, S.P., 2002. Phenotypic and  
307 genetic characterization of *Escherichia coli* O26 isolates from North America  
308 using carbohydrate fermentation and genes encoding virulence factors. [On-Line]  
309 <[http://animalscience.ag.utk.edu/annual\\_reports\\_all.htm](http://animalscience.ag.utk.edu/annual_reports_all.htm)>

310

311 Beebakhee, G., Louie, M., De Azavedo, J., Brunton, J., 1992. Cloning and  
312 nucleotide sequence of the *eae* gene homologue from enterohemorrhagic  
313 *Escherichia coli* serotype O157:H7. FEMS Microbiol. Lett. 70, 63-68.

314

315 Bettleheim, K.A., Whipp, M., Djordjevic, S.P., Ramachandran, V., 2002. First  
316 isolation outside Europe of sorbitol-fermenting verocytotoxigenic *Escherichia coli*  
317 (VTEC) belonging to O group O157. J. Med. Microbiol. 51, 713-714.

318

319 DebRoy, C., Roberts, E., Kundrat, J., Davis, M.A., Briggs, C.E., Fratamico, P.M.,  
320 2004. Detection of *Escherichia coli* serogroups O26 and O113 by PCR  
321 amplification of the *wzx* and *wzy* genes. Appl. Environ. Microbiol. 70, 1830-1832.

322

323 Hiramatsu, R., Matsumoto, M., Miwa, Y., Suzuki, Y., Saito, M., Miyazaki, Y.,  
324 2002. Characterization of Shiga Toxin-Producing *Escherichia coli* O26 Strains  
325 and Establishment of Selective Isolation Media for These Strains. J. Clin.  
326 Microbiol. 40, 922-925.

327



328 Gannon, V.P., King, R.K., Kim, J.Y., Thomas, E.J., 1992. Rapid and sensitive  
329 method for detection of Shiga-like toxin-producing *Escherichia coli* in ground beef  
330 using the polymerase chain reaction. Appl. Environ. Microbiol. 58, 3809-15.  
331

332 Karch, H., Bohm, H., Schmidt, H., Gunzer, F., Aleksic, S., Heesemann, J., 1993.  
333 Clonal structure and pathogenicity of shiga-like toxin-producing, sorbitol-  
334 fermenting *Escherichia coli* O157:H-. J. Clin. Microbiol. 31, 1200-1205.  
335

336 Karch, H., Bielaszewska, M., 2001. Sorbitol-fermenting shiga-toxin-producing  
337 *Escherichia coli* O157:H- strains: Epidemiology, phenotypic and molecular  
338 characteristics and microbiological diagnosis. J. Clin. Microbiol. 39, 2043-2049.  
339

340 Louie, M., Read, S., Simor, A.E., Holland, J., Louie, L., Ziebell, K., Brunton, J.,  
341 Hii, J., 1998. Application of Multiplex PCR for Detection of Non-O157  
342 Verocytotoxin-producing *E. coli* in bloody stools: Identification of Serogroups O26  
343 and O111. J. Clin. Microbiol. 36, 3375-3377.  
344

345 McMaster, C., Roch, E.A., Willshaw, G.A., Doherty, A., Kinnear, W., Cheasty, T.,  
346 2001. Verocytotoxin-producing *Escherichia coli* serotype O26:H11 Outbreak in  
347 an Irish crèche. Eur. J. Clin. Microbiol. Infect. Dis. 20, 430-432.  
348

349 Murphy, M., Carroll, A., Whyte, P., Mahony, M., Anderson, W., McNamara, E.,  
350 Fanning, S., 2005. Prevalence and Characterization of *Escherichia coli* O26

351 and O111 in Retail Minced Beef in Ireland. Foodborne Pathogens and  
352 Disease. 2, 357-360.

353

354 O'Hanlon, K.A., Catarama, T.M.G., Duffy, G., Blair, I.S., McDowell, D.A., 2004.  
355 Rapid detection and quantification of *E. coli* O157/O26/O111 in minced beef by  
356 real-time PCR. J. Appl. Microbiol. 96, 1013-1023.

357

358 Paton, A.W., Paton, J.C., 1998. Detection and Characterization of Shiga  
359 Toxigenic *Escherichia coli* by using Multiplex PCR Assays for *stx*<sub>1</sub>, *stx*<sub>2</sub>, *eaeA*,  
360 Enterohemorrhagic *E. coli hlyA*, *rfb*<sub>O111</sub>, and *rfb*<sub>O157</sub>. J. Clin. Microbiol. 36, 598-  
361 602.

362

363 Riley, L.W., Remis, R.S., Helgerson, S.D., McGee, H.B., Wells, J.G., Davis B.R.,  
364 Hebert, R.J., Olcott, E.S., Johnson, L.M., Hargrett, N.T., Blake, P.A., Cohen,  
365 M.L., 1983. Hemorrhagic colitis associated with a rare *Escherichia coli* serotype.  
366 N. Engl. J. Med. 308, 681-685.

367

368 Safarikova, M., Safarik, I., 2001. Immunomagnetic separation of *Escherichia coli*  
369 O26, O111 and O157 from vegetables. Lett. in Appl. Microbiol. 33, 36-39.

370

371 Sharma, V.K., 2002. Detection and quantitation of enterohemorrhagic  
372 *Escherichia coli* O157, O111 and O26 in beef and bovine feces by real-time  
373 polymerase chain reaction. J. Food Protect. 65, 1371-1380.

374

375 Tarr, P.I., Neill, M.A., 1996. Perspective: the problem of non-O157 Shiga toxin  
376 (verocytotoxin)-producing *Escherichia coli*. J. Infect. Dis. 174, 1136-1139.

377

378 Ternent, H.E., Innocent, G.T., Filshie, L.M., Taylor, D.J., Steele, W.B., McEwen,  
379 S.A., Reilly, W.J., Gunn, G.J., Reid, S.W.J., Mellor, D.J., 2004. Frozen storage of  
380 *Escherichia coli* O157 in buffered peptone water and its detection on bovine  
381 carcasses. J. Food Protect. 67, 40-45.

382

383

384

385

386

387

388

389

390

391

392

393

394

395

396

397 **Figure Legends to Illustrations**

398 **Fig. 1a.** Results of the limit of detection dilution assay for *Escherichia coli* (*E.*  
399 *coli*) O157, *E. coli* O111 and *E. coli* O26 in minced beef using this rapid  
400 technique. Lane M = molecular weight marker VIII (Roche); Lane 1 = *E. coli* O26  
401  $10^{-7}$  (15 CFU/ml); Lane 2 = *E. coli* O26  $10^{-8}$  (0 CFU/ml); Lane 3 = *E. coli* O111  
402  $10^{-7}$  (10 CFU/ml); Lane 4 = *E. coli* O111  $10^{-8}$  (0 CFU/ml); Lane 5 = *E. coli* O157  
403  $10^{-7}$  (40 CFU/ml); Lane 6 = *E. coli* O157  $10^{-8}$  (4 CFU/ml); Lane 7 = *E. coli* O157  
404  $10^9$  (0 CFU/ml); Lane 8 = Control (sample); Lane 9 = multiplex positive control  
405 consisting of equivalent template DNA from three independent *E. coli* serogroups  
406 [including O26, O111 and O157]; Lane 10 = No template PCR control.  
407 A 2% (w/v) agarose gel in 0.5 X Tris-Borate-EDTA (TBE) buffer containing 0.5  
408 mg/ml ethidium bromide.

409

410 **Fig. 1b.** Melt curve graph of SYBR-1 green real-time PCR assay for *vt2*. Melting  
411 temperature ( $T_m$ ), 89°C. *E. coli* O157 positive control (NCTC 12900), (red peak);  
412 Isolate 4, (green peak). Other isolates and the negative control are shown as flat  
413 lines.

414

415 **Fig. 1c.** Melt curve graph of SYBR-1 green real-time PCR assay for *eae* and  
416 *hlyA* genes. Melting temperature ( $T_m$ ) of 88°C for *eae* and 87°C for *hlyA*. *E. coli*  
417 O157 positive control, (pink peak, *hlyA*; green peak, *eae*); Isolate 4, (red peak,  
418 *hlyA*); Isolate 6, (blue peak, *eae*). Other isolates and the negative control are  
419 shown as flat lines.

**Table 1.** List of the micro-organisms used to test the specificity of the primers, with their corresponding reactions.

Micro-organism	Source	*O26 <i>wzx</i>	*O111 <i>rfb</i>	*O157 <i>rfb</i>
<i>Clostridia perfringens</i>	NCTC 8237	-	-	-
<i>Vibrio parahaemolyticus</i>	NCTC 10885	-	-	-
<i>Salmonella poona</i>	NCTC 4840	-	-	-
<i>Staphylococcus aureus</i>	NCTC 6571	-	-	-
<i>Listeria monocytogenes</i>	NCTC 11994	-	-	-
<i>Bacillus cereus</i>	NCTC 7464	-	-	-
<i>Pseudomonas aeruginosa</i>	NCTC 10662	-	-	-
<i>Citrobacter freundii</i>	NCTC 9750	-	-	-
<i>Enterobacter faecalis</i>	NCTC 775	-	-	-
<i>E. coli</i> O126	Clinical Isolate	-	-	-
<i>E. coli</i> O128	Clinical Isolate	-	-	-
<i>E. coli</i> O142	Clinical Isolate	-	-	-
<i>E. coli</i> O124	Clinical Isolate	-	-	-
<i>E. coli</i> O44	Clinical Isolate	-	-	-
<i>E. coli</i> O55	Clinical Isolate	-	-	-
<i>E. coli</i> O111	NCTC 9111	-	+	-
<i>E. coli</i> O26	NCTC 8781	+	-	-
<i>E. coli</i> O157	NCTC 12900	-	-	+

\* target present ( + ), absent ( - ).

**Table 2.** Oligonucleotide primers used to identify *Escherichia coli* serogroups.

<u>Oligonucleotide primers</u>	<u>Gene</u>	<u>Sequence (5' to 3')</u>	<u>Size</u>	<u>% G+C</u>	<u>Amplicon (bp)</u>
O111-F	<i>rfb E<sub>O111</sub></i>	TAGAGAAATTATCAAGTTAGTTCC	[24-mer]	29	406
O111-R		ATAGTTATGAACATCTTGTTTAGC	[24-mer]	29	
O157-F	<i>rfb E<sub>O157</sub></i>	CGGACATCCATGTGATATGG	[20-mer]	50	259
O157-R		TTGCCTATGTACAGCTAATCC	[21-mer]	43	
O26-F	<i>wzx E<sub>O26</sub></i>	GCGCTGCAATTGCTTATGTA	[20-mer]	45	152
O26-R		TTTCCCCGCAATTTATTCAG	[20-mer]	40	


**Table 3.** Characteristics of the *Escherichia coli* O26 isolates detected in raw minced beef utilizing the triplex PCR.

Isolate number	Package type	Detection Method		*Virulence Markers			
		IMS	New PCR Technique	vt1	vt2	eae	hlyA
1	Prepacked	Detected	Detected	-	-	-	-
2	Prepacked	Detected	Detected	-	-	-	-
3	Prepacked	Not Detected	Detected	-	-	-	-
4	Loose	Not Detected	Detected	-	+	-	+
5	Loose	Not Detected	Detected	-	-	-	-
6	Prepacked	Not Detected	Detected	-	-	+	-


\* virulence target present (+), absent (-).

# Figures

[a].



[b].





[c].

