1978-01-01

A Topological Approach to a Problem of Nunke

Brendan Goldsmith

Technological University Dublin, brendan.goldsmith@tudublin.ie

Follow this and additional works at: <https://arrow.tudublin.ie/scschmatart>

Part of the [Mathematics Commons](#)

Recommended Citation

Goldsmith, B. (1978). A topological approach to a problem of Nunke. *Archiv. Math. (Basel)*, 30, (pg. 70-74). doi:10.1007/BF01226052

This Article is brought to you for free and open access by the School of Mathematics at ARROW@TU Dublin. It has been accepted for inclusion in Articles by an authorized administrator of ARROW@TU Dublin. For more information, please contact arrow.admin@tudublin.ie, aisling.coyne@tudublin.ie, gerard.connolly@tudublin.ie.

A topological approach to a problem of Nunke

By

B. GOLDSMITH

1. Introduction. If P denotes the Specker group i.e. the direct product of countably many infinite cyclic groups, then the following question was raised by Nunke [6]: "If P has a subgroup A such that $P/A \cong \mathbb{Q}$, the additive group of rationals, is $A \cong P$?"

The question has been answered negatively by Meijer [4]. However the problem arose from essentially homological work of Nunke and Meijer's solution was also homological. We present here a solution to the problem which uses a topological idea introduced by Lady [3] and avoids all homological machinery.

The word group will be understood to refer to an additively written abelian group and we refer to the standard works [1] and [2] of Fuchs for notation and terminology.

2. The Main Result. A Specker group P may be topologized by using the product topology obtained by considering each component as being discretely topologized. We refer to this topology on P simply as the product topology. P may also be topologized by choosing as a basis of neighbourhoods of 0 the subgroups nP ($n \in \mathbb{Z}$, $n \neq 0$). This is the familiar $\mathbb{Z}$ -adic topology (see [1] page 30). We will make use of a topology, which, following Lady [3], we call the strong topology. This is the topology on P which is the supremum of the $\mathbb{Z}$ -adic and product topologies on P .

The result we require can be stated as

Theorem 1 (Meijer). *Let G be a torsion-free group of at most countable rank and let X be an extension of a Specker group P by G . Then $X \cong P$ if and only if G is free of finite rank.*

We use the following results:

Lemma 2. *If $Y \leq P$ and $Y \cong P$ then there are integers k_n and elements a_n of P ($n = 1, 2, 3, \dots$) such that*

$$P = \prod_{n=1}^{\infty} \langle a_n \rangle \quad \text{and} \quad Y = \prod_{n=1}^{\infty} \langle k_n a_n \rangle$$

Proof. Suppose $Y = \prod_{n=1}^{\infty} \langle y_n \rangle$. Define elements w_n of P by

$$\begin{aligned} w_1 &= (y_1, 0, 0, \dots), \\ w_2 &= (0, y_2, y_3, y_4, \dots), \\ w_3 &= (0, 0, y_3, y_4, \dots), \\ &\vdots \\ w_n &= (0, 0, \dots, 0, y_n, y_{n+1}, \dots) \quad \text{etc.} \end{aligned}$$

Since, for every positive integer j , the j 'th components of almost all the w_n vanish, the infinite sum $\sum_{n=1}^{\infty} s_n w_n$ (s_n integers) makes sense and so the subgroup W consisting of all such infinite sums is a product in the sense defined by Fuchs [2] § 95.

Claim $W = Y$. For if $w \in W$ then $w = \sum_{n=1}^{\infty} s_n w_n$ ($s_n \in \mathbb{Z}$) and so

$$w = (s_1 y_1, s_2 y_2, (s_2 + s_3) y_3, \dots, (s_2 + s_3 + \dots + s_n) y_n, \dots) \in Y.$$

Thus $W \leq Y$. While conversely if $y \in Y$ then $y = (t_1 y_1, t_2 y_2, \dots, t_n y_n, \dots)$ for integers t_n , and so $y = \sum_{n=1}^{\infty} s_n w_n$ where the s_n are given by

$$s_1 = t_1, \quad s_2 = t_2, \quad s_3 = t_3 - s_2, \quad s_4 = t_4 - s_3, \dots \quad \text{etc.}$$

Thus $y \in W$ and so $W = Y$ as claimed.

Thus Y is a product in the sense of Fuchs [2] § 95 and the result follows from [2] Lemma 95.1.

Proposition 3 (Nunke). *Every epimorphic image of a Specker group P is the direct sum of a cotorsion group and a direct product of at most countably many infinite cyclic groups.*

Proof. See Nunke [5] or Fuchs [2] 95.2.

Proof of Theorem 1. Suppose X is isomorphic to P , then applying Proposition 3 to G where $X/P \cong G$, we see that G is a direct sum of a cotorsion group and a product of infinite cyclic groups. Since G has at most countable rank it is clear that the product of infinite cyclic groups must be a finite product. Moreover since a reduced torsion-free cotorsion group must contain a summand isomorphic to the p -adic integers, for some p , it is clear that the cotorsion summand of G must be divisible. Thus $G = D \oplus F$ where D is divisible and F is a finite product of infinite cyclic groups i.e. F is free of finite rank. It remains to show that $D = 0$.

Choose a subgroup Y of X such that $Y/P \cong F$. Clearly $Y \cong P$ and $X/Y \cong D$. Now applying Lemma 2 we may write

$$X = \prod_{n=1}^{\infty} \langle a_n \rangle \quad \text{and} \quad Y = \prod_{n=1}^{\infty} \langle k_n a_n \rangle.$$

We topologize X with the strong topology. (We remark that we are only concerned with this presentation of X and we don't need to show that the product topology on X is independent of the presentation although this is indeed true.) The proof is completed by the following result:

Lemma 4. *Y is closed and dense in X in the strong topology on X .*

Proof. (i) *Y is dense.* If x is in X then a basis for the neighbourhoods of x in the strong topology is $x + m \prod_{n \notin J} \langle a_n \rangle$, where J is a finite subset of $\{1, 2, \dots\}$ and $m \in \mathbb{Z} \setminus \{0\}$.

To show density it suffices to show that for any finite subset J and any integer $m \neq 0$,

$$Y \cap (x + m \prod_{n \notin J} \langle a_n \rangle) \text{ is non-null.}$$

Since Y is dense in X in the $\mathbb{Z}$ -adic topology, we may write for any non-zero integer m , $x = mx' + y$, some $x' \in X$, $y \in Y$. Say $x' = (\dots, r_i a_i, \dots)$. Then set $x' = u + v$ where $u = (\dots, r_i a_i, \dots)_{i \notin J}$, and $v = (\dots, r_i a_i, \dots)_{i \in J}$. Then $x - mu = mv + y$. But Y is pure in X and $k_n a_n \in Y \cap k_n X$, so $a_n \in Y$ each n . Since v is just a finite sum of multiples of elements of Y , v is in Y . Thus $x - mu \in Y$ i.e.

$$Y \cap (x + m \prod_{n \notin J} \langle a_n \rangle) \text{ is non-null.}$$

(ii) *Y is closed in X .* Since the strong topology is the supremum of the product and $\mathbb{Z}$ -adic topologies, it is clearly sufficient to show Y is closed in the product topology on X . If $\bar{Y}$ denotes the closure of Y in the product topology on X then we see that $x \in \bar{Y}$ if, and only if, for each $n = (1, 2, \dots)$ there is a u^n in Y such that $x_i = u_i^n$ for all $i < n$. (Here the subscript i denotes the i -th component.)

So if $x \in \bar{Y}$ then there is a u^2 in Y such that

$$x_1 = u_1^2 = r_1 k_1 a_1 \quad \text{some } r_1 \in \mathbb{Z}.$$

Similarly $x_2 = u_2^3 = r_2 k_2 a_2 \quad \text{some } r_2 \in \mathbb{Z} \quad \text{etc.}$

But then $x = (r_1 k_1 a_1, r_2 k_2 a_2, \dots, r_i k_i a_i, \dots)$ is clearly in Y . Thus Y is closed as required.

We can easily deduce

Corollary 5. *If X is a pure subgroup of the Specker group P such that P/X is of at most countable rank, then $X \cong P$ if and only if $P/X \cong \mathbb{Z}^n$ for some non-negative integer n .*

Remark. It is hoped that this paper, along with Lady's paper [3], will illustrate how useful simple topological techniques may be in areas which have hitherto used homological machinery.

Acknowledgement. I would like to express my thanks to my colleague Raymond Flood for several interesting and helpful conversations on some of the topological aspects of this work.

References

- [1] L. FUCHS, Infinite Abelian Groups, Vol. I. New York and London 1970.
- [2] L. FUCHS, Infinite Abelian Groups, Vol. II. New York and London 1973.
- [3] E. L. LADY, Slender rings and modules. *Pacific J. Math.* **49**, 397—406 (1973).
- [4] A. R. MEIJER, A note on separable homogeneous extensions of the Specker group. *Arch. Math.* **24**, 123—125 (1973).
- [5] R. J. NUNKE, Slender groups. *Bull. Amer. Math. Soc.* **67**, 274—275 (1961); *Acta Sci. Math. (Szeged)* **23**, 67—73 (1962).
- [6] R. J. NUNKE, Problem posed in: *Topics in Abelian Groups*, Chicago 1963.

Eingegangen am 10. 2. 1977

Anschrift des Autors:

B. Goldsmith
College of Technology
Kevin Street
Dublin 8, Ireland
and
School of Theoretical Physics
Dublin Institute for Advanced Studies
Dublin 4, Ireland