2009-01-01

Macular Pigment and its Contribution to Spatial Vision

James Loughman

Technological University Dublin, james.loughman@tudublin.ie

Peter Davison

Technological University Dublin, Peter.Davison@tudublin.ie

Grainne Scanlon

Technological University Dublin, grainne.scanlon@tudublin.ie

See next page for additional authors

Follow this and additional works at: <https://arrow.tudublin.ie/scschphycon>

Part of the [Medicine and Health Sciences Commons](#)

Recommended Citation

Akkali, M., Nolan, J., & Beatty, S., (2009). Macular Pigment and Its Contribution to Spatial Vision. *Association for Research in Vision and Ophthalmology Conference Investigative Ophthalmology and Visual Science*. doi:10.1111/j.1755-3768.2008.644.x

This Conference Paper is brought to you for free and open access by the School of Physics & Clinical & Optometric Science at ARROW@TU Dublin. It has been accepted for inclusion in Conference Papers by an authorized administrator of ARROW@TU Dublin. For more information, please contact arrow.admin@tudublin.ie, aisling.coyne@tudublin.ie, gerard.connolly@tudublin.ie.
Funder: Bausch & Lomb; Enterprise Ireland

Authors

James Loughman, Peter Davison, Grainne Scanlon, and Veronica O'Dwyer

Macular Pigment and its Contribution to Spatial Vision

Mukunda C Akkali¹, John M Nolan¹, Stephen Beatty¹, Veronica O'Dwyer², Peter A Davison², Grainne Scanlon², Philip Major¹, James Loughman²

¹Macular Pigment Research Group, Department of Chemical & Life Sciences, Waterford Institute of Technology, Waterford, Ireland;

²Macular Pigment Research Group, Optometry Department, Dublin Institute of Technology, Dublin, Ireland

INTRODUCTION

Macular pigment (MP), which is composed of three dietary carotenoids, lutein (L), zeaxanthin (Z) and *meso*-zeaxanthin (MZ), is predominantly located at the site of maximum visual acuity in the human retina. MP absorbs short wavelength light prior to photoreceptor stimulation. Filtering such defocused short wavelength (blue) light could enhance visual performance by reducing the effects of chromatic aberration and light scatter.¹ This hypothesis, in relation to MP, remains unproven and poorly investigated to date.

Figure 1.

Central, pre-receptor retinal location of MP. Adapted from Trieschmann et al., 2008²

PURPOSE

The **Collaborative Optical Macular Pigment ASsessment Study** (COMPASS) is a double blind, placebo controlled and randomized clinical trial, designed to investigate whether MP optical density (MPOD) influences spatial vision and visual comfort by means of selective short wavelength light absorption prior to photoreceptor light capture.

SUBJECTS and METHODS

We recruited 142 young healthy subjects (mean $\pm$ SD age = 28.85 ± 6.37 years). The spatial profile of MPOD was assessed by customized heterochromatic flicker photometry (cHFP), which has been shown to be a reliable and repeatable method of quantifying MPOD.³ Visual performance was assessed using visual acuity, mesopic and photopic contrast sensitivity, glare sensitivity and photostress recovery time measures. Each subject also completed a visual performance questionnaire, generating a performance index of the subject's perception of their functional vision in terms of colour, acuity, glare, light/dark adaptation and daily visual tasks.

MPOD measurement

MPOD measurements were obtained by cHFP (using the Macular Densitometer™) at five loci (0.25°, 0.50°, 1°, 1.75° and 3° of retinal eccentricity), with a reference point at 7°, to generate a complete spatial profile of MP.

Best corrected visual acuity (BCVA)

BCVA was assessed using a computer-generated LogMAR, high contrast chart, employing a SLOAN ETD RS letterset. A visual acuity rating (VAR) was computed to quantify precise acuity limits.

Contrast sensitivity function (CSF)

Mesopic (3 cd/m²) and photopic (100cd/m²) CSF curves were plotted using spatial frequencies of 1 cycle per degree (cpd), 4.1 cpd, 7.5 cpd, 11.8 cpd and 20.7 cpd. Gabor stimuli were displayed on a CRT monitor and sensitivities were determined using a linear staircase procedure.

Glare sensitivity

Mesopic (3 cd/m²) and photopic (85 cd/m²) contrast sensitivity was also tested under glare conditions using a sine grating functional acuity contrast test, and a radial glare source. CSF curves were plotted using spatial frequencies of 1.5 cpd, 3 cpd, 6 cpd, 12 cpd and 18 cpd. Glare effects on the CSF were determined for medium (35 Lux) and high (128 Lux) glare intensities.

Photostress Recovery Time (PRT)

PRT was evaluated using a previously described macular automated photostress test⁴ using a Humphrey 745 visual field analyzer.

RESULTS

These results represent the baseline findings from COMPASS. Mean $\pm$ SD MPOD obtained at peak (0.25° eccentricity) was 0.48 ± 0.19 . There was a statistically significant positive relationship between BCVA and MPOD at all retinal eccentricities ($r = 0.236$ to 0.354 , $p < 0.01$) [Figures 2 and Figure 3 illustrate the relationship at 0.25° and 0.50° respectively].

Figure 2.

Figure 3.

There was a statistically significant positive relationship between mesopic and photopic contrast sensitivities and MPOD, but confined to the central 0.50° of retinal eccentricity (mesopic contrast sensitivity @ 7.5 cpd, 11.8 cpd and MPOD at 0.25°, 0.50°: $r = 0.171$ to 0.210 , $p < 0.05$; photopic contrast sensitivity @ 1.0 cpd, 7.5 cpd, 11.8 cpd and MPOD at 0.25°, 0.50°: $r = 0.174$ to 0.197 , $p < 0.05$; [Figures 4 and 5 illustrate the relationship between MPOD at 0.25° and both mesopic and photopic contrast sensitivity at 7.5cpd respectively].

Figure 4.

Figure 5.

There was an inverse relationship between mesopic contrast sensitivity under glare conditions for all spatial frequencies and MPOD at all degrees of retinal eccentricity, but was not statistically significant ($p > 0.05$).

There was no statistically significant relationship observed between MPOD and any of the functional aspects of subjective visual function questionnaire ($r = -0.188$ to 0.148 , $p > 0.05$). There was also no statistically significant relationship observed between MPOD and PRT. ($r = 0.002$ to 0.102 , $p > 0.05$).

CONCLUSION

Given its short wavelength absorption characteristics and central, prereceptorial location [see Figure 1], MP exhibits ideal properties to optimize visual performance. Visual performance, as assessed by visual acuity and contrast sensitivity measures, does appear to be influenced by MPOD. Photostress recovery and visual performance under glare conditions however show no significant relationship with MPOD.

The lack of consistency among the results possibly reflects the inherent difficulty associated with efforts to clarify the role of MP with respect to visual performance using a cross sectional methodology. A placebo controlled, randomized MP supplementation clinical trial, has the capacity to more adequately address the critical research question.

REFERENCES

- (1) Wooten BR, Hammond BR. Macular pigment: influences on visual acuity and visibility. Prog Ret Eye Res 2002 Mar;21(2):225-240.
- (2) Trieschmann M, van Kuijk FJ, Alexander R et al. Macular pigment in the human retina: histological evaluation of localization and distribution. Eye 2008 January;22(1):132-7.
- (3) Kirby ML, Galea M, Loane E, Stack J, Beatty S, Nolan JM. Foveal Anatomic Associations with the Secondary Peak and the Slope of the Macular Pigment Spatial Profile. Invest Ophthalmol Vis Sci 2009 Mar;50(3):1383-1391.
- (4) Dhalla MS, Fantin A. Macular photostress testing: sensitivity and recovery with an automated perimeter. Retina 2005 February;25(2):189-92.