2006-01-01

Apoptosis is Initiated in Human Keratinocytes Exposed to Signalling Factors from Microbeam Irradiated Cells

Fiona Lyng

Technological University Dublin, fiona.lyng@tudublin.ie

Paula Maguire

Technological University Dublin

A. Kilmurray

Technological University Dublin

See next page for additional authors

Follow this and additional works at: <https://arrow.tudublin.ie/radart>

Recommended Citation

Lyng, F. et al. (2006) Apoptosis is Initiated in Human Keratinocytes Exposed to Signalling Factors from Microbeam Irradiated Cells. *International Journal of Radiation Biology*, Vol. 82, no.6, pp.393-399.
doi:10.1080/09553000600803904

This Article is brought to you for free and open access by the Radiation and Environmental Science Centre at ARROW@TU Dublin. It has been accepted for inclusion in Articles by an authorized administrator of ARROW@TU Dublin. For more information, please contact arrow.admin@tudublin.ie, aisling.coyne@tudublin.ie, gerard.connolly@tudublin.ie.

Authors

Fiona Lyng, Paula Maguire, A. Kilmurray, Carmel Mothersill, C. Shao, M. Folkard, and K. Prise

Apoptosis is initiated in human keratinocytes exposed to signalling factors from microbeam irradiated cells

Lyng FM^{a,1}, Maguire P^a, Kilmurray N^a, Mothersill C^{a,2}, Shao C^b, Folkard M^b and Prise KM^b

^aRadiation and Environmental Science Centre, Focas Institute, Dublin Institute of Technology, Kevin St, Dublin 8, IRELAND

^bCell and Molecular Biophysics Group, Gray Cancer Institute, Mount Vernon Hospital, Northwood, UK

Running Head: Bystander factor induced apoptosis in microbeam irradiated cells

Keywords: Radiation, Bystander Effects, Cell Signalling, Apoptosis

¹Person to whom all correspondence should be sent:

Dr Fiona M Lyng

Radiation and Environmental Science Centre

Focas Institute

Dublin Institute of Technology

Kevin St

Dublin 8

IRELAND

Email: fiona.lyng@dit.ie

²Present address: Medical Physics & Applied Radiation Sciences, McMaster University, Hamilton, Ontario, Canada

Abstract

Purpose: There is now no doubt that bystander signalling from irradiated cells occurs and causes a variety of responses in cells not targeted by the ionising track. However, the mechanisms underlying these processes are unknown and the relevance to radiotherapy and risk assessment remains controversial. Previous research by our laboratory has shown bystander effects in a human keratinocyte cell line, HPV-G cells, exposed to medium from γ irradiated HPV-G cells. The aim of this work was to investigate if similar mechanisms to those identified in medium transfer experiments occurred in these HPV-G cells when they are in the vicinity of microbeam irradiated cells. Demonstration of a commonality of mechanisms would support the idea that the process is not artifactual.

Materials and Methods: HPV-G cells were plated as two separate populations on mylar dishes. One population was directly irradiated using a charged particle microbeam (1 - 10 protons). The other population was not irradiated. Bystander factor induced apoptosis was investigated in both populations following treatment by monitoring the levels of reactive oxygen species and mitochondrial membrane potential using fluorescent probes. Expression of the anti-apoptotic protein, bcl-2, and cytochrome c were determined, as well as apoptosis levels.

Results: Microbeam irradiation induced increases in reactive oxygen species and decreases in mitochondrial membrane potential at 6 hours post exposure, increased expression of bcl-2 and cytochrome c release at 6.5 hours and increased apoptosis at 24 hours.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Conclusion: This study shows that similar bystander signalling pathways leading to apoptosis are induced following microbeam irradiation and following medium transfer. This demonstrates that the mechanisms involved are common across different radiation qualities and conditions and indicates that they may be relevant *in vivo*.

Introduction

There has been considerable interest recently in non-targeted effects of radiation that cannot be attributed to direct DNA damage. Recent research has shown that low doses of ionising radiation can cause a “bystander effect”, where radiation damage occurs in cells not directly irradiated (see reviews; Mothersill and Seymour 2003, Lorimore *et al*, 2003, Little 2003, Morgan 2003).

Studies with very low doses of α particles have shown increases in the number of sister chromatid exchanges (SCE) (Nagasawa and Little 1992, Deshpande *et al*, 1997), chromosomal aberrations (Lorimore *et al*, 1998) and activation of stress inducible signalling pathways (Azzam *et al*, 1998, 2001, 2003) in considerably more cells than would have been traversed by an α particle.

Microbeams, which allow the targeting of individual cells or subcellular locations within a population with a charged particle beam, have become useful tools in the study of radiation induced bystander effects. Microbeam studies have shown that effects of single cell irradiation are not limited to the exposed cell but affect other cells in the vicinity. Micronucleus formation and apoptosis (Prise *et al*, 1998, Belyakov *et al*, 2001, 2003), mutation (Zhou *et al*, 2000, 2001) and oncogenic transformation (Sawant *et al*, 2001) have all been shown in cells distant from the target cell.

Medium transfer experiments have shown that bystander effects may be mediated by damage signals released into the culture medium by irradiated cells. Non-irradiated cells incubated with conditioned medium from irradiated cells have shown similar effects to the directly irradiated cells. Mothersill and Seymour (1997, 1998) reported that cell death was induced in unirradiated cells treated with medium from directly irradiated

1
2
3 epithelial cells. This was further shown to be associated with early apoptotic events such
4
5 as calcium fluxes, loss in mitochondrial membrane permeability and the induction of
6
7 reactive oxygen species (ROS) (Lyng *et al*, 2000, 2002). Lehnert *et al* (1997) and
8
9 Narayanan *et al* (1997) also used medium transfer experiments to show that extracellular
10
11 factors including ROS were released by α particle irradiated cells leading to increased
12
13 SCE in nonirradiated cells. Conditioned medium from α particle irradiated cells has also
14
15 been shown to stimulate proliferation in nonirradiated cells (Iyer *et al*, 2000). A recent
16
17 study by Suzuki *et al* (2004) demonstrated that cells irradiated with α particles released
18
19 medium borne factors which induced chromatin damage in bystander cells plated on the
20
21 other side of a medium filled double mylar dish.
22
23
24
25
26
27
28
29

30 To date the majority of reports on bystander effects have used either direct exposure to
31
32 high linear energy transfer (LET) helium ions delivered by low fluence sources
33
34 (Nagasawa and Little 1992), direct irradiation using microbeam approaches (Prise *et al*,
35
36 1998, Zhou *et al*, 2000, Sawant *et al*, 2001, Shao *et al*, 2003) or medium transfer after
37
38 low LET exposure (Mothersill and Seymour 1997).
39
40

41 Two main models have emerged on the mechanisms of bystander responses; cell - cell
42
43 communication through gap junctions and secretion of a cytotoxic factor into the
44
45 medium. The model appears to depend on the cell type used for the experiments and on
46
47 the way the experiments were performed. Some groups have demonstrated a requirement
48
49 for gap junctional intercellular communication (GJIC) to mediate a bystander response
50
51 (Azzam *et al*, 2001, Shao *et al*, 2003b) but other groups have shown GJIC not to be
52
53 involved (Mothersill and Seymour 1997, Lehnert *et al*, 1997).
54
55
56
57
58
59
60

A role for ROS in radiation induced bystander effects has been reported by many groups (Lehnert *et al* 1997, Azzam *et al* 2002, Lyng *et al* 2006). Specifically, nitric oxide, an important signalling molecule, has been shown to induce bystander effects (Matsumoto *et al* 2000, Shao *et al* 2001, 2002, 2003a, 2004)

The aim of the present study was to investigate bystander responses following microbeam irradiation in a human keratinocyte cell line (HPV-G cells) which has previously been shown to undergo apoptosis when exposed to medium from γ irradiated cells (Lyng *et al*, 2000, 2002, Maguire *et al*, 2005). Mitochondrial membrane potential depolarisation, cytochrome c release, bcl-2 expression, induction of reactive oxygen species (ROS) and apoptosis levels were measured in HPV-G cells which were in the vicinity of microbeam irradiated HPV-G cells.

Materials and Methods

Cell Culture

Human keratinocytes, HPV-G cells, immortalised with the human papilloma virus (HPV) were originally obtained as a kind gift from Dr. J. Di Paolo, NIH Bethesda (Pirisi *et al*, 1988). HPV-G cells were cultured in Dulbecco's Modified Eagle Medium (DMEM) : F12 (1:1) medium (Sigma, Dorset, U.K.) containing, 10% fetal bovine serum (Gibco, Irvine, U.K.), 1% penicillin-streptomycin solution 1000 IU (Gibco, Irvine, U.K.), 2mM L- glutamine (Gibco, Irvine, U.K.) and 1µg/ml hydrocortisone (Sigma, Dorset, U.K.) Cells were maintained in an incubator at 37°C, with 95% humidity and 5% CO₂. Subculture was routinely performed when cells were 80-100% confluent, using a 1:1 solution of 0.25% trypsin and 1mM versene (Sigma, Dorset, U.K.) at 37° C.

Microbeam irradiation

For microbeam experiments, plateau phase cells were seeded into specially designed dishes (Folkard *et al*, 1997) consisting of a 34 mm diameter base composed of a 4 µm thick mylar membrane. Two areas of the dish diagonally opposed to each other had been pretreated with 1 µg/ml CellTak adhesive (Becton Dickinson, Oxford, U.K.). Each area was about 5 mm in diameter. Cells were seeded 16 h prior to irradiation to allow full attachment. Typically cells were seeded at a density to allow 600–800 cells in each area of the dish. One hour prior to irradiation cells were incubated with 1 µM Hoechst 33258 (Molecular Probes, Leiden, The Netherlands). At the time of irradiation the cell culture medium was replaced with fresh medium containing 20 mM 4-(2-hydroxyethyl)-1-piperazineethanesulfonic acid (HEPES) (Sigma, Dorset, U.K.) and irradiation was

performed at room temperature. All the cells in one area of the dish had exact numbers of 3.2 MeV protons delivered through the centre of the cell nucleus. Details of the automatic cell finding and imaging system are described in Folkard *et al* (1997). The cells in the unirradiated area of the dish were bystander cells. The irradiation procedure typically took around 10 min after which fresh medium was added to the cells and incubation continued at 37°C for 6 or 24 hours prior to scoring. Control cells were sham irradiated by incubating with 1 µM Hoechst 33258 (Molecular Probes, Leiden, The Netherlands) as before and scanning, finding and following the same number of cells as in the irradiated dishes but without actual irradiation.

Measurement of mitochondrial membrane potential

Mitochondrial membrane potential was determined using rhodamine 123, a green fluorescent dye that accumulates in active mitochondria with high membrane potential. Cultures were washed twice with a buffer containing 130 mM NaCl, 5 mM KCl, 1 mM Na₂HPO₄, 1 mM CaCl₂, 1 mM MgCl₂ and 25 mM HEPES (pH 7.4). Cells were loaded with 5 µM Rhodamine 123 (Sigma, Dorset, U.K.) for 30 min in the buffer at 37 °C. Subsequently, the cultures were washed three times with buffer. Fluorescence images were recorded using a Zeiss Axioskope epifluorescence microscope with a BP 450-490 nm excitation filter and LP515 emission filter (Carl Zeiss Ltd, Welwyn Garden City, U.K.) and a cooled charge coupled device (CCD) camera system (Photonic Science, UK). The mean fluorescence intensity (or mean grey value) was determined using the software package, NIH Image (National Institutes of Health, USA).

Measurement of reactive oxygen species

Induction of reactive oxygen species was measured using 2,7 - dichlorofluorescein diacetate (DCF-DA) (Sigma, Dorset, U.K.). Once inside a cell, the acetate group is cleaved by cellular esterases leaving dichlorofluorescein which emits green fluorescence when oxidized by the reactive oxygen species, hydrogen peroxide and nitric oxide. Cultures were washed twice with a buffer containing 130 mM NaCl, 5 mM KCl, 1 mM Na₂HPO₄, 1 mM CaCl₂, 1 mM MgCl₂ and 25 mM HEPES (pH 7.4). Cells were loaded with 5 µM 2,7 DCF-DA for 30 min in the buffer at 37 °C. Subsequently, the cultures were washed three times with buffer. The cells were then returned to the incubator for 30 mins and washed once more prior to scoring. Fluorescence images were recorded using a Zeiss Axioskope epifluorescence microscope with a BP 450-490 nm excitation filter and LP515 emission filter (Carl Zeiss Ltd, Welwyn Garden City, U.K.) and a cooled CCD camera system (Photonic Science, UK). The mean fluorescence intensity (or mean grey value) was determined using the software package, NIH Image (National Institutes of Health, USA).

Immunocytochemistry

Following the fluorescence measurements as described above (ie. approx 6.5 hours after microbeam irradiation), the cells were washed twice in phosphate buffered saline (PBS) to remove any debris and then fixed in 10% buffered formalin. The mylar on which the cells were attached was removed from the microbeam dish and adhered to a glass slide with the cells facing upwards. The slides were marked to indicate the irradiated and

bystander areas. The slides were immersed in PBS in plastic coplin jars and transported back to Dublin, stored at 4°C and assayed within 48 hours. Immunocytochemical analysis was performed using the Streptavidin Peroxidase method for cell culture using the Vectastain ABC kits (Vector Laboratories, UK). The primary antibody, mouse monoclonal, anti-Bcl-2 (Dako, Denmark) or anti-cytochrome c (Dako, Denmark) was applied for one hour (1:50 and 1:250 dilution respectively). Biotinylated anti-mouse reagent was then added to the cells for 30 minutes followed by Streptavidin Peroxidase for a further 30 minutes, with a wash in PBS in between each step. The chromagen, 0.02% DAB (Sigma, Dorset, U.K.) was then added for 10 minutes in darkness and washed off in distilled water. Cells were then counterstained with Harris haematoxylin and mounted with glycergel. A negative control, where no primary antibody was added was included in each experimental run. Positive staining was determined by brown staining in the cytoplasm. Numbers of cells positive for bcl-2 or cytochrome c were scored blind in both direct and bystander areas on each of three replicate slides and expressed as the percentage of the total cells counted (approx. 400 - 500 cells were counted in each area).

Quantification of apoptosis

Twenty four hours after microbeam irradiation, the cells were washed twice in PBS to remove any debris and then fixed in 10% buffered formalin. The mylar was removed and adhered to a glass slide as described above. The slides were transported back to Dublin immersed in PBS in plastic coplin jars, stored at 4°C and assayed within 48 hours. The cells were stained for 15 mins with 1 µg / ml Propidium Iodide (Sigma, Dorset, U.K.) to visualise the nuclei. Slides were scored for the presence of apoptotic cells using a Zeiss

Axioplan epifluorescence microscope equipped with BP546/12 nm excitation filter and LP590nm emission filter (Carl Zeiss Ltd, Welwyn Garden City, U.K.). Cells were defined as apoptotic if they displayed evidence of two or more of the following; cell volume shrinkage and pycnotic nucleus (chromatin condensation), nuclear fragmentation and formation of apoptotic bodies (Kerr and Harmon 1991). Numbers of apoptotic cells were scored blind in both direct and bystander areas on each of three replicate slides and expressed as the percentage of the total cells counted (approx. 500 cells were counted in each area).

Statistical analysis

Microscope and detector parameters were standardised to allow quantitative comparisons of the relative fluorescence intensity of the cells between groups. All measurements are presented as mean values $\pm$ S.E. of 3 independent experiments with 3 replicate dishes per measurement. Significance of differences was determined by a student's unpaired t-test and the differences were considered significant if $p \leq 0.05$.

Results

A significant reduction in rhodamine 123 fluorescence, indicating mitochondrial membrane potential depolarisation, was observed in HPV-G cells 6 hours after direct microbeam irradiation (figure 1). Cells that were not directly irradiated but were in the same dish (bystander cells) also showed a similar significant reduction in fluorescence (figure 1).

A significant increase in DCF-DA fluorescence, indicating an increase in ROS, was observed in HPV-G cells 6 hours after direct microbeam irradiation (figure 2). Cells in another area of the same dish (bystander cells) also showed a significant increase in fluorescence (figure 2).

Both the directly irradiated cells and the bystander cells showed increased expression of bcl-2 6.5 hours after microbeam irradiation (table I). Similarly increased levels of cytochrome c were observed 6.5 hours after microbeam irradiation in both the directly irradiated cells and the bystander cells (table I).

HPV-G cells showed increased apoptosis 24 hours after direct microbeam irradiation (figure 3). Cells that were not directly irradiated but were in the same dish (bystander cells) also showed a similar increase in apoptosis levels (figure 3).

For all endpoints measured, there was no significant difference between the effect in the directly irradiated cells and the bystander cells. Similarly there was no significant difference between the effect following irradiation with 1 or 10 protons to the directly exposed cells.

Discussion

This study has shown a significant reduction in mitochondrial membrane potential and a significant increase in ROS in both directly irradiated and bystander HPV-G cells 6 hours after microbeam irradiation. Increased bcl-2 expression and cytochrome c release, after approx 6.5 hours, and increased apoptosis, after 24 hours, were also observed. No significant differences were observed between the effects of different doses.

These results are very similar to those previously reported for HPV-G cells exposed to medium from γ irradiated cells (Lyng *et al*, 2000, 2002, Maguire *et al*, 2005). This finding is important as it shows the mechanisms are similar for medium transfer and for microbeam irradiation and therefore more likely to be universal.

The findings from this study and from previous studies (Lyng *et al*, 2000, 2002, Maguire *et al*, 2005) have shown that early apoptotic events, such as mitochondrial membrane potential depolarisation, induction of ROS, expression of bcl-2 and release of cytochrome c are induced in normal human keratinocytes either exposed to medium from γ irradiated cells or in the vicinity of microbeam irradiated cells. The effects observed appear to be independent of the dose or number of protons delivered to the irradiated cells. A medium borne factor is likely to be involved in both cases. It is unlikely that gap junctional communication is involved in the medium transfer approach and in the present microbeam study the directly irradiated cells and the bystander cells were two distinct populations separated by on average 6 mm.

Most of the studies on bystander effects have used either direct exposure to low fluences of α particles, direct irradiation using microbeam approaches or medium transfer after

low LET exposure. There have been very few reports on the LET dependence of the bystander effect. Hickman *et al* (1994) observed a bystander response, evidenced by increased p53 expression, in rat lung epithelial cells exposed to low fluences of α particles. No increase was seen in cells exposed to similar doses (less than 10 cGy) of X-rays, indicating the existence of a relatively higher damage threshold for sparsely ionizing radiation. Shao *et al* (2002, 2003b) reported an LET dependent induction of micronuclei and cell proliferation in human neoplastic epithelial cells. High LET (100 keV/ μ m) carbon-ion irradiation was found to be more efficient at inducing the medium-mediated bystander effect than low LET (13 keV/ μ m) carbon-ion irradiation. Further studies by the same group compared the bystander responses in primary human fibroblasts individually targeted by a high LET heavy particle microbeam of ^{40}Ar (1260 keV/ μ m) or ^{20}Ne (380 keV/ μ m) (Shao *et al*, 2003b). An increase in micronuclei was observed independent of the LET and the number of particles delivered to the targeted cells. Previous studies by our group has shown increased apoptosis in human keratinocytes exposed to a medium borne factor from cells irradiated with low LET γ rays (Lyng *et al*, 2000, 2002, Maguire *et al*, 2005). The protons used in the present study (3.2 MeV with an LET of ~ 13 keV/ μ m) are essentially low LET and importantly, the degree of bystander responses observed was similar to that obtained with γ -rays.

An important observation from these studies is that the level of effect is the same regardless of whether cells were directly exposed or were neighboring non-targeted bystander cells. This agrees with other studies showing that at least after low dose exposure, the bystander response predominates the overall effect (Schettino *et al*, 2003, Seymour and Mothersill 2000).

References

Azzam EI, de Toledo SM, Gooding T, Little JB. 1998. Intercellular communication is involved in the bystander regulation of gene expression in human cells exposed to very low fluences of alpha particles. *Radiation Research* 150: 497-504

Azzam EI, de Toledo SM, Little JB. 2001. Direct evidence for the participation of gap junction-mediated intercellular communication in the transmission of damage signals from alpha -particle irradiated to nonirradiated cells. *Proceedings of the National Academy of Sciences U.S.A.* 98: 473-478

Azzam EI, de Toledo SM, Spitz DR, Little JB. 2002. Oxidative metabolism modulates signal transduction and micronucleus formation in bystander cells from alpha-particle-irradiated normal human fibroblast cultures. *Cancer Research* 62: 5437-5442

Azzam EI, de Toledo SM, Little JB. 2003. Oxidative metabolism, gap junctions and the ionizing radiation-induced bystander effect. *Oncogene* 22: 7050-7057

Belyakov OV, Malcolmson AM, Folkard M, Prise KM and Michael BD. 2001. Direct evidence for a bystander effect of ionizing radiation in primary human fibroblasts. *British Journal of Cancer* 84: 674-679

Belyakov OV, Folkard M, Mothersill C, Prise KM, Michael BD. 2003. A proliferation-dependent bystander effect in primary porcine and human urothelial explants in response to targeted irradiation. *British Journal of Cancer* 88: 767-774

Deshpande A, Goodwin EH, Bailey SM, Marrone BL and Lehnert BE. 1997. Alpha-particle-induced sister chromatid exchange in normal human lung fibroblasts: evidence for an extranuclear target. *Radiation Research* 145: 260-267

Folkard M, Vojnovic B, Hollis KJ, Bowey AG, Watts SJ, Schettino G, Prise KM and Michael BD. 1997. *International Journal of Radiation Biology* 72: 387-395

Hickman AW, Jaramillo RJ, Lechner JF, Johnson NF. 1994. Alpha-particle-induced p53 protein expression in a rat lung epithelial cell strain. *Cancer Research* 54: 5797-5800

Iyer R, Lehnert BE and Swensson R. 2000. Factors underlying the cell growth-related bystander responses to alpha particles. *Cancer Research* 60: 1290-1298

Kerr JFR and Harmon BV. 1991. Definition and incidence of apoptosis. An historical perspective. In: Tomei LD and Cope FO, editors. *Apoptosis. The Molecular Basis of Cell Death*, New York: Cold Springs Harbour Laboratory, pp 5 - 9

1
2
3 Lehnert BE, Goodwin EH, Desppande A. 1997. Extracellular factor(s) following
4 exposure to alpha-particles can cause sister chromatid exchanges in normal human cells.
5
6 Cancer Research 57: 2164-2171
7
8
9

10
11
12 Little JB. 2003. Genomic instability and bystander effects: a historical perspective.
13
14 Oncogene 22: 6978-6987
15
16
17

18
19
20 Lorimore SA, Coates PJ, Wright EG. 2003. Radiation-induced genomic instability and
21
22 bystander effects: inter-related nontargeted effects of exposure to ionizing radiation.
23
24 Oncogene 22: 7058-7069
25
26
27

28
29
30 Lyng FM, Seymour CB, Mothersill C. 2000. Production of a signal by irradiated cells
31
32 which leads to a response in unirradiated cells characteristic of initiation of apoptosis.
33
34 British Journal of Cancer 83: 1223-1230.
35
36
37

38
39
40 Lyng FM, Seymour CB, Mothersill C. 2002. Initiation of apoptosis in cells exposed to
41
42 medium from the progeny of irradiated cells: a possible mechanism for bystander-
43
44 induced genomic instability? Radiation Research 157: 365-370
45
46
47

48
49
50 Lyng FM, Maguire P, McClean B, Seymour C, Mothersill C. 2006. The involvement of
51
52 calcium and MAP kinase signalling pathways in the production of radiation induced
53
54 bystander effects, Radiation Research (in press)
55
56
57
58
59
60

Maguire P, Mothersill C, Seymour C and Lyng FM. 2005. Medium from irradiated cells induces dose dependent mitochondrial changes and bcl-2 responses in unirradiated human keratinocytes. *Radiation Research* 163: 384-390

Matsumoto H, Hayashi S, Hatashita M, Shioura H, Ohtsubo T, Kitai R, Ohnishi T, Yukawa O, Furusawa Y, Kano E. 2000. Induction of radioresistance to accelerated carbon-ion beams in recipient cells by nitric oxide excreted from irradiated donor cells of human glioblastoma. *International Journal of Radiation Biology* 76: 1649-1657

Morgan WF. 2003. Non-targeted and delayed effects of exposure to ionizing radiation: I. Radiation-induced genomic instability and bystander effects in vitro. *Radiation Research* 159: 567-580

Mothersill C, Seymour C. 1997. Medium from irradiated human epithelial cells but not human fibroblasts reduces the clonogenic survival of unirradiated cells. *International Journal of Radiation Biology* 71 421-427

Mothersill C, Seymour CB. 1998. Cell-cell contact during gamma irradiation is not required to induce a bystander effect in normal human keratinocytes: evidence for release during irradiation of a signal controlling survival into the medium. *Radiation Research* 149 256-262

1
2
3 Mothersill C and Seymour CB. 2004. Radiation-induced bystander effects--implications
4 for cancer. *Nature Reviews Cancer* 4: 158-164
5
6
7

8
9
10 Nagasawa H and Little JB. 1992. Induction of sister chromatid exchanges by extremely
11 low doses of alpha particles. *Cancer Research* 52: 6394 – 6396
12
13
14

15
16
17 Narayanan PK, Goodwin EH, Lehnert BE. 1997. α particles initiate biological
18 production of superoxide anions and hydrogen peroxide in human cells. *Cancer Research*
19
20
21
22 57: 2963-3971
23
24

25
26
27 Pirisi L, Creek KE, Doniger J, DiPaolo J. 1988. Continuous cell lines with altered growth
28 and differentiation properties originate after transfection of human keratinocytes with
29 human papillomavirus type 16 DNA. *Carcinogenesis*. 9: 1573-1579
30
31
32

33
34
35
36
37 Prise KM, Belyakov OV, Folkard M, and Michael B. 1998. Studies of bystander effects
38 in human fibroblasts using a charged particle microbeam. *International Journal of*
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Schettino G, Folkard M, Prise KM, Vojnovic B, Held KD and Michael BD. 2003. Low-dose studies of bystander cell killing with targeted soft X rays, *Radiation Research* 160: 505-511

Seymour CB, Mothersill C. 2000. Relative contribution of bystander and targeted cell killing to the low-dose region of the radiation dose-response curve. *Radiation Research* 153: 508-511.

Shao C, Aoki M, Furusawa Y. 2001. Medium-mediated bystander effects on HSG cells co-cultivated with cells irradiated by X-rays or a 290 MeV/u carbon beam. *Journal of Radiation Research (Tokyo)* 42: 305-316

Shao C, Furusawa Y, Aoki M, Matsumoto H, Ando K. 2002. Nitric oxide-mediated bystander effect induced by heavy-ions in human salivary gland tumour cells. *International Journal of Radiation Biology* 78: 837-844

Shao C, Stewart V, Folkard M, Michael BD and Prise KM. 2003a. Nitric oxide-mediated signaling in the bystander response of individually targeted glioma cells. *Cancer Research* 63 8437-8442

Shao C, Furusawa Y, Kobayashi Y, Funayama T, Wada S. 2003b. Bystander effect induced by high-LET particles in confluent human fibroblasts: a mechanistic study. *FASEB Journal* 17: 1422-1427

Shao C, Aoki M, and Furusawa Y. 2004. Bystander effect in lymphoma cells vicinal to irradiated neoplastic epithelial cells: Nitric oxide is involved. *Journal of Radiation Research* 45: 97-103

Suzuki M, Zhou H, Geard CR, Hei TK. 2004 Effect of medium on chromatin damage in bystander mammalian cells. *Radiation Research* 162: 264- 269

Zhou H, Randers-Pehrson G, Waldren CA, Vannais D, Hall EJ and Hei TK. 2000. Induction of a bystander mutagenic effect of alpha particles in mammalian cells. *Proceedings of the National Academy of Sciences U.S.A.* 97: 2099-2104

Zhou H, Suzuki M, Randers-Pehrson G, Vannais D, Chen G, Trosko JE, Waldren CA, Hei TK. 2001. Radiation risk to low fluences of alpha particles may be greater than we thought. *Proceedings of the National Academy of Sciences U.S.A.* 98:14410-14415

Acknowledgements

The authors acknowledge support from Science Foundation Ireland, the Royal Irish Academy, the Royal Society, the Gray Cancer Institute and Cancer Research UK.

Figure Legends

Figure 1 % Fluorescence from rhodamine 123 in both directly irradiated and bystander HPV-G cells 6 hours after microbeam irradiation. A decrease in fluorescence levels is indicative of a decrease in mitochondrial membrane potential. * $p < 0.001$

Figure 2 % Fluorescence from 2,7 dichlorofluorescein diacetate in both directly irradiated and bystander HPV-G cells 6 hours after microbeam irradiation. An increase in fluorescence levels is indicative of an increase in reactive oxygen species. * $p < 0.001$

Figure 3 % Apoptotic cells in both directly irradiated and bystander HPV-G cells 24 hours after microbeam irradiation. * $p < 0.05$, ** $p < 0.01$

Table I % HPV-G cells positive for bcl-2 and cytochrome c 6.5 hours following microbeam irradiation. * p<0.05, ** p<0.01, *** p<0.005

Dose	% bcl-2 positive cells		% cytochrome c positive cells	
	Direct	Bystander	Direct	Bystander
Control	2.17 ± 0.33	1.67 ± 0.60	1.83 ± 0.44	2.50 ± 0.73
1 proton	7.17 ± 0.73 **	7.00 ± 0.29 *	13.83 ± 0.60 ***	14.17 ± 0.88 **
10 protons	9.00 ± 0.59**	7.67 ± 1.01*	15.67 ± 0.73***	14.33 ± 0.19 **