



2005-01-01

# Minimal modules over valuation domains

Brendan Goldsmith

*Technological University Dublin*, [brendan.goldsmith@dit.ie](mailto:brendan.goldsmith@dit.ie)

P. Zanardo

Follow this and additional works at: <https://arrow.dit.ie/scschmatart>

 Part of the [Mathematics Commons](#)

## Recommended Citation

Goldsmith, B. and Zenardo, P.: Minimal modules over valuation domains. *Journal of Pure and Applied Algebra*, Vol.199, (2005), pp.95-109.

This Article is brought to you for free and open access by the School of Mathematics at ARROW@TU Dublin. It has been accepted for inclusion in Articles by an authorized administrator of ARROW@TU Dublin. For more information, please contact [yvonne.desmond@dit.ie](mailto:yvonne.desmond@dit.ie), [arrow.admin@dit.ie](mailto:arrow.admin@dit.ie), [brian.widdis@dit.ie](mailto:brian.widdis@dit.ie).



This work is licensed under a [Creative Commons Attribution-NonCommercial-Share Alike 3.0 License](#)



# MINIMAL MODULES OVER VALUATION DOMAINS

B. GOLDSMITH AND P. ZANARDO

ABSTRACT. Let  $R$  be a valuation domain. We say that a torsion-free  $R$ -module is *minimal* if it is isomorphic to all its submodules of finite index. Here the usual concept of finite index for groups is replaced by the more appropriate (for module theory) definition: a submodule  $H$  of the module  $G$  is said to be of finite index in  $G$  if the quotient  $G/H$  is a finitely presented torsion module. We investigate minimality in various settings and show *inter alia* that over a maximal valuation domain all torsion-free modules are minimal. Constructions of non-minimal modules are given by utilizing realization theorems of May and the authors. We also show that direct sums of minimal modules may fail to be minimal.

## INTRODUCTION.

The motivation for the present investigation of modules over valuation domains with certain special properties comes, via Abelian group theory, from topology. Recall that a topological space  $Y$  is said to be  $H$ -connected if, and only if every proper local homeomorphism  $\phi$  from a space  $X \rightarrow Y$  is a (global) homeomorphism. If the space  $Y$  is assumed to be a compact manifold then  $Y$  is simply connected provided it has dimension 1, 2. For dimensions  $\geq 4$ , this is known not to be so. In dimension 3 however one encounters the so-called Poincaré Conjecture since an  $H$ -connected 3-manifold is a homology sphere. In an attempt to circumvent this difficulty the notion of an  $h$ -connected space was considered: this is a connected first countable Hausdorff space  $Y$  which has the property that given any connected covering  $\phi : X \rightarrow Y$ ,  $X$  is homeomorphic to  $Y$  via *some* homeomorphism. If  $Y$  is an  $h$ -connected manifold then its fundamental group  $\pi_1(Y)$  has the property: if  $\varphi : H \rightarrow \pi_1(Y)$  is a monomorphism and  $|G : H^\varphi| < \infty$ , then  $H^\varphi \cong \pi_1(Y)$  via *some* isomorphism. Thus a (not necessarily Abelian) group which has the property that it is isomorphic to all its normal subgroups of finite index, is called an *hc-group*. (See [13] and the references therein for further details.) Interestingly, in the category of Abelian groups, the same class of groups arose from considerations of a certain type of ‘quasi-minimality’ arising naturally in point-set topology. Recall that if  $\mathcal{C}$  is a category with an equivalence relation on the objects of  $\mathcal{C}$  and  $\preceq$  is the quasi-order induced from a given property  $\mathcal{P}$  by setting  $A \preceq B$  if  $A$  is equivalent to some subobject of  $B$  having the property  $\mathcal{P}$ , then an object  $M$  is minimal with respect to  $\preceq$  if  $X \preceq M$  implies  $X$  is equivalent to  $M$ . When  $\mathcal{C}$  is the category of Abelian groups with isomorphism of groups as the equivalence and  $\mathcal{P}$  is the property of being of finite index, then *hc-groups* are precisely the minimal objects. These groups have been investigated by Ó hÓgáin and Goldsmith in ([7] and [11]) where they have been termed *minimal*.

Since every proper quotient of the ring of integers is finite, the notion of an Abelian group  $G$  having a subgroup of  $H$  of finite index is both straightforward and natural: the quotient group  $G/H$  is finite. However for modules over an arbitrary

---

1991 *Mathematics Subject Classification*. 13G05, 13A15, 13A17.  
Research supported by MIUR, PRIN 2002.

ring  $R$  this concept is no longer so natural and one needs to find a meaningful extension of the notion if it is to play any significant role.

In the present paper we focus on torsion-free modules over a valuation domain  $R$ . In the first section we introduce the notion of a submodule  $H$  of  $G$  being of ‘finite index’ if the quotient  $G/H$  is *finitely presented and torsion*. With this new understanding of finite index, we say that a torsion-free  $R$ -module  $G$  is *minimal* if it is isomorphic to each of its submodules of finite index. Our principal result in that section is that every torsion-free module over a maximal valuation is minimal. We find classes of torsion-free  $R$ -modules which are always minimal, for instance, free modules or modules with basic rank one.

The second section is devoted to the construction of non-minimal indecomposable torsion-free modules of finite rank over largely general valuation domains. We utilize the so-called realization theorems of May and the authors, [6] and [9]. On the other hand, we show that over special valuation domains (first constructed by Nagata, [10]), every torsion-free module is minimal.

In the final section we consider the problem of whether direct sums and summands of minimal modules remain minimal. Although we obtain some positive results, direct sums of minimal  $R$ -modules may fail to be minimal when  $R$  is not almost maximal.

## 1. PRELIMINARIES AND FIRST RESULTS.

Our notation is standard and any undefined terms may be found in the texts [4] and [5].

Throughout the sequel  $R$  will denote a valuation domain and  $Q$  its field of quotients; the maximal ideal of  $R$  shall always be denoted by  $P$ . Since in the present paper there is no danger of confusion, when we say that  $R$  is a discrete valuation domain we automatically mean that  $R$  has rank one, i.e.  $R$  is Noetherian.

Let  $M$  and  $H$  be torsion-free  $R$ -modules, with  $H \leq M$ . Inspired by the corresponding definition in the Abelian groups case, and with a little abuse of language, we say that  $H$  has *finite index in  $M$*  if  $M/H$  is a *finitely presented torsion  $R$ -module*, that is:  $M/H \cong \bigoplus_{i=1}^n R/r_i R$ , for suitable  $0 \neq r_i \in R$ . An  $R$ -module  $M$  is said to be *minimal* if  $M$  is isomorphic to all of its submodules of finite index.

Throughout the paper we shall deal with torsion-free modules. Thus an  $R$ -module is automatically assumed to be torsion-free, if not otherwise specified.

It is useful to recall that, over a valuation domain  $R$ , finitely generated submodules of finitely presented  $R$ -modules are finitely presented, as well. Indeed, if  $G/K < F/K$ , where  $F$  is free of finite rank, and  $K$  and  $G/K$  are finitely generated, then  $G$  is a finitely generated submodule of the free module  $F$ , whence  $G$  is free, since  $R$  is a valuation domain. Therefore  $G/K$  is finitely presented.

It is convenient to summarize some properties of this new notion of finite index; many standard ideas remain valid, one exception being that if  $A$  is of finite index in  $C$  and  $B$  is a submodule of  $C$  containing  $A$ , then  $B$  is not necessarily of finite index in  $C$ . For instance, take a non-complete valuation domain  $R$  with non-finitely generated maximal ideal  $P$ . In this situation, there exists a rank-two indecomposable  $R$ -module  $M$  with free basic submodule  $X$  of rank one, such that  $M/X$  is divisible. Such a construction is reminiscent of the classical one by Pontryagin for torsion-free Abelian groups (see e.g. [15]). Choose  $0 \neq t \in P$ . From  $M/X$  divisible and  $X$  pure it follows that  $M = X + tM$  and  $M/tM \cong R/tR$ , whence  $tM$  has finite index in  $M$ . On the other hand,  $PM$  contains  $tM$ ,  $M/PM$  is isomorphic to  $R/P$  (see the next Lemma 1.8), and  $R/P$  is not finitely presented, since  $P$  is not finitely generated.

**Proposition 1.1.** *Let  $A < B < C$  be  $R$ -modules.*

(i) If  $A$  has finite index in  $C$ , then  $A$  has finite index in  $B$  if, and only if  $B/A$  is finitely generated.

(ii) If  $A$  has finite index in  $B$  and  $B$  has finite index in  $C$ , then  $A$  has finite index in  $C$ .

*Proof.* If  $A$  has finite index in  $B$  then it is immediate that  $B/A$  is finitely generated. Conversely suppose  $B/A$  is finitely generated. Then  $B/A$  is also finitely presented, being contained in the finitely presented module  $C/A$ . We conclude that  $A$  is of finite index in  $B$ .

To establish (ii) note that we have an exact sequence

$$0 \rightarrow B/A \rightarrow C/A \rightarrow C/B \rightarrow 0,$$

where  $B/A$  and  $C/B$  are finitely presented. We have to show that  $C/A$  is also finitely presented. The above sequence shows that  $C/A$  is finitely generated; write  $C/A = F/K$ , where  $F$  is free of finite rank. Then  $B/A = H/K$  for a suitable submodule  $H$  of  $F$ , and  $C/B \cong F/H$ . Since  $C/B$  is finitely presented, we derive that  $H$  is finitely generated, in view of Proposition 2.1, page 152 of [5] (an immediate application of Schanuel's lemma). Since  $R$  is a valuation domain, finitely generated submodules of free modules are free, so  $H$  is free. Again by Proposition 2.1 of [5], from  $H/K$  finitely presented we get  $K$  finitely generated. We conclude that  $C/A = F/K$  is finitely presented, as desired.  $\square$

**Remark 1.** One may wonder why we did not give a more obvious definition for a submodule  $H$  to be finite index in  $M$ , *viz.* requiring that  $M/H$  be a finite direct sum of cyclic torsion modules. We avoided this definition mainly because in such a case not even  $R$  would be a minimal module (unless  $R$  is a discrete valuation domain). In fact, whenever  $R$  contains a non-finitely generated ideal  $I$ , we have  $R/I$  cyclic and  $R \not\cong I$ .

**Proposition 1.2.** *Let  $H$  be a submodule of finite index of a torsion-free  $R$ -module  $M$ . Let  $x_1, \dots, x_n$  be elements of  $M$  such that  $M/H = \bigoplus_{i=1}^n R(x_i + H)$ , where  $\text{Ann}(x_i + H) = r_i R \neq R$ , for suitable  $r_i \in R$ . Then  $X = \langle x_1, \dots, x_n \rangle$  is a free pure submodule of  $M$ , and  $\{x_1, \dots, x_n\}$  is a basis of  $X$ . Moreover,  $Y = \bigoplus_{i=1}^n R(r_i x_i)$  coincides with  $X \cap H$ . In particular,  $Y$  is a pure submodule of  $H$ .*

*Proof.* Note that  $\text{Ann}(x_i + H) = r_i R \neq R$ , implies that  $x_i \notin H$  for all  $i \leq n$ ; set  $y_i = r_i x_i \in H$ .

Firstly we verify that  $X = \langle x_1, \dots, x_n \rangle = Rx_1 \oplus \dots \oplus Rx_n$ . In fact, assume for a contradiction that  $\sum_{i=1}^n a_i x_i = 0$ , for suitable  $a_i \in R$  not all zero. One of the nonzero coefficients, say  $a_j$ , divides all the  $a_i$ ,  $1 \leq i \leq n$ . Then, since  $M$  is torsion-free, possibly dividing the preceding equality by  $a_j$ , we may assume that  $a_j = 1$ . Reducing modulo  $H$  we get  $x_j + H \in \langle x_i + H : i \neq j \rangle$ , which is impossible. As a consequence, we also have  $Y = \langle y_1, \dots, y_n \rangle = Ry_1 \oplus \dots \oplus Ry_n$ .

Our next step is to prove that  $X$  is a pure submodule of  $M$ . Assume that  $z \in M$  is such that  $0 \neq tz = \sum_{i=1}^n b_i x_i$ ; we have to show that  $t$  divides all the coefficients  $b_i$ . Suppose not: let  $b_j$  be a coefficient which does not divide all the  $b_i$ ,  $1 \leq i \leq n$ ; then  $t$  is a proper multiple of  $b_j$ . Since  $M$  is torsion-free, dividing the preceding equality by  $b_j$ , we may reduce to the case when  $b_j = 1$  and  $t \in P$ . Recall now that  $z + H = \sum_{i=1}^n c_i (x_i + H)$ , for suitable  $c_i \in R$ , whence  $t(z + H) = \sum_{i=1}^n t c_i (x_i + H) = \sum_{i=1}^n b_i (x_i + H)$ . It follows that  $(1 - t c_j)(x_j + H) \in \langle x_i + H : i \neq j \rangle$ , a contradiction, since  $t \in P$ , whence  $1 - t c_j$  is a unit of  $R$ .

It remains to prove that  $Y = X \cap H$ , so that  $X$  pure in  $M$  implies  $Y$  pure in  $H$ , since all the modules are torsion-free. Pick an arbitrary  $h \in X \cap H$ ; we have  $h = \sum_{i=1}^n d_i x_i$ , for suitable  $d_i \in R$ . Then  $\sum_{i=1}^n d_i (x_i + H) = 0$ , and therefore

$d_i \in r_i R$ , for all  $i \leq n$ . It follows that  $h = \sum_{i=1}^n (d_i/r_i)y_i \in Y$ . Since  $h$  was arbitrary, we conclude that  $X \cap H \subseteq Y$ . The opposite inclusion is trivial.  $\square$

The proof of the following lemma may be found in the book by Fuchs and Salce [5].

**Lemma 1.3.** *Let  $F$  be a free  $R$ -module of finite rank, which is a pure submodule of a finite direct sum  $W$  of uniserial  $R$ -modules. Then  $F$  is a direct summand of  $W$ .*

The preceding Proposition 1.2 and Lemma 1.3 allow us to find some interesting classes of modules which are automatically minimal.

**Proposition 1.4.** *Let  $M$  be a torsion-free  $R$ -module.*

(i) *If  $M = PM$ , then  $M$  is minimal. In particular, divisible modules are minimal.*

(ii) *If  $M$  is free, then  $M$  is minimal.*

(iii) *If  $I$  is a fractional ideal of  $R$ , then  $I$  is minimal.*

*Proof.* (i) We will show that in this case a submodule of finite index of  $M$  necessarily coincides with  $M$ . Therefore  $M$  is trivially minimal. In the notation of Proposition 1.2, let us assume for a contradiction that  $M = \langle x_1, \dots, x_n \rangle + H$ , where  $M/H = \bigoplus_{i=1}^n R(x_i + H)$  and  $0 \neq R(x_i + H) \cong R/r_i R$ , for all  $i \leq n$ . In view of Proposition 1.2, the submodule  $X = \bigoplus_{i=1}^n R x_i$  is free and pure in  $M$ . In particular,  $x_i \notin PM$  for all  $i$ . But this is contrary to the assumption  $M = PM$ .

(ii) In the same notation, let  $M = X + H$ , where  $M$  is free. Then  $X$  is a free pure submodule of  $M$ . Since  $X$  has finite rank, it is contained in a finite rank direct summand  $N$  of  $M$ . Therefore, by Lemma 1.3,  $X$  is a direct summand of  $N$ , whence  $X$  is also a direct summand of  $M$ , say  $M = X \oplus M_1$ . Let  $Y = X \cap H$ . By Proposition 1.2,  $Y = \bigoplus_{i=1}^n R(r_i x_i)$  is free. We have

$$M_1 \cong M/X = (X + H)/X \cong H/(X \cap H) = H/Y.$$

Since  $M_1$  is free, we get  $H \cong Y \oplus M_1$ . We conclude that  $H \cong M$ , since  $X \cong Y$ .

(iii) Since  $R$  is a valuation domain, then either  $I$  is principal, and so is free, or  $I = PI$ . In either case the desired result follows from the previous sections of the proposition.  $\square$

Recall that a torsion-free  $R$ -module  $M$  is said to be *separable* if every finite subset of  $M$  is contained in a direct summand  $W$  of  $M$ , where  $W$  is a finite direct sum of uniserial  $R$ -modules (see [4]).

**Theorem 1.5.** *Let  $H$  be a submodule of finite index of a torsion-free  $R$ -module  $M$ . If  $H$  is separable, then  $H$  is isomorphic to  $M$ . In particular,  $M$  is separable, too.*

*Proof.* We follow the notation of Proposition 1.2. Write  $M = X + H$  and  $Y = X \cap H$ . Since  $H$  is separable, and  $Y \cong X$  is free of finite rank, then  $Y$  is contained in a finite direct sum of uniserial modules  $W$  which is a direct summand of  $H$ . By the preceding lemma,  $Y$  is a direct summand of  $W$ , and therefore of  $H$ . Write  $H = Y \oplus H_1$  and observe that  $X \cap H_1 = 0$ , since  $M = X + H$ . It follows readily that  $M = X \oplus H_1$ . We conclude that  $M$  is isomorphic to  $H$ , as desired.  $\square$

**Corollary 1.6.** *Let  $R$  be a maximal valuation domain. Then every torsion-free  $R$ -module is minimal.*

*Proof.* Every torsion-free  $R$ -module is separable. This follows from the fact that uniserial modules over maximal valuation domains are pure-injective (see [5]).  $\square$

**Corollary 1.7.** *If  $M$  is a torsion-free separable minimal  $R$ -module, then a direct summand of  $M$  is again minimal*

*Proof.* Suppose that  $M = A \oplus B$  and that  $H$  is a submodule of  $A$  of finite index. Then  $H \oplus B$  is clearly of finite index in  $M$  and hence is isomorphic to  $M$ . In particular  $H \oplus B$  is separable which in turn implies  $H$  is separable since over valuation domains, summands of separable modules are again separable (see [5]). It now follows directly from the theorem above that  $H$  is isomorphic to  $A$  and so the latter is again minimal.  $\square$

For the general notion of *basic submodule* of an  $R$ -module  $M$  we refer to [4], [5]. We recall that, when  $M$  is torsion-free, a basic submodule  $B$  of  $M$  is any submodule which is maximal with respect to the properties:

- (1)  $B$  is a direct sum of uniserial modules;
- (2)  $B$  is a pure submodule of  $M$ .

In the torsion-free setting, basic submodules always exist and are unique, up to isomorphism. The *basic rank* of  $M$  is defined to be the rank of its basic submodules; it plays a role somewhat analogous to that played by the  $p$ -rank of a torsion-free Abelian group.

In this context we have a result analogous to Ó hÓgáin's result (see [11]) that a torsion-free Abelian group with  $p$ -rank at most 1 for all primes  $p$ , is minimal.

We shall need a preliminary lemma, apparently not yet openly stated in the literature.

**Lemma 1.8.** *If  $B$  is a basic submodule of the torsion-free  $R$ -module  $M$ , then  $M = B + PM$ .*

*Proof.* Pick an arbitrary  $x \in M$ ,  $x \notin B$ . Since  $B$  is pure in  $M$ , we have  $Rx \cap B = 0$ . Then  $Rx \oplus B$  is not pure in  $M$ , by the defining properties of basic submodules. Now  $Rx \oplus B$  not pure amounts to the existence of  $y \in M$  and  $t \in P$  such that  $ty \in Rx \oplus B \setminus P(Rx \oplus B)$ . Let  $ty = rx + b$ , where  $r \in R$  and  $b \in B$ . Then  $r \notin P$ ; otherwise  $b \in PB$ , since  $B$  is pure, whence  $ty \in P(Rx \oplus B)$ , contrary to our assumption. It follows that  $r$  is a unit, so that  $x = r^{-1}(-b + ty) \in B + PM$ . Since  $x$  was arbitrary, we get the desired conclusion  $M = B + PM$ .  $\square$

**Theorem 1.9.** *Let  $M$  be a torsion-free  $R$ -module with basic rank 1. Then  $M$  is minimal.*

*Proof.* Suppose that there exists a proper submodule  $H$  of  $M$  such that  $M/H$  is finitely presented. Then, by Proposition 1.2, we may write  $M = X + H$ , where  $X$  is free and pure in  $M$ . Since  $M$  has basic rank 1, it follows that necessarily  $X$  is a basic submodule of  $M$ , whence  $X$  is cyclic, say  $X = Rx$ . Let  $\text{Ann}(x + H) = tR$ . We want to show that, necessarily,  $H = tM$ , whence  $M \cong H$  follows at once.

We readily see that  $tM \subseteq H$ , since  $tM = t(Rx + H)$  and  $tx \in H$ .

We want to verify the opposite inclusion  $H \subseteq tM$ . First we show that  $H \subseteq PM$ . Indeed, since  $Rx$  is basic in  $M$ , by the preceding lemma we have  $M = Rx + PM$ . Now assume, for a contradiction, that there exists  $h \in H \setminus PM$ . We may write

$$h = ax + qm$$

for suitable  $a \in R$ ,  $q \in P$ ,  $m \in M$ . Then  $a$  is a unit, since  $h \notin PM$ . Moreover  $m = bx + h'$ , where  $b \in R$ ,  $h' \in H$ . It follows that  $(a + qb)x \in H$ , which is impossible since  $a + qb$  is a unit and  $x \notin H$ . We may now verify that  $H \subseteq tM$ . Take any  $h \in H$ . Since the basic submodule of  $M$  is free of rank 1, the purification  $\langle h \rangle_*$  of  $h$  in  $M$  is also free. Accordingly, there exists  $q \in R$  such that  $h \in qM \setminus qPM$ . Therefore we may write

$$h = q(cx + h''), \quad c \in R, h'' \in H.$$

Since  $h'' \in PM$  and  $h \notin qPM$ , we get that  $c$  is a unit. Then  $qcx \in H$  implies  $qx \in H$ , whence  $q \in tR$ . It follows that  $h \in tM$ , as desired.  $\square$

## 2. NON-MINIMAL MODULES.

The purpose of the present section is to show how to construct non-minimal indecomposable modules of finite rank over largely general valuation domains. However, we will see that the important class of Nagata valuation domains constitutes a noticeable exception.

We begin with a pair of lemmas valid for commutative domains.

**Lemma 2.1.** *Let  $A$  be a commutative ring,  $\mathfrak{M}_1, \dots, \mathfrak{M}_k$  pairwise distinct maximal ideals of  $A$ , and  $M$  an  $A$ -module such that  $\mathfrak{M}_i M \neq M$  for all  $i \leq k$ . Then there exists  $z \in M$  such that  $z \notin \bigcup_{i \leq k} \mathfrak{M}_i M$ .*

*Proof.* Since  $\mathfrak{M}_j \not\supseteq \bigcap_{i \neq j} \mathfrak{M}_i$  for any  $j \leq k$ , we get  $M = \mathfrak{M}_j M + (\bigcap_{i \neq j} \mathfrak{M}_i) M$ , whence, in particular,  $\mathfrak{M}_j M \not\supseteq (\bigcap_{i \neq j} \mathfrak{M}_i) M$ . It follows that for every  $j \leq k$  there exists  $z_j \in (\bigcap_{i \neq j} \mathfrak{M}_i) M \setminus \mathfrak{M}_j M$ . It is now clear that the element  $z = z_1 + \dots + z_k$  fulfills our requirements.  $\square$

The following lemma and its proof are inspired by [1], Theorem 5.9 (a), page 52.

**Lemma 2.2.** *Let  $R$  be a commutative domain and let  $A$  be a commutative torsion-free  $R$ -algebra with  $\mathfrak{M}$  a maximal ideal of  $A$ . If  $M$  is a torsion-free  $R$ -module such that: (1)  $\text{End}_R(M) = A$  and (2)  $M/tM$  is a finitely generated  $R$ -module for some  $t \in R \cap \mathfrak{M}$ , then  $\mathfrak{M}M \neq M$ .*

*Proof.* Assume for a contradiction that  $\mathfrak{M}M = M$ . If  $N = M/tM$ , we have  $\mathfrak{M}N = N$ , as well. Since  $N$  is a finitely generated  $A$ -module, a classical result shows that  $(1+y)N = 0$ , for some  $y \in \mathfrak{M}$  (see e.g. [8]). It follows that  $(1+y)M \subseteq tM$ . Since  $M$  is a torsion-free  $R$ -module, we infer that  $(1+y)/t$  is a well-defined endomorphism of  $M$ , so that  $1+y \in tA \subseteq \mathfrak{M}$  and hence we get the desired contradiction  $1 \in \mathfrak{M}$ .  $\square$

**Theorem 2.3.** *Let  $R$  be a valuation domain and  $A$  a free  $R$ -algebra of rank  $n \geq 2$  containing only finitely many maximal ideals  $\mathfrak{M}_1, \dots, \mathfrak{M}_k$ ,  $k \geq 1$ . Suppose that there exists a torsion-free finite rank  $R$ -module  $M$  satisfying the following conditions: (1)  $\text{End}_R(M) = A$  and (2) there exists  $t \in P$  such that  $M/tM$  and  $A/tA$  are isomorphic as  $R$ -modules. Then if  $z \in M$  is such that  $z \notin \bigcup_{i \leq k} \mathfrak{M}_i M$ , the module  $N = Rz + tM$  is non-minimal. Moreover, if  $A$  is an integral domain then both  $M$  and  $N$  are indecomposable.*

*Proof.* It is clear that  $P \subset \mathfrak{M}_i$ , for every  $i \leq k$ . Also observe that the existence of  $z$  is ensured by Lemmas 2.1 and 2.2; indeed  $\mathfrak{M}_i M < M$ , for every  $i \leq k$ , since  $M/tM \cong A/tA \cong (R/tR)^n$  is a finitely generated  $R$ -module.

Let us first show that  $\text{Ann}_R(z + tM) = tR$ , so that  $tM$  has finite index in  $N$ . Suppose not and assume that  $sz \in tM$  with  $t/s \in P$ . It follows that  $z \in (t/s)M \subset PM \subset \mathfrak{M}_i M$  – contradiction.

We now verify that  $N$  is non-minimal, by proving that  $N \not\cong M \cong tM$ . Assume, for a contradiction, that there exists  $f \in \text{End}_R(M) = A$  such that  $f(M) = N$ . Now  $f \notin \mathfrak{M}_i$  for all  $i \leq k$ , since  $z \in f(M) \setminus \bigcup_{i \leq k} \mathfrak{M}_i M$ . Therefore  $f$  is a unit of  $A$  and  $M = N$ . It follows that

$$M/tM = N/tM \cong Rz/(tM \cap Rz)$$

is a cyclic  $R$ -module. We have reached the desired contradiction, since  $M/tM \cong A/tA \cong (R/tR)^n$  is a direct sum of  $n \geq 2$  cyclic modules.

Finally, assume that  $A$  is an integral domain. Then  $M$  is trivially indecomposable. However,  $N$  is also indecomposable. Suppose  $\pi : N \rightarrow N$  is an idempotent in  $\text{End}_R(N)$ . Then the composition  $t\pi : M \rightarrow N$  can be regarded as an endomorphism of  $M$ , say  $f$ . Thus  $f^2 = tf$  in  $\text{End}_R(M) = A$ , whence either  $f = 0$  or  $f = t1_A$ . In the first case we get  $f(N) = 0$ , whence  $\pi = 0$ , since  $N$  is torsion-free. If  $f = t1_A$ ,

then for any  $u \in N$  we get  $tu = f(u) = t\pi(u)$ . By torsion-freeness we infer that  $\pi$  is the identity on  $N$ . Thus  $\text{End}_R(N)$  contains only trivial idempotents and  $N$  is indecomposable.  $\square$

Before proving our next theorem, let us make some remarks. Assume that the field of quotients  $Q$  of  $R$  is not algebraically closed. Then there exists a non-trivial finite field extension  $L = Q[x]$  of  $Q$ , where we may clearly assume that  $x$  is integral over  $R$ . Consider the integral closure  $D$  of  $R$  in  $L$ ; by classical results on integral closures (for instance, see [3]), we know that  $D$  has only finitely many maximal ideals. Since  $x$  is integral over  $R$ , we have  $D \supseteq R[x]$ . The equality does not hold, in general. However,  $R[x]$  has only finitely many maximal ideals, since  $D$  is integral over  $R[x]$  and the ‘lying over’ property holds (see Theorem 44 of [8]).

In the remainder of this section we shall denote by  $\hat{Q}$ ,  $\hat{R}$ ,  $\hat{A}$  the completions as  $R$ -modules of  $Q$ ,  $R$ ,  $A$ , respectively.

**Theorem 2.4.** *Let  $R$  be a valuation domain such that  $Q$  is not algebraically closed and  $[\hat{Q} : Q] \geq 6$ . Then there exist non-minimal indecomposable torsion-free  $R$ -modules of finite rank.*

*Proof.* Since  $Q$  is not algebraically closed there exists, as noted above, a non-trivial finite-dimensional extension  $L = Q[x]$  of  $Q$  such that  $R[x]$  is not local and contains only finitely many maximal ideals. Set  $A = R[x]$ , and let  $n = [L : Q] \geq 2$ . Then  $A = R \oplus Rx \oplus \dots \oplus Rx^{n-1}$  is a free  $R$ -module.

Our goal is to construct an  $R$ -module  $M$  satisfying the conditions (1) and (2) of Theorem 2.3.

In this situation we are in a position to apply the techniques of [6] (see also [9]). We follow the notation of those papers. By Theorem 1 of [6] (see also Lemma 1 of [9]), for a suitable choice of  $\alpha, \delta \in \hat{R} \setminus R$ , the torsion-free  $R$ -module

$$M = \langle A1_A + A\alpha + A\delta \rangle_* \subset \hat{A}.$$

is such that  $\text{End}_R(M) = A$  (here the symbol  $\langle \rangle_*$  denotes the purification in  $\hat{A}$ ).

Fix  $t \in P$ ; to complete the proof we have to show that  $M/tM \cong A/tA$ . Consider firstly any  $z = b_1 1_A + b_2 \alpha + b_3 \delta \in A1_A + A\alpha + A\delta$ , where  $b_i \in A$ . For every  $r \in P$  there exist  $a_r, d_r \in R$  such that  $\alpha - a_r \in r\hat{R}$  and  $\delta - d_r \in r\hat{R}$ . If we set  $\lambda_r = b_1 + b_2 a_r + b_3 d_r \in A$ , it follows that

$$z - \lambda_r 1_A \in r\hat{A} \cap M = rM.$$

Now an arbitrary element of  $M$  may be written into the form  $z/r$ , with  $z = b_1 1_A + b_2 \alpha + b_3 \delta \in A1_A + A\alpha + A\delta$ , where the  $b_i \in A$  and  $r \in P$  are suitably chosen. In a similar way as above, we write  $\lambda_{rt} = b_1 + b_2 a_{rt} + b_3 d_{rt} \in A$ . Then  $z - \lambda_{rt} 1_A \in rt\hat{A}$ , and from this relation we first get  $\lambda_{rt}/r 1_A \in \hat{A}$ , whence  $\mu_{rt} = \lambda_{rt}/r \in A$ . Therefore we also see that

$$z/r \equiv \mu_{rt} 1_A \pmod{t\hat{A} \cap M = tM}.$$

Since  $z/r \in M$  was arbitrary, we conclude that the map  $\phi : A \rightarrow M/tM$  defined by  $\phi(\mu) = \mu(1_A + tM)$ , for  $\mu \in A$ , is an epimorphism. Let  $Q(A)$  be the field of fractions of  $A$ . If  $\mu 1_A \in tM \subset t\hat{A}$  then  $\mu/t \in \hat{A} \cap Q(A) = A$ , whence  $\mu \in tA$ . Thus we see that  $tA$  is the kernel of  $\phi$ , so that  $M/tM$  and  $A/tA$  are isomorphic as  $A$ -modules, and therefore isomorphic as  $R$ -modules, as well.  $\square$

The fact that the non-minimal modules furnished by the above theorem are indecomposable is particularly interesting in the light of the forthcoming Theorem 3.5. In fact in that result we will produce easy examples of non-minimal decomposable modules of rank two over any not almost maximal valuation domain.



It is worth noting that  $Q$  is not algebraically closed when  $R$  is not Henselian. Consequently Theorem 2.4 applies to any non-Henselian valuation domain  $R$  satisfying the degree condition.

Our next aim is to show that the preceding results do not hold, in general, if we deal with Henselian valuation domains. We will provide examples of non-maximal valuation domains, which are even discrete valuation domains, such that all their torsion-free modules of finite rank are minimal.

We consider the important class of discrete valuation domains called ‘‘Nagata valuation domains’’ in [17]. These are discrete valuation rings  $R$  of rank one such that  $[\hat{Q} : Q] = p^k$ , where  $p > 0$  is the characteristic of  $Q$ ,  $k$  is a positive integer, and, as above,  $\hat{Q}$  denotes the completion of  $Q$  in the topology of the valuation. These type of discrete valuation domains were first constructed in Nagata’s book [10], Example E33, page 207. They are Henselian, since, by construction,  $\hat{Q}$  is a purely inseparable extension of  $Q$ . Since they are not complete, they are, of course, not maximal,

However, note that Theorem 2.4 applies to Nagata valuation domains, whenever, in the above notation,  $p^k \geq 6$ .

Let  $R$  be a Nagata valuation domain such that  $[\hat{Q} : Q] = 2$ . Then it was shown in [17] (see also [2] for generalizations) that the finite-rank torsion-free indecomposable modules all have rank  $\leq 2$  and moreover those of rank 2 are all isomorphic ([17], Theorem 8). We are now in a position to establish the following result.

**Proposition 2.5.** *Let  $R$  be a Nagata valuation domain such that  $[\hat{Q} : Q] = 2$ . Then all torsion-free  $R$ -modules of finite rank are minimal.*

*Proof.* Let  $M$  be a torsion-free  $R$ -module of finite rank, and let  $H$  be any submodule of  $M$  of finite index. Thus we may write  $M = Z + H$ , where  $tZ \subseteq H$ , for a suitable  $t \in R$ , so that  $tM \subseteq H$ . Since  $R$  is a discrete valuation domain, we conclude that  $M$  and  $H$  are quasi-isomorphic, in the classical sense (see, for instance, [1]). In particular, by the results in [1],  $M$  and  $H$  have the same rank and basic rank (called  $p$ -rank in the classical setting). As shown in [17],  $M$  and  $H$  are direct sums of indecomposable submodules of rank  $\leq 2$ . We write  $M = F \oplus M_1$  and  $H = G \oplus H_1$ , where  $F, G$  are free modules, and  $M_1, H_1$  are direct sums of rank-two indecomposable modules. Let  $\text{rk } F = f$ ,  $\text{rk } G = g$ ,  $\text{rk } M_1 = 2h$ ,  $\text{rk } H_1 = 2k$ . Then the  $p$ -ranks of  $M_1, H_1$  are  $h, k$ , respectively. The equalities of the ranks and  $p$ -ranks yield the equations  $f + 2h = g + 2k$  and  $f + h = g + k$ , from which it follows that  $f = g$  and  $h = k$ . Therefore we at once get  $F \cong G$ . Moreover  $M_1$  and  $H_1$  have the same number of indecomposable rank-two direct summands. Since indecomposable rank-two modules are all isomorphic, we have  $M_1 \cong H_1$  as well. We conclude that  $M \cong H$ , as desired.  $\square$

Our final results in this section show that over certain discrete valuation domains the so-called Baer-Specker module:  $\mathcal{P} = \prod_{\aleph_0} R$  is not minimal. Note that  $\mathcal{P}$  has infinite rank. We are presently assuming that  $R$  is a discrete valuation domain. So, if  $R$  is complete the product  $\mathcal{P}$  will, of course, be minimal by Corollary 1.6. Clearly then it is necessary to assume that  $R$  is not complete but there are technical reasons relating to the combinatorial nature of our proof, which necessitate some cardinality restrictions. So suppose that  $R$  is a discrete valuation domain of cardinality  $\lambda$  with maximal ideal  $pR$  and that the residue class field  $R/pR$  has cardinality  $\mu$ . We shall call the following cardinal inequality, the residue class field cardinality condition:

$$\lambda^{\aleph_0} < \mu^{\mu^{\aleph_0}}.$$

Notice that  $\mu^{\mu^{\aleph_0}} = 2^{\mu^{\aleph_0}}$  and that if  $\lambda \leq 2^{\aleph_0}$  then this condition is always satisfied.

**Theorem 2.6.** *If  $R$  is a non-complete discrete valuation domain satisfying the residue class field cardinality condition, then  $\mathcal{P} = \prod_{\aleph_0} R$  is not minimal.*

*Proof.* Since  $R$  is not complete it is a slender module – see Example XVI 6.10 in [5] – and so the algebra of endomorphisms of  $\mathcal{P}$  is, as an  $R$ -module, isomorphic to  $\prod_{\aleph_0} \bigoplus_{\aleph_0} R$ ; in particular it has cardinality  $\lambda^{\aleph_0}$ . Consequently there are at most  $\lambda^{\aleph_0}$  submodules of  $\mathcal{P}$  which are isomorphic to  $\mathcal{P}$ .

Now consider the module  $\mathcal{P}/p\mathcal{P} \cong \prod_{\aleph_0} R/pR$ . This is a vector space over the residue class field  $R/pR$  and as such is isomorphic to  $\bigoplus_{\kappa} R/pR$  for some cardinal  $\kappa$ . Since  $\prod_{\aleph_0} R/pR$  is the dual space of  $\bigoplus_{\aleph_0} R/pR$ , its dimension  $\kappa$  is well known to be  $\mu^{\aleph_0}$ . Fix a one-dimensional summand, say,  $\mathcal{P}/p\mathcal{P} = \langle e \rangle \oplus V$  where  $V$  is a  $R/pR$ -space of dimension  $\kappa$ . It is well known that the number of direct complements for  $\langle e \rangle$  is equal to the dimension of the dual space of  $V$  i.e. there are  $\mu^{\kappa} = \mu^{\mu^{\aleph_0}}$  subspaces  $V_{\alpha} (\alpha < \mu^{\kappa})$  with  $\mathcal{P}/p\mathcal{P} = \langle e \rangle \oplus V_{\alpha}$ . Let  $H_{\alpha}$  be the pre-image of  $V_{\alpha}$  so that  $H_{\alpha}/p\mathcal{P} = V_{\alpha}$ . Thus the family  $\{H_{\alpha} : \alpha < \mu^{\kappa}\}$  is a family of  $\mu^{\mu^{\aleph_0}}$  submodules of  $\mathcal{P}$  each of which is of finite index in  $\mathcal{P}$ . By assumption  $\mu^{\mu^{\aleph_0}} > \lambda^{\aleph_0}$ , so not all of these submodules  $H_{\alpha}$  can be isomorphic to  $\mathcal{P}$ . Hence  $\mathcal{P}$  is not minimal as required.  $\square$

**Corollary 2.7.** *If  $R$  is a discrete valuation domain of non-measurable cardinality  $\lambda$  and  $R$  is not complete, then the product  $\prod_{\lambda} R$  is not minimal.*

*Proof.* Since  $\lambda$  is not measurable, the slenderness properties used above will still hold, giving the cardinality of the endomorphism algebra of the product  $\prod_{\lambda} R$  to be  $\lambda^{\lambda} = 2^{\lambda}$ . But now the number of submodules of finite index in  $\prod_{\lambda} R$  will be  $\mu^{\mu^{\lambda}} \geq 2^{2^{\lambda}} > \lambda^{\lambda}$  and the result follows as above.  $\square$

### 3. DIRECT SUMMANDS AND DIRECT SUMS.

We have already noted in Corollary 1.6 that a direct summand of a separable minimal module is again minimal.

It is worth noting and easily seen that a direct summand of a non-minimal module may be minimal. Just take any valuation domain  $R$  admitting a non-minimal  $R$ -module  $M_1$ . If  $F$  is a free  $R$ -module of finite rank, then  $M = M_1 \oplus F$  is not minimal (see the proof of the next proposition). Of course, the direct summand  $F$  of  $M$  is a minimal module.

**Proposition 3.1.** *Let the  $R$ -module  $M$  be of the form  $M = G \oplus F$ , where  $F$  is free of finite rank. Then  $M$  minimal implies  $G$  minimal.*

*Proof.* Suppose for a contradiction that  $M$  is minimal and  $G$  is not minimal. Then there is a submodule  $H$  of finite index in  $G$  which is not isomorphic to  $G$ . However the submodule  $H \oplus F$  is of finite index in  $M$ . Then one has  $H \oplus F \cong M \cong G \oplus F$  and this implies  $H \cong G$  since  $F$  is finitely generated, and finitely generated modules over a valuation domain have the substitution, and hence the cancellation, property (see [5], Corollary V.8.3).  $\square$

We will soon see that the converse of the preceding proposition is false, whenever  $R$  is not almost maximal.

In the case of a Henselian domain we can say somewhat more.

**Proposition 3.2.** *Let  $R$  be a Henselian valuation domain and let  $M$  be a minimal  $R$ -module of finite rank. Then the indecomposable direct summands of  $M$  are minimal.*

*Proof.* Recall that Vámos ([16]) proved that indecomposable torsion-free modules of finite rank over a Henselian valuation domain have local endomorphism rings. Let us write  $M = M_1 \oplus \cdots \oplus M_k$ , where the  $M_i$  are indecomposable. This decomposition is unique, up to isomorphism, since  $\text{End}_R(M_i)$  is local for all  $i \leq k$ , and we may apply Azumaya's Theorem. Let us now assume, by contradiction, that  $M_1$  (say) is non-minimal. Let  $H_1$  be a submodule of finite index of  $M_1$  such that  $H_1 \not\cong M_1$ . It is then clear that  $H = H_1 \oplus M_2 \oplus \cdots \oplus M_k$  is a submodule of  $M$  of finite index. Then  $M \cong H$ . Since  $M_2, \dots, M_k$  have local endomorphism rings, by iterated use of the substitution property (see [5] or [1], Ch. 8) we get  $M_1 \cong H_1$ . We have thus reached the desired contradiction.  $\square$

In the general situation one can derive a closure property under the operation of direct sums for the class of minimal modules provided one is willing to impose some restrictions.

**Theorem 3.3.** *If  $M = F \oplus X$ , where  $F$  is a finite rank free  $R$ -module and  $X$  is a minimal  $R$ -module with  $\text{Ext}_R^1(X, F)$  torsion-free, then  $M$  is minimal.*

*Proof.* Suppose that  $H$  is of finite index in  $M$ . Then since  $F$  is free of finite rank, the quotient  $(H + F)/H$  is a finitely generated submodule of  $M/H$  and so by Proposition 1.1,  $H$  is of finite index in  $H + F$ . It follows from the usual Noether isomorphism that  $H \cap F$  is of finite index in  $F$ , and hence we have  $F \cong (H \cap F)$ .

Since  $(H + F)/H$  is finitely generated and  $M/H$  is finitely presented torsion,  $M/(H + F)$  is finitely presented and so  $(H + F)/F$  is of finite index in  $M/F$ . But  $M/F \cong X$  and so  $(H + F)/F \cong X$  since  $X$  is minimal. Thus  $H/(H \cap F) \cong X$ . Moreover  $M/H$  is finitely presented so there is an element  $r \in R$  such that  $rM \leq H$ . Thus  $rM + (H \cap F) = r(F \oplus X) + (H \cap F) = rX \oplus (H \cap F)$ . But now  $(H \cap F) \leq rH + (H \cap F) \leq rX \oplus (H \cap F)$  and so  $rH + (H \cap F)$  splits over  $(H \cap F)$ . It follows from Lemma I.5.6 in [5], that this is equivalent to the extension  $0 \rightarrow (H \cap F) \rightarrow H \rightarrow H/(H \cap F) \rightarrow 0$  being in the kernel of the mapping induced on  $\text{Ext}_R^1(H/(H \cap F), H \cap F)$  by multiplication by the scalar  $r$  i.e. it is a torsion element of the module of extensions and this latter is just  $\text{Ext}_R^1(X, F)$  since  $H/(H \cap F) \cong X$  and  $F \cong (H \cap F)$ . However, by assumption,  $\text{Ext}_R^1(X, F)$  is torsion-free and so the extension  $0 \rightarrow (H \cap F) \rightarrow H \rightarrow H/(H \cap F) \rightarrow 0$  must be the null element, i.e.  $H$  splits as  $H = (H \cap F) \oplus Y$ , where  $Y \cong H/(H \cap F) \cong X$ . It follows that  $H \cong M$  and thus  $M$  is minimal as required.  $\square$

Since  $\text{Ext}_R^1(X, F) \cong \prod \text{Ext}_R^1(X, R)$  when  $F$  is free of finite rank, we can easily deduce the simple:

**Corollary 3.4.** *If  $M = F \oplus X$ , where  $F$  is free of finite rank and  $X$  is minimal with  $\text{Ext}_R^1(X, R) = 0$  then  $M$  is again minimal.*

**Remark 2.** In the previous theorem and corollary the only use made of  $F$  was that it was finitely generated. This was used to show that  $H \cap F$ ,  $(H + F)/F$  were of finite index in  $F$ ,  $M/F$ , respectively. We shall see shortly that for discrete valuation domains the above results can be significantly generalized.

Our next result shows the crucial fact that in general the direct sum of minimal modules may fail to be minimal. As a by-product, we will see that one cannot drop the condition relating to the vanishing of  $\text{Ext}$  in the above results. Even in the simplest possible case where  $F$  is free of rank 1, the direct sum of a minimal module and  $R$  need not be minimal; in fact this phenomenon always occurs when  $R$  is not an almost maximal valuation domain.

For the convenience of the reader, we recall some notions and results that are explained in full detail in [15] and [14], or, more briefly, in [5].

Let  $J$  be an ideal of the valuation domain  $R$ . Let us assume that  $R/J$  is not complete in the topology of its ideals. Then, necessarily,  $J$  is a v-ideal, that is  $J = \bigcap_{r \notin J} rR$ . Thus the set of ideals  $\mathcal{B} = \{rR/J : r \notin J\}$  forms a basis of neighborhoods of 0 for the ideal topology of  $R/J$ . Since  $R/J$  is not complete, there exists a Cauchy net, with respect to  $\mathcal{B}$ , with no limit in  $R/J$ . We may assume it to have the form  $\{u_r + J : r \notin J\}$ , where all the  $u_r$  are units of  $R$ .

For  $J$  an ideal of  $R$ , we denote by  $J^{-1}$  its inverse, namely  $J^{-1} = (R : J) = \{a \in Q : aJ \leq R\}$ . We have  $J^{-1} = \langle r^{-1} : r \notin J \rangle$ . Note that, if  $J$  is a non-zero v-ideal, then  $J^{-1}$  is a non-principal fractional ideal which is, by Proposition 1.4 (iii), a minimal  $R$ -module.

We also recall that  $R$  is not almost maximal exactly if there exists a non-zero ideal  $J$  such that  $R/J$  is not complete.

**Theorem 3.5.** *Let  $R$  be a valuation domain which is not almost maximal. Then direct sums of minimal  $R$ -modules may not be minimal. Specifically, if the non-zero ideal  $J$  is such that  $R/J$  is not complete, then the  $R$ -module  $R \oplus J^{-1}$  is not minimal.*

*Proof.* Let us first note that, since  $J$  is a non-zero v-ideal, then  $J^{-1}$  is a fractional ideal, whence, in particular,  $J^{-1}$  is minimal. Since  $R/J$  is not complete, we may choose a family of units  $\{u_r : r \notin J\}$  of  $R$  in the way described above. Consider the vector space  $Qx \oplus Qy$ , where  $x, y$  are indeterminates. By the results in [15] the following submodule of  $Qx \oplus Qy$  is indecomposable of rank two:

$$H = \langle x, z_r = r^{-1}(x + u_r y) : r \notin J \rangle;$$

moreover  $Rx$  is a basic submodule of  $H$ . Now choose  $t \in J$  and note that the  $R$ -module  $M = Rt^{-1}x \oplus J^{-1}y$  clearly contains  $H$  and is isomorphic to  $R \oplus J^{-1}$ . Since  $H$  is indecomposable, it is not isomorphic to  $M$ . To complete the proof, it remains to show that  $H$  has finite index in  $M$ . We will show that  $M = Rt^{-1}x + H$  and  $M/H \cong R/tR$ . Take  $r^{-1}y \in J^{-1}y$ , where  $r$  is any element of  $R$  not in  $J$ . By direct computation we see that  $r^{-1}y = u_r^{-1}(-(t/r)t^{-1}x + z_r) \in Rt^{-1}x + H$  (note that  $t/r \in P$ ). This suffices to show that  $M \subseteq Rt^{-1}x + H$ . Since  $x \in H$ , it is clear that  $\text{Ann}(t^{-1}x + H) \supseteq tR$ . Assume now that  $at^{-1}x \in H$  for a suitable  $a \in R$ . Since  $Rx$  is pure in  $H$ , we have  $x \notin PH$ , and therefore  $x \in ta^{-1}H$  implies  $ta^{-1} \notin P$ , so that  $at^{-1} \in R$  and  $a \in tR$ . Thus it follows that  $\text{Ann}(t^{-1}x + H) \supseteq tR$ , whence equality holds, and we get the desired conclusion.  $\square$

The requirement in the preceding theorem that the valuation domain  $R$  is not almost maximal is necessary. If  $R$  is a discrete valuation domain and thus is automatically almost maximal, we can derive a far-reaching generalization of Theorem 3.3 and Corollary 3.4 above.

**Theorem 3.6.** *Let  $R$  be a discrete valuation domain and let  $G$  and  $X$  be minimal  $R$ -modules such that  $\text{Ext}_R^1(X, G)$  is torsion-free. Then  $M = G \oplus X$  is minimal.*

*Proof.* As noted in the Remark after Corollary 3.4, the proof of this result will follow immediately from Theorem 3.3 (replacing  $F$  with  $G$ ) if we can show that  $H \cap G, (H + G)/G$  are of finite index in  $G$  and  $M/G$  respectively, where  $H$  is an arbitrary submodule of finite index in  $M$ . Since  $R$  is now Noetherian, the notions of finitely presented and finitely generated coincide, as a submodule of a finitely generated module is again finitely generated. But then  $(H + G)/H \leq M/H$  is finitely presented and so  $H \cap G$  is of finite index in  $G$ . Moreover  $M/(H + G)$  as the quotient of two finitely presented modules is again finitely presented and so  $(H + G)/G$  is of finite index in  $M/G$  as required.  $\square$

**Corollary 3.7.** *Let  $R$  be a discrete valuation domain and  $G$  a minimal  $R$ -module. If  $F$  is a free  $R$ -module of arbitrary rank, then  $M = G \oplus F$  is minimal.*

*Proof.* Since  $F$  is free,  $\text{Ext}_R^1(F, G) = 0$ , and hence is trivially torsion-free.  $\square$

In special circumstances the preceding results enable one to characterize summands of minimal groups.

**Corollary 3.8.** *Let  $R$  be a discrete valuation domain and let  $M$  be a  $R$ -module of the form  $M = G \oplus F$ , where  $F$  is free of finite rank. Then  $M$  is minimal if and only if  $G$  is minimal.*

*Proof.* One implication follows from the corollary above. The converse follows from Proposition 3.1.  $\square$

#### REFERENCES

- [1] D. Arnold, *Finite Rank Torsion Free Abelian Groups and Rings*, Lecture Notes in Math., Springer-Verlag, Berlin - Heidelberg - New York, 1982.
- [2] D. Arnold and M. Dugas, *Indecomposable modules over Nagata valuation domains*, Proc. Amer. Math. Soc. 122 (1994), 689–696.
- [3] N. Bourbaki, *Commutative Algebra, Chapters 1–7*, Springer-Verlag, Berlin - Heidelberg - New York, 1989.
- [4] L. Fuchs and L. Salce, *Modules over Valuation Domains*, Marcel Dekker, New York and Basel 1985.
- [5] L. Fuchs and L. Salce, *Modules over Non-Noetherian Domains*, Math. Surveys and Monographs, Vol. 84, AMS, Providence, Rhode Island 2001.
- [6] B. Goldsmith and W. May, *The Krull-Schmidt problem for modules over valuation domains*, J. Pure Appl. Algebra 140 (1999), 57–63.
- [7] B. Goldsmith and S. Ó hÓgáin, *On Torsion and Mixed Minimal Abelian Groups*, Rocky Mountain J. Mathematics 32 (2002) 1451–1465.
- [8] I. Kaplansky, *Commutative Rings*, The University of Chicago Press, Chicago and London, 1974.
- [9] W. May and P. Zanardo, *Non-henselian valuation domains and the Krull-Schmidt property for torsion-free modules*, Forum Math. 10 (1998) 249–257.
- [10] M. Nagata, *Local Rings*, Wiley/Interscience, New York, 1962.
- [11] S. Ó hÓgáin, *On Torsion-Free Minimal Abelian Groups*, submitted 2004.
- [12] P. Ribenboim, *Equivalent forms of Hensel’s lemma*, Expo. Math. 3 (1985) 3–24.
- [13] D. J. S. Robinson and M. Timm, *On Groups that are Isomorphic with Every Subgroup of Finite Index and Their Topology*, J. London Math. Soc. 57 (1988) 91–104.
- [14] L. Salce and P. Zanardo, *On two-generated modules over valuation domains*, Archiv der Math. 46 (1986) 408 - 418.
- [15] L. Salce and P. Zanardo, *Rank-two torsion-free modules over valuation domains*, Rend. Sem. Mat. Univ. Padova 80 (1988) 175 - 201.
- [16] P. Vámos, *Decomposition problems for modules over valuation domains*, J. London Math. Soc. 41 (1) (1990) 10–26.
- [17] P. Zanardo, *Kurosch invariants for torsion-free modules over Nagata valuation domains*, J. Pure Appl. Algebra 82 (1992), 195–209.

DUBLIN INSTITUTE OF TECHNOLOGY, KEVIN STREET, DUBLIN 8, IRELAND AND  
 DIPARTIMENTO DI MATEMATICA PURA E APPLICATA, VIA BELZONI 7, 35131 PADOVA, ITALY  
*E-mail address:* `brendan.goldsmith@dit.ie` and `pzanardo@math.unipd.it`