2010-01-01

Influence of Cultivation Conditions on the Production of a Thermostable Extracellular Lipase from *Amycolatopsis Mediterranei* DSM 43304

Dharmendra Dheeman

Technological University Dublin, dharmendra.dheeman@student.dit.ie

Jesus Maria Frias

Technological University Dublin, Jesus.Frias@tudublin.ie

Gary Henahan

Technological University Dublin, gary.henahan@tudublin.ie

Follow this and additional works at: <https://arrow.tudublin.ie/schfsehart>

Part of the [Biochemistry Commons](#)

Recommended Citation

Dheeman D.S., Frias J.M, Henahan, G.T.M. (2010) Influence of Cultivation Conditions on the Production of a Thermostable Extracellular Lipase from *Amycolatopsis Mediterranei* DSM 43304. *Journal of Industrial Microbiology & Biotechnology*, 37, (1),pp. 1-17. doi:10.1007/s10295-009-0643-7 doi:10.1007/s10295-009-0643-7

This Article is brought to you for free and open access by the School of Food Science and Environmental Health at ARROW@TU Dublin. It has been accepted for inclusion in Articles by an authorized administrator of ARROW@TU Dublin. For more information, please contact arrow.admin@tudublin.ie, aisling.coyne@tudublin.ie, gerard.connolly@tudublin.ie.

Funder: ABBEST Research Scholarship, Technological University Dublin (PB 03557/2007).

**Influence of cultivation conditions on the production of a thermostable extracellular
lipase from *Amycolatopsis mediterranei* DSM 43304**

Dharmendra S. Dheeman, Jesus M. Frias and Gary T. M. Henehan*

School of Food Science and Environmental Health, Dublin Institute of Technology, Cathal
Brugha Street, Dublin 1, Ireland.

*Corresponding Author:

E-mail: gary.henehan@dit.ie

Tel: +353 1 4024408

Fax: +353 1 4024495

Abstract

Among several lipase producing actinomycete strains screened, *Amycolatopsis mediterranei* DSM 43304 was found to produce a thermostable, extracellular lipase. Culture condition and nutrient source modification studies involving carbon sources, nitrogen sources, incubation temperature and medium pH were carried out. Lipase activity of 1.37 ± 0.103 IU/ml of culture medium was obtained in 96 h at 28 °C and pH 7.5 using linseed oil and fructose as carbon sources and a combination of phytone peptone and yeast extract (5:1) as nitrogen sources. In optimal culture conditions the lipase activity was enhanced 12-fold with a 2-fold increase in lipase specific activity. The lipase showed maximum activity at 60 °C and pH 8.0. The enzyme was stable between pH 5.0–9.0 and temperatures up to 60 °C. Lipase activity was significantly enhanced by Fe^{3+} and strongly inhibited by Hg^{2+} . Li^+ , Mg^{2+} and PMSF significantly reduced lipase activity, whereas other metal ions had no significant effect at 0.01 M concentration. *A. mediterranei* DSM 43304 lipase exhibited remarkable stability in the presence of a wide range of organic solvents at 25% (v/v) concentration for 24h. These features render this novel lipase attractive for potential biotechnological applications in organic synthesis reactions.

Keywords: *Amycolatopsis mediterranei*; Screening; Organic solvent-tolerant; Thermostable; Lipase

1. Introduction

Lipases (triacylglycerol acylhydrolases E.C.3.1.1.3) are serine hydrolases of considerable physiological significance and industrial potential [46]. Interest in lipases has greatly increased in recent years, mainly because they present a broad range of biotechnological applications due to their unique characteristics of substrate specificity, regio-specificity and chiral-selectivity [41]. Many microorganisms, including bacteria, yeast, and fungi, have been shown to secrete lipases during their growth on hydrophobic substrates [25]. Among lipases of various origins, those from bacteria show the highest degree of versatility, reactivity and stability in the catalysis of reactions in organic solvents [19]. Lipase applications range from the hydrolysis of fats in wastewaters to the synthesis of chiral pharmaceutical intermediates [42]. These applications often require harsh reaction conditions *e.g.* high temperatures may be required to either favour stereoselectivity or solubilise high melting point lipids [17]. Biocatalysis with lipases is often carried out in organic solvents to promote synthetic reactions by reducing water activity. Many lipases lose activity in organic solvents, and therefore, there is an ongoing interest in lipases that are solvent-tolerant [7]. Despite the advantages of biocatalysis in organic solvent based systems, the catalytic activities of enzymes in these systems are typically much lower than in aqueous solutions [43]. Currently, since microbial lipases do not have the desirable combination of thermostability and stability in both hydrophobic and hydrophilic organic solvents, the search for new lipases is required [31, 33]. There are two main strategies for obtaining lipases with improved properties either protein engineering of currently known lipases [40] or the search for novel lipolytic activities in previously unexplored

microorganisms [11]. As each industrial application requires specific properties of lipases, there is still an active interest in finding novel lipases for specific applications.

To date, a large number of lipases from filamentous fungi, yeasts and unicellular bacteria have been extensively studied, both from the biochemical and from genetic point of view [8]. However, despite their high biotechnological potential for the production of secondary metabolites and enzymes, the actinomycetes have not been widely studied for lipase production [1, 30, 50].

The present paper deals with the screening of lipase producing actinomycete strains and the culture conditions for optimum enzyme production by a selected strain of *Amycolatopsis mediterranei* DSM 43304. Medium composition, initial pH, temperature, and time of incubation were examined for the optimization of lipase production. Lipase characteristics with respect to the optimal temperature and pH for both activity and stability are examined. We also characterize the activity and stability of the lipase in the presence of various organic solvents, metal ions, detergents and inhibitors. To our knowledge, the present work is the first report of an organic solvent-tolerant lipase from the genus *Amycolatopsis*.

2. Materials and methods

2.1 Materials

Analytical reagent grade chemicals were purchased from commercial sources at the highest purity. Unless mentioned otherwise, all culture media and chemicals used were from Sigma (Dublin, Ireland). Phytone peptone was obtained from BBL Microbiology Systems (Cockeysville, MD, USA) and Bacto-peptone was obtained from Difco Laboratories (Detroit, MI, USA). Natural oils were purchased from the local retail in Dublin.

2.2 Microorganisms

Actinomycete strains were obtained from Divisional Culture Collection, School of Biology, Newcastle University, UK.

2.3 Maintenance of microorganisms

Actinomycete strains were grown on GYM agar slants (g/l: glucose 4.0 g; yeast extract 4.0 g; malt extract 10 g; CaCO₃ 2.0 g; agar 12.0 g; pH 7.2). The working stock cultures were maintained and stored on GYM slants at 4 °C.

2.4 Culture conditions

The composition of basal medium used was (g/l): NaNO₃ 0.5 g; KCl 0.5 g; MgSO₄·7H₂O 0.5 g; KH₂PO₄ 2.0 g; yeast extract 1.0 g; and Bacto-peptone 5.0 g. The pH was adjusted to 7.2 with 1 M NaOH or 1 M HCl. Then 1.0% (v/v) olive oil was added. Media were sterilized for 15 min at 121 °C at 15 psi. Submerged microbial cultures were incubated in 250 ml Erlenmeyer flasks containing 50 ml of basal medium with 5 ml inoculum on a rotary shaker (130 rpm) at 28 °C.

2.5 Rhodamine B agar screening

The primary screening for the detection of lipolytic activity on solid media was carried out on rhodamine B agar (RBA) as described by Kouker and Jaeger [29] with some modifications. The growth medium containing 0.9% (w/v) peptone water, 0.25 % (w/v) of yeast extract, 2% (w/v) of agar, was adjusted to pH 7.2, autoclaved and cooled to 60 °C. Then, filter sterilized rhodamine B stock solution (1.0 mg/ml) in distilled water was added to a substrate lipoidal emulsion to yield a final concentration of 0.001% (w/v). The substrate lipoidal emulsion consisted of 1.5% (w/v) olive oil with 0.25% (v/v) Tween 80 in distilled water that was sterilized by autoclaving. The resulting mixture of lipoidal emulsion with growth medium (1:10) was vigorously stirred to emulsify for 15 min. The medium

was allowed to stand for 10 min at 60 °C to reduce foaming before pouring 20 ml of medium into plastic petri plates. Fresh RBA plates were spot inoculated with 72 h-old actinomycete cultures in GYM broth (g/l: glucose 4.0 g; yeast extract 4.0 g; malt extract 10 g; pH 7.2) and incubated at 28 °C for 6 days. The plates with visible growth were UV irradiated (350 nm). Lipase production was identified as orange fluorescence under UV light.

2.6 Screening in submerged fermentation broth

For screening in submerged cultivation conditions, 50 ml of basal medium in 250 ml Erlenmeyer flasks was inoculated with 5 ml, 72 h-old actinomycete culture in GYM broth and incubated at 28 °C on a reciprocal shaker (130 rpm). After 96 h, samples were processed for lipase activity assay. One ml of culture was centrifuged at 10,000 ×g, at 4 °C, for 10 min to obtain a cell free supernatant. The clear supernatant was filtered through 0.2 µm filter before lipase activity assay.

2.7 Spectrophotometric *p*-NPP assay

Lipase activity was quantitatively assayed in cell free supernatant using *p*-nitrophenyl palmitate (*p*-NPP) as substrate. This assay was performed as described by Winkler and Stuckman [55] with some modifications. A stock solution of *p*-NPP was freshly prepared in 2-propanol at a concentration of 0.3% (w/v). This solution (Solution A) was subjected to 3 min sonication (135W, 42 kHz, Branson 5510E-MT). Then, 900 µl of 1:20 dilution of the substrate stock solution A in solution B (0.1% (w/v) gum arabic, 0.4% (v/v) Triton X100 in distilled water) with 50 µl of appropriate buffer were preincubated for 2 min at the assay temperature before adding 50 µl of enzyme sample. This mixture was incubated at the assay temperature for 10 min, and the reaction was terminated by addition of 2 ml of 0.2 M

Na₂CO₃ solution. Released *p*-nitrophenol (*p*-NP) was immediately determined by measuring the absorbance at 410 nm in a Unicam UV-VIS spectrophotometer (Model UV2000E, Cambridge, UK). Appropriate blanks were used to subtract the absorbance corresponding to the reaction mixture other than that produced by the specific hydrolysis of *p*-NPP. The molar extinction coefficient of *p*-NP ($\epsilon_{410\text{nm}} = 16,900 \text{ M}^{-1} \text{ cm}^{-1}$) was estimated from the absorbance of standard solutions of *p*-NP. One international unit of lipase activity was expressed as the amount of enzyme liberating 1 μM of *p*-NP per minute under the conditions of the assay.

2.8 Biomass concentration analysis

After centrifugation at 10,000 $\times g$, at 10 °C, for 10 min and washing the pellet in 0.9% (w/v) NaCl solution, the pellet of 5 ml suspension sample was dried to a constant weight at 80 °C for 48 h and the dry biomass weight was determined gravimetrically.

2.9 Statistical analysis

Data were analyzed using analysis of variance. In all these cases the analyses were conducted using SPSS (version 15.0) using the procedure of general linear model (Tukey test). The level of tested significance was at $p \leq 0.05$.

3. Time course of lipase production by *A. mediterranei* DSM 43304

Time course of lipase production was studied in the basal medium using shake flask cultures. A 10% (v/v) of 72 h-old inoculum grown in GYM broth was added to 50 ml medium, in a 250 ml Erlenmeyer flask and incubated at 130 rpm on a rotary shaker, at 28 °C. Samples were analyzed at 24 h intervals to determine pH, dry biomass and lipase activity in the culture supernatant. The effects of the cultivating conditions in shake flask experiments are commonly investigated by subjecting the microorganism to different

environmental conditions [56]. In the present work the effect of initial pH and incubation temperature on the culture was studied using shake flask cultures at different temperatures (20–45 °C) and initial values of pH (5.0–9.0). The effect of inoculum size on lipase production was investigated by varying culture inoculum size from 2% to 12% of total volume. All experiments were carried out at least in triplicate.

4. Nutritional factors affecting lipase production by *A. mediterranei* DSM 43304

The general procedure for cultivation was as follows: 10% (v/v) of 72 h-old inoculum grown in GYM broth was inoculated into 50 ml of culture medium and incubated at 28 °C for 96 h. The culture was harvested by centrifugation at 10,000 ×g, 4 °C for 10 min. The cell free supernatant was filtered (0.2 µm filter, Millipore) before spectrophotometric determination of lipase activity. Each experiment was carried out in triplicate.

4.1 Effect of inducers on lipase production

To determine the effects of substrate related compounds, the olive oil in the basal medium was substituted with natural oils (jojoba, corn, cottonseed, grapeseed, groundnut, linseed, rapeseed, soybean and sunflower) at 1.0% (v/v) concentration.

4.2 Effect of surfactants on lipase production

The following detergents were added to the basal medium as lipase inducers, replacing olive oil, at 0.5% (w/v) concentration: Span 40, Span 65, Span 80, Tween 20, Tween 21, Tween 40, Tween 80 and Triton X100.

4.3 Effect of carbon source additives on lipase production

In order to elucidate the effect of carbon source additives on lipase production, olive oil in the basal medium was combined with 1.0% (w/v) of the following sugars: arabinose,

dextrin, fructose, galactose, glucose, lactose, mannitol, maltose, maltotetraose, mannose, raffinose, rhamnose, sorbitol, sucrose, starch and xylose.

4.4 Effect of organic nitrogen sources on lipase production

To assess the effects of organic nitrogen source in combination with yeast extract on lipase production, Bacto-peptone in the basal medium, was substituted with phytone peptone, yeast extract, corn steep liquor, beef extract, skim milk, wheat peptone, fish peptone, tryptone, casein hydrolysate, casein, and wheat gluten, each at a concentration of 0.5% (w/v).

4.5 Effect of phytone peptone and yeast extract on lipase production

Phytone peptone with yeast extract was the best nitrogen source for *A. mediterranei* DSM 43304 lipase production. The influence of the amount of phytone peptone and yeast extract in basal medium was tested by varying their concentrations in the basal medium.

4.6 Effect of inorganic nitrogen sources on lipase production

In order to assess the effects of inorganic nitrogen sources on lipase production, NaNO₃ in the basal medium was substituted with: NH₄Cl, (NH₄)₂SO₄, (NH₄)₂HPO₄, NH₄H₂PO₄, CH₃COONH₄, NH₄NO₃ and urea, each at 0.05% (w/v).

4.7 Effect of metal ions on lipase production

Metal ions individually and in combination were tested for their effects on optimal lipase production in basal medium. Mg²⁺, Ca²⁺ and Fe³⁺ salts were incorporated into the basal medium at 0.05% (w/v) concentration. Also combinations of Mg²⁺ and Ca²⁺, Ca²⁺ and Fe³⁺, and Mg²⁺ and Fe³⁺ were added at individual final concentration of 0.025% (w/v) to assess their cumulative effect on lipase production.

5. Characterization of *A. mediterranei* DSM 43304 lipase

5.1 Effect of temperature activity and stability

Lipase activity was measured at various temperatures (30–80 °C) under standard assay conditions. Thermal stability of the enzyme was investigated by preincubating the enzyme at various temperatures (30–80 °C) for 3 h. The samples were then assayed for residual lipase activity under standard assay conditions.

5.2 Effect of pH on activity and stability

The effect of pH on lipolytic activity was determined in the following buffers (all at 50 mM): HCl–KCl (pH 2.0), sodium citrate (pH 3.0), succinate–NaOH (pH 4.0 and 5.0), sodium phosphate (pH 6.0 and 7.0), Tris–HCl (pH 8.0 and 9.0) and glycine–NaOH (pH 10.0). The optimum pH obtained was used for investigation of thermostability and other parameters. These buffers were used to determine pH stability of the crude lipase preparation. A mixture (1:1) of crude enzyme and buffers (0.1 M) was incubated for 24 h at 4 °C and the standard enzyme assay described previously was performed. The residual activities were calculated by comparison with the activity in 50 mM Tris–HCl buffer, pH 8.0, without pre-incubation.

5.3 Effect of organic solvents on lipase stability

The effect of various polar and non-polar organic solvents with different $\log P$ values on crude lipase stability was investigated. One ml of organic solvent was added to 3.0 ml of cell free supernatant and incubated at 30 °C, while shaking at 200 rpm for 24 h to ensure the continuous mixing of enzyme and the solvents. The enzyme stability was expressed as the remaining activity relative to the control without solvent.

5.4 Effect of metal ions and effector molecules on lipase stability

226 The effect of metal ions and effector molecules on lipase activity was studied at pH 8.0 by
227 incubating the enzyme in presence of 1 mM of metal ions (Ag^+ , Ba^{2+} , Co^{2+} , Li^+ , Mn^{2+} , Ni^{2+} ,
228 Pb^{2+} , Ca^{2+} , Fe^{3+} , Cu^{2+} , Zn^{2+} , Mg^{2+} and Hg^{2+}) and effectors (PMSF, EDTA, SDS, NH_4^+ and
229 urea). Incubation was carried out at 60 °C for 10 min and assayed for lipase activity.

230 Residual lipase activity was calculated as a percentage of that without metal ions/effectors.

231 **6. Results and discussion**

232 **6.1 Screening of strains for lipase production**

233 Figure 1 shows rhodamine B agar screening for lipase production by actinomycete strains.

234 Except for *Amycolatopsis coloradensis* DSM 44225 and *Streptomyces aureoverticillatus*

235 NRRL B-3326, all actinomycete strains showed brilliant pink-red/orange fluorescence on

236 UV irradiation of RBA plates. Lipase screening in basal medium using shake flask cultures

237 showed presence of lipolytic activity in cell free supernatants prepared from actinomycete

238 culture broths (Table 1). Of 18 actinomycetes strains, 7 showed more than 0.05 IU/ml at 96

239 h. Among these, *A. rubida* DSM 44637, *S. rochei* DSM 40231 and *S. griseus* subsp.

240 *griseus* DSM 40236 produced the highest activities at 0.149 ± 0.017 IU/ml, 0.141 ± 0.004

241 IU/ml and 0.116 ± 0.010 IU/ml, respectively.

242 Actinomycetes are Gram-positive bacteria with a remarkable genetic repertoire for

243 producing secondary metabolites and enzymes. Despite their high biotechnological

244 potential, the actinomycetes have not been widely investigated for lipase activity [8, 13,

245 50]. Only a few studies have been reported on their esterase and lipolytic activities [6].

246 Large et al. described lipase activity associated with the cells of different *Streptomyces* sp.,

247 which were found to be induced and enhanced by the presence of a lipid substrate in the

248 fermentation medium [30]. Gandolfi et al. carried out carboxylesterase screening of

Streptomyces strains on solid media using tributyrin, triolein and Tween 60 as substrate and evaluated their cell bound and extracellular hydrolytic activities [18]. Cardenas et al. isolated novel actinomycetes and fungal strains and demonstrated their lipolytic activity by employing screening techniques on solid and liquid media using agar plates supplemented with emulsified olive oil and tributyrin [8]. The present investigation identified a number of actinomycetes producing significant extracellular lipolytic activities.

6.2 Effect of culture conditions on lipase production by *A. mediterranei* DSM 43304

The preliminary characterization of extracellular lipases from actinomycete strains in terms of pH optimum, temperature optimum and thermostability identified the lipase from *A. mediterranei* DSM 43304 as the most thermostable and analysis by zymography indicated presence of a single lipase in extracellular culture broth (data not shown). Therefore, *A. mediterranei* DSM 43304 was selected for further characterization and various culture parameters were studied to improve lipase production.

6.2.1 Time course of lipase production

It has been reported that the lipase synthesis of *S. exfoliatus* M11 and *S. coelicolor* A3(2) is growth phase dependent [47]. Therefore, the time course of lipase synthesis in basal medium by *A. mediterranei* DSM 43304 was monitored by measurement of lipase activity, dry biomass and pH. Figure 2 shows that substantial lipase production commenced at 24 h and reached a maximum at 96 h. Further incubation lead to increase in lipase activity with a slow decrease in the lipase specific activity (data not shown). Loss of lipase specific activity may be due to secretion of other proteins at the late logarithmic phase leading to an apparent decrease in lipase specific activity. Swift et al. reported that once cell densities have reached certain threshold level, generally in the late logarithmic phase, the expression

of genes encoding exoproteins and secretion system is induced [49]. There was a shift in pH from 7.20 ± 0.02 to 8.02 ± 0.057 during the first 24 h of incubation, rising to pH 8.96 ± 0.11 at 96 h.

6.2.2 Effect of inoculum size on lipase production

Low inoculum density may give insufficient biomass causing reduced product formation, whereas a higher inoculum may produce too much biomass leading to poor product formation [37]. Increased enzyme production was observed with the increase in inoculum size and showed maximum enzyme activity (0.105 ± 0.003 IU/ml) and biomass (5.8 ± 0.6 mg/ml) production at 96 h with 10% inoculum (Table 2). Further increases in inoculum size resulted in decreased enzyme synthesis, probably due to nutrient limitation.

6.2.3 Effect of incubation temperature on lipase production

Temperature may affect lipase production [34]. Submerged fermentation was carried out at 20–45 °C with 10% inoculum for 96 h to evaluate the effect of incubation temperatures on growth and enzyme production. Maximum enzyme and biomass production was observed in the mesophilic range at 28–35 °C (Figure 3). The optimal temperature determined for lipase production by *A. mediterranei* DSM 43304 (28 °C) is comparable to those of *S. erythraea*, *S. clavuligerus* [30] and *Pseudomonas aeruginosa* PseA [45].

6.2.4 Effect of initial medium pH on lipase production

The pH of the culture broth was found to be one of the most critical environmental parameters affecting the growth and enzyme production by *A. mediterranei* DSM43304. The results showed maximum biomass and lipase production at initial medium pH of 7.5 (Figure 4). Lipase activity dropped significantly at alkaline and acidic pH of 9.0 and 5.0, respectively. The optimum pH (7.5) is close to optimum pH of 7.0 for lipase production by

295 *Candida* sp. [52] but is lower than that of other reported lipase-producing organisms.
296 *Bacillus mycoides* showed optimal lipase production at pH 8.0. [53] whereas, maximal
297 lipase production by *A. terreus* was observed at an initial medium pH of 9.0 [22]. The pH
298 change observed during growth of the organism may affect the enzyme stability in the
299 medium [23].

300 **6.2.5 Effect of inducers on lipase production**

301 Natural oils as carbon sources had different effects on lipase production. The results (Table
302 3) indicate that all lipidic sources supported lipase activity, ranging from 0.014 ± 0.001
303 IU/ml to 0.128 ± 0.004 IU/ml. The highest lipase production (0.128 ± 0.004 IU/ml) and
304 biomass (16.0 ± 1.4 mg/ml) was found using linseed oil followed by sunflower oil ($0.104 \pm$
305 0.004 IU/ml). Soybean oil gave the lowest lipase activities. There are very few examples in
306 literature where linseed oil has been used as a lipid source for lipase production. Linseed oil
307 was tested as one of the oils for lipase production by *Burkholderia cepacia* [44] and in case
308 of *P.camembertii* Thom PG-3 it was found to be the second best inducer for lipase
309 production [51]. Natural oils such as soybean, corn, sunflower, olive, palm and cotton seed
310 oils, amongst others, are cited as inducers of lipase production, comprising at times, the
311 sole source of carbon in the medium [34, 38, 44, 51].

312 Surfactants as lipase inducers in the medium did not enhance lipase production when
313 compared to olive oil (Table 3). Similar effects were reported on lipase production in
314 *Rhizopus* sp. BTNT-2 [3] and *Yarrowia lipolytica* [15]. Surfactants do not always increase
315 lipolytic enzyme production [35] and their effect varies with microorganism, surfactant
316 type and its concentration [15, 52]. However, the addition of surfactant to the culture

medium has been shown to increase the secretion of lipolytic enzymes in a number of microorganisms, attributable to alteration of cell permeability leading to increased protein secretion or to surface effects on cell bound enzymes [52]. A wide variety of surfactants like Tweens, Triton, SDS, PEG and gum arabic have been studied by different investigators [15, 35, 52].

6.2.6 Effect of carbon source additives on lipase production

The effect of carbon source in the basal medium on lipase production by *A. mediterranei* DSM43304 is shown in Table 4. Lipase production and biomass was significantly enhanced with the addition of sugars to the basal medium in most cases. Fructose gave the highest activity of 0.467 ± 0.007 IU/ml with biomass of 11.2 ± 0.9 mg/ml, whereas lactose produced the highest biomass of 14.9 ± 0.4 mg/ml with lipase activity of 0.399 ± 0.021 IU/ml. Lipase production was increased 3-fold with lactose, sorbitol, maltose and xylose as carbon source additives. However, addition of arabinose, dextrin, and sucrose had no significant effect on lipase production and showed significant decrease in biomass. In contrast, addition of rhamnose showed increased biomass with no significant improvement in lipase activity. A range of different carbon sources (i.e. carbohydrates, alcohols, acids, lipids) have been reported to support both growth of lipolytic enzyme producers and lipase/esterase production [4, 22]. Although lipidic carbon sources seem to be generally essential for obtaining a high enzyme yield, some authors have indicated good results in the absence of fats and oils [34], whereas in some cases, a mixture of compounds has been proposed as optimum carbon source [14, 18]. Similar to the present study, fructose as carbon source was found to significantly increase lipase activity in *Issatchenkia orientalis* [11]. Lactose was reported to support high to moderate lipase production in various

microorganisms [26], which is similar to the present study where lactose led to a significant increase in lipase production.

6.2.7 Effect of nitrogen sources on lipase production

Nitrogen sources, including organic nitrogen and inorganic nitrogen sources play an important role in the biosynthesis of lipase [51]. Lipase production and biomass were found to be highest with phytone peptone (0.131 ± 0.021 IU/ml) and yeast extract (0.104 ± 0.005 IU/ml) as organic nitrogen source followed by Bacto-peptone (0.071 ± 0.002 IU/ml) and cornsteep liquor (0.061 ± 0.006 IU/ml) (Table 5). Significant biomass was produced with beef extract (8.4 ± 0.7 mg/ml) and tryptone (8.1 ± 0.9 mg/ml) as nitrogen source but the lipase activities obtained were low. Wheat gluten gave the lowest lipase (0.017 ± 0.001 IU/ml) yield followed by casein and casein hydrolysate. Similar studies were carried out by other investigators to find the best nitrogen source for lipase production [3, 22]. Soybean meal was reported to be the best nitrogen source for lipase production by *P. camembertii* PG-3 [51]. Higher lipase production was reported using yeast extract as nitrogen source for *Saccharomyces cerevisiae* [48]. Contrary to the present results, casein and corn gluten were the best sources for lipase production by *A. terreus* [22].

Phytone peptone with yeast extract was identified as the best nitrogen sources for lipase production by *A. mediterranei* DSM 43304. The effect of different concentrations of phytone peptone and yeast extract on lipase production is shown in Table 5. There was no significant difference in terms of lipase activities and biomass production by different concentrations of phytone peptone and yeast extract in the basal medium. By contrast, a similar study identified a combination of 0.6% tryptone and 0.2% yeast extract as the best nitrogen source for lipase production by *Bacillus* sp. strain 42 [16]. The effect of different

inorganic nitrogen sources showed $(\text{NH}_4)_2\text{HPO}_4$ as the best inorganic nitrogen source producing maximum lipase activity followed by NaNO_3 (Table 6). In agreement with the present observation, higher specific activity of *Burkholderia cepacia* lipase was observed in presence of $(\text{NH}_4)_2\text{HPO}_4$ as inorganic nitrogen source followed by KNO_3 [44]. Among the inorganic nitrogen sources tested for *Pseudomonas* sp. G6, NaNO_3 supported the best lipase production [27]. Urea showed the lowest lipase and biomass yield followed by NH_4Cl . Urea was also reported to be inhibitory for lipase synthesis by *P. camembertii* Thom PG-3 [51].

6.2.8 Effect of metal ions on lipase production

Besides physical and nutritional parameters, metal ions may play an important role in lipase production [39]. The effect of different metal ions and metal ion combinations on lipase production is shown in Table 4. Metal ions did not make any significant difference to lipase activity and biomass production. Contrary to the present results, iron was reported to be critical for the production of lipase by *Pseudomonas* sp. G6 [27] and Ca^{2+} was reported to have strong stimulatory effect on extracellular lipase production by *P. fluorescens* 2D [36]. Similarly, Ca^{2+} in presence of Mg^{2+} was reported to produce a significant increase in lipase production by *Burkholderia cepacia* [44].

6.2.9 Lipase production in modified medium

The extent of improvement in lipase activity yield by fermentation using the optimal culture parameters was investigated. The modified culture medium for lipase production by *A. mediterranei* DSM 43304 consisted of 1% (v/v) linseed oil, 1% (w/v) fructose, 0.05% (w/v) $(\text{NH}_4)_2\text{HPO}_4$, 0.05% (w/v) KCl, 0.05% (w/v) $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$, 0.2% (w/v) KH_2PO_4 , 0.5% (w/v) phytone peptone, 0.1% (w/v) yeast extract, initial pH 7.5 and the culture was

incubated at 28 °C for 96 h. The lipase activity produced increased from 0.108 ± 0.002 IU/ml (3.80 ± 0.14 IU/mg) in the basal medium to 1.372 ± 0.103 IU/ml (7.27 ± 0.52 IU/mg) in the modified medium. Thus, a 12-fold increase in lipase activity yield and 2-fold increase in lipase specific activity was achieved by employing newly formulated production medium.

7. Characterization of lipase

7.1 Effect of temperature on activity and stability

The effect of temperature on the activity and stability of lipase is shown in Figure 5. The lipase from *A. mediterranei* DSM 43304 was found to be quite thermostable with a temperature optimum of 60 °C at pH 8.0. It retained 90% of activity at 60 °C after 3 h incubation and had a half-life of more than 30 min at 70 °C. High thermal stability of lipolytic activity detected in this study may be useful for several industrial applications. Bacterial lipases generally have temperature optima in the range 30–60 °C. However, reports exist of bacterial lipases with optima both in lower and higher ranges [24]. Thermal stability of a lipase is obviously related to its tertiary structure and is influenced by environmental factors such as pH and the presence of metal ions [62]. A few thermostable lipases from actinomycetes have been reported. *S. fradiae* var. k11 was reported to produce a proteolysis-resistant lipase having a temperature optimum of 55 °C [60] and a thermostable lipase, most active at 50–60 °C, was purified from *S. rimosus* [1]. Two different lipases from *S. coelicolor* A3(2) were characterized showing temperature optima at 20–30 °C (SCO1725) and 45–55 °C (SCO7513) [12].

7.2 Effect of pH on activity and stability

Figure 6 shows the effect of pH on lipolytic activity and stability. The lipase was found to be active over the pH range 5.0–8.0. Maximal lipolytic activity was at pH 8.0 and showed retention of 88%, 98%, 95% and 89% activity at pH 5.0, 6.0, 7.0 and 9.0, respectively. The pH stability profile showed the highest stability at pH 8.0 and 9.0 with 96% residual activity. The activity was also stable at pH 6.0 and 7.0 showing 94% residual activity after 24 h. The stability of the enzyme in acidic and alkaline pH suggests its usefulness in industrial applications. Generally, lipases of bacterial origin have neutral or alkaline pH optima with the exception of *P. fluorescence* SIK W1 lipase which has an acidic pH optimum of 4.8 [2]. Most of the lipases characterized from actinomycetes so far show alkaline pH optima. *S. rimosus* lipase was reported to show a pH optimum range of 9.0–10.0 [1] and *S. fradiae* var. k11 lipase was most active at pH 9.8 [60]. Similarly, *S. coelicolor* A3(2) lipases showed pH optima between 7.5–10.0 [12].

7.3 Effect of organic solvents on lipase stability

The effect of various organic solvents on the stability of *A. mediterranei* DSM 43304 lipase is shown in Table 6. The lipase exhibited considerable stability in the presence of polar solvents ($\log P < 0.3$) as well as non-polar hydrophobic solvents ($\log P$ 0.85–6.6) with significant activation observed in most cases. The highest degree of activation was in benzene and toluene. Lipase was significantly activated after 1 h in *p*-xylene and *n*-hexane showing a 61.9% and 56.7% increase in activity, respectively. Further incubation for 24 h led to 28.2% and 45% decrease in activation indicating destabilizing effects of these solvents. The lipase was least stable in pyridine and showed 49.8% residual activity after 24 h. Similarly, a destabilizing effect was seen with *t*-butanol, dodecane and DMSO. It has been shown that many enzymes retain activity in organic solvents [59] and have interesting

catalytic properties such as higher thermostability and altered stereoselectivity [54]. Despite many advantages of enzymatic reactions in organic solvents, in most cases the catalytic activity in organic solvents is orders of magnitude lower than in aqueous systems [10] because of diffusional limitations, changes in protein flexibility, or destabilization of the enzyme [28]. *A. mediterranei* DSM 43304 lipase was activated and stable in DMF, methanol, ethanol, 2-propanol and acetone. Though solvent stability has been reported for a few actinomycete lipases [5, 8, 32], the high stability and indeed, activation in polar solvents like methanol and ethanol has rarely been observed [8]. *S. rimosus* lipase was reported to show stimulation of activity upon addition of 2.5% (v/v) 1,4-dioxane, THF, acetone and DMF [32]. Similarly, *S. coelicolor* lipase was found to be stable with different water-miscible solvents and showed 50% increase in activity in acetone after 18 h and similar to the present study, significant loss of activity was observed with DMSO, while in contrast, DMF led to almost complete loss of activity [5]. It is reported that polar solvents strip off the essential water molecules from the active site of enzymes [21]. For this reason, use of polar solvents is avoided and hydrophobic solvents are more often employed in non-aqueous enzymology [20]. The polar solvent tolerant lipases therefore appear promising for catalysis in low water media. This property is a novel attribute of *A. mediterranei* DSM 43304 lipase, which has been reported in few cases [5, 32, 61]. The high stability and activation of *A. mediterranei* DSM 43304 lipase in a wide range of polar and non-polar organic solvents may make it useful for practical applications in synthetic biotransformation reactions.

7.4 Effect of metal ions and effectors on lipase stability

453 The effect of different metal salts and effector molecules on *A. mediterranei* DSM 43304
 454 lipase is shown in Table 7. The crude lipase was strongly inhibited by Hg^{2+} losing 80% of
 455 activity in 10 min suggesting it is able to alter enzyme conformation as has been reported
 456 for other lipases [9]. Li^+ and Mg^{2+} also significantly reduced enzyme activity. The other
 457 metal ions tested did not produce adverse effect on the activity of the enzyme. Compared to
 458 the control, Ca^{2+} did not cause significant reduction in lipase, even though Ca^{2+} has been
 459 reported to stabilize lipolytic activity [9, 20]. Cu^{2+} has been reported to be a strong inhibitor
 460 of lipase activity [19, 26], but no inhibition was observed with Cu^{2+} at a concentration of 1
 461 mM under the conditions tested. Similarly, *S. fradiae* var. k11 lipase was reported to retain
 462 99.4% lipase activity in the presence of 1 mM Cu^{2+} for 30 min [60]. Cu^{2+} did not
 463 significantly reduce the lipolytic activity of SCO1725 and SCO7513 from *S. coelicolor*
 464 A3(2) [12]. The lipase activity was unaffected by the metal-chelating agent EDTA
 465 indicating that *A. mediterranei* DSM 43304 lipase is probably not a metalloenzyme. The
 466 activities of lipases from *S. rimosus* and *S. fradiae* var. k11 have also been reported to be
 467 unaffected by EDTA [1, 60]. The effect of a serine inhibitor PMSF, at 1 mM concentration
 468 gave a 15% reduction in lipase activity possibly suggesting the presence of a hydrophobic
 469 lid hindering access to the catalytic site [12]. The extracellular lipase from *S. rimosus* was
 470 found to be marginally affected by 1 mM PMSF showing retention of 90% of residual
 471 activity [1]. Similarly, 89–91% residual activity was retained in presence of 1 mM PSMF
 472 by *S. coelicolor* A3(2) lipases [12]. The lipase showed 88% retention of activity in presence
 473 of 1 mM urea. SDS was found not to affect lipase activity. In agreement with the present
 474 study lipase activity was reported to be stable with 79% residual activity in the presence of
 475 1 mM SDS by *B. cepacia* lipase [57], whereas *Yarrowia lipolytica* lipase showed

significant inhibition of lipase activity with 1 mM SDS [58]. Contrary to the present results, lipase from *B. thermoleovorans* CCR11 was reported to be completely inhibited by 1 mM SDS in 1 h [9].

8. Conclusions

A lipolytic actinomycete strain, *Amycolatopsis mediterranei* DSM 43304 was identified as producing an extracellular organic solvent-tolerant thermostable lipase. A significant 12-fold increase in lipase activity and 2-fold increase in lipase specific activity was achieved by employing the newly formulated culture medium. The extracellular lipase was highly stable in a broad range of polar and non-polar organic solvents, with maximum enhancement of activity in non-polar solvents. The properties of thermostability, stability and enhancement of activity in organic solvents, and stability with a broad range of effector molecules shows that this lipase may be a promising enzyme for applications in a range of *in vitro* biotransformation reactions.

Acknowledgements

This work was supported by ABBEST Research Scholarship from Dublin Institute of Technology (PB 03557/2007). The authors want to express their thanks to Prof. Michael Goodfellow and Dr. Amanda L. Jones, School of Biology, Newcastle University, UK for providing the strains for this study.

References:

1. Abramic M, Lescic I, Korica T, Vitale L, Saenger W, Pigac J (1999) Purification and properties of extracellular lipase from *Streptomyces rimosus*. Enzym Microb Technol 25:522–529

2. Andersson RE, Hedlund GB, Jensson V (1979) Thermal inactivation of a heat resistant lipase produced by the psychrotrophic bacterium *Pseudomonas fluorescens*. J Dairy Sci 62:361–367
3. Bapiraju KVVS, Sujatha P, Ellaiah P, Ramana T (2005) Sequential parametric optimisation of lipase production by a mutant strain *Rhizopus* sp, BTNT-2. J Basic Microbiol 45:257–273
4. Benjamin S, Pandey A (1996) Optimization of liquid media for lipase production by *Candida rugosa*. Bioresour Technol 55:167–170
5. Bielen A, Cetkovic H, Long PF, Schwab H, Abramic M, Vujaklija D (2009) The SGNH-hydrolase of *Streptomyces coelicolor* has (aryl)esterase and a true lipase activity. Biochimie 91:390–400
6. Bormann C, Nikoleit K, Potgeter M, Tesch C, Sommer P, Goetz F (1993) Investigation of lipolytic enzymes from *Streptomyces*. In: DECHEMA-monographs. VCH, Weinheim, Germany, pp 237–247
7. Boutaiba S, Bhatnagar T, Hacene H, Mitchell DA, Baratti JC (2006) Preliminary characterisation of a lipolytic activity from an extremely halophilic archaeon, *Natronococcus* sp. J Mol Catal B: Enzym 41:21–26
8. Cardenas F, Castro MS, Sanchez-Montero JM, Sinisterra JV, Valmaseda M, Elson SW, Alvarez E (2001) Novel microbial lipases: catalytic activity in reactions in organic media. Enzym Microb Technol 28:145–154
9. Castro-Ochoa LD, Rodriguez-Gomez C, Valerio-Alfaro G, Ros RO (2005) Screening, purification and characterization of the thermoalkalophilic lipase

produced by *Bacillus thermoleovorans* CCR11. Enzyme Microb Technol 37:648–
654

10. Clark DS (2004) Characteristics of nearly dry enzymes in organic solvents:
implications for biocatalysis in the absence of water. Philos T Roy Soc B
359:1299–1307

11. Costas M, Deive FJ, Longo MA (2004) Lipolytic activity in submerged cultures
of *Issatchenkia orientalis*. Process Biochem 39:2109–2114

12. Cote A, Shareck F (2008) Cloning, purification and characterization of two lipases
from *Streptomyces coelicolor* A3(2). Enzym Microb Technol 42:381–388

13. Cruz H, Perez C, Wellington E, Castro C, Servin-Gonzalez L (1994) Sequence of
the *Streptomyces albus* G lipase-encoding gene reveals the presence of a
prokaryotic lipase family. Gene 144:141–142

14. Dalmau E, Montesinos JL, Lotti M, Casas C (2000) Effect of different carbon
sources on lipase production by *Candida rugosa*. Enzyme Microb Technol
26:657–663

15. Dominguez A, Deive FJ, Sanroman MA, Longo MA (2003) Effect of lipids and
surfactants on extracellular lipase production by *Yarrowia lipolytica*. J Chem
Technol Biotechnol 78:1166–1170

16. Eltaweel MA, Rahman RNZRA, Salleh AB, Basri M (2005) An organic solvent-
stable lipase from *Bacillus* sp. strain 42. Ann Microbiol 55:187–192

17. Ema T (2004) Rational strategies for highly enantioselective lipase-catalyzed
kinetic resolutions of very bulky chiral compounds: Substrate design and high-
temperature biocatalysis. Tetrahedron: Asymmetry 15:2765–2770

- 543 18. Gandolfi R, Marinelli F, Lazzarini A, Molinari F (2000) Cell-bound and
544 extracellular carboxylesterases from *Streptomyces*: hydrolytic and synthetic
545 activities. J App Microb 89: 870–875
- 546 19. Gao XG, Cao SG, Zhang KC (2000) Production, properties and application to
547 nonaqueous enzymatic catalysis of lipase from a newly isolated *Pseudomonas*
548 strain. Enzym Microb Technol 27:74–82
- 549 20. Gaur R, Gupta A, Khare SK (2008) Purification and characterization of lipase
550 from solvent tolerant *Pseudomonas aeruginosa* PseA. Process Biochem 43:1040–
551 1046
- 552 21. Gorman LAS, Dordick JS (1992) Organic solvents strip water off enzymes.
553 Biotechnol Bioeng 39:392–397
- 554 22. Gulati R, Saxena RK, Gupta R, Yadav RP, Davidson WS (1999) Parametric
555 optimisation of *Aspergillus terreus* lipase production and its potential in ester
556 synthesis. Process Biochem 35:459–464
- 557 23. Gupta R, Gigras P, Mohapatra H, Goswami VK, Chauhan B (2003) Microbial α -
558 amylases: a biotechnological perspective. Process Biochem 38:1599–1616
- 559 24. Gupta R, Gupta N, Rathi P (2004) Bacterial lipases: an overview of production,
560 purification and biochemical properties. Appl Microbiol Biotechnol 64:763–781.
- 561 25. Haba E, Bresco D, Ferrer C, Margues A, Busquets M, Manresa A (2000) Isolation
562 of lipase-secreting bacteria by deploying used frying oil as selective substrate.
563 Enzym Microb Technol 26:40–44

26. Kamini NR, Fujii T, Kurosu T, Iefuji H (2000) Production, purification and characterization of an extracellular lipase from *Cryptococcus* sp. S-2. *Process Biochem* 36:317–324
27. Kanwar L, Gogoi BK, Goswami P (2002) Production of a *Pseudomonas* lipase in *n*-alkane substrate and its isolation using an improved ammonium sulphate precipitation technique. *Bioresour Technol* 84:207–211
28. Klibanov AM (1997) Why are enzymes less active in organic solvents than in water? *Trends Biotechnol* 15:97–101
29. Kouker G, Jaeger KE (1987) Specific and sensitive plate assay for bacterial lipases. *Appl Environ Microbiol* 53:211–213
30. Large KP, Mirjalili N, Osborne M, Peacock LM, Zormpaidis V, Walsh M, Cavanagh ME, Leadlay PF, Ison AP (1999) Lipase activity in *Streptomyces*. *Enzym Microb Technol* 25:569–575
31. Lee M-H, Lee CH, Oh TK, Song JK, Yoon JH (2006) Isolation and characterization of a novel lipase from a metagenomic library of tidal flat sediments: evidence for a new family of bacterial lipases. *Appl Environ Microbiol* 72:7406–7409
32. Lescic I, Vukelic B, Majeric-Elenkov M, Saenger W, Abramic M (2001) Substrate specificity and effects of water-miscible solvents on the activity and stability of extracellular lipase from *Streptomyces rimosus*. *Enzym Microb Technol* 29: 548–553

33. Lima VMG, Krieger N, Mitchell DA, Baratti JC, Filippis I, Fontana JD (2004) Evaluation of the potential for use in biocatalysis of a lipase from a wild strain of *Bacillus megaterium*. J Mol Catal B: Enzym 31:53–61
34. Lin ES, Ko HC (2005) Glucose stimulates production of the alkaline-thermostable lipase of the edible basidiomycete *Antrodia cinnamomea*. Enzyme Microb Technol 37:261–265
35. Lin SF, Chiou CM, Tsai YC (1995) Effect of Triton X-100 on alkaline lipase production by *Pseudomonas pseudoalkaligenes* F-111. Biotechnol Lett 17:959–962
36. Makhzoum A, Knapp JS, Owusu RK (1995) Factor affecting growth and lipase production by *Pseudomonas fluorescens* 2D. Food Microbiol 12:277–290
37. Mudgetti RE (1986) In: Demain AL, Solmen NA (ed) Manual of industrial biotechnology. Washington, DC: American Society for Microbiology, pp 66–83
38. Muralidhar RV, Chirumamila RR, Marchant R, Nigam P (2001) A response surface approach for the comparison of lipase production from *Candida cylindracea* using two different carbon sources. Biochem Eng J 9:17–23
39. Obido FJC, Okereke UO, Oyeka CA (1995) Influence of culture conditions on the production of lipase of *Hendersonula toruloidea*. Bioresour Technol 54:81–83.
40. Ogino H, Ishikawa H (2001) Enzymes which are stable in the presence of organic solvents. J Biosci Bioeng 91:109–116
41. Pandey A, Benjamin S, Soccol CR, Nigam P, Krieger N, Soccol VT (1999) The realm of microbial lipases in biotechnology. Biotechnol Appl Biochem 29:119–131

42. Patel RN (2008) Synthesis of chiral pharmaceutical intermediates by biocatalysis. *Coord Chem Rev* 252:659–701
43. Pencreac'h G, Baratti JC (2001) Comparison of hydrolytic activity in water and heptane for thirty-two commercial lipase preparations. *Enzym Microb Technol* 28:473–479
44. Rathi P, Saxena RK, Gupta R (2001) A novel alkaline lipase from *Burkholderia cepacia* for detergent formulation. *Process Biochem* 37:187–192
45. Ruchi G, Anshu G, Khare SK (2008) Lipase from solvent tolerant *Pseudomonas aeruginosa* strain: production optimization by response surface methodology and application. *Bioresour Technol* 99:4796–4802
46. Schmid RD, Verger R (1998) Lipases: interfacial enzymes with attractive applications. *Angew Chem Int Ed Engl* 37:1608–1633
47. Servin-Gonzalez L, Castro C, Perez C, Rubio M, Valdez F (1997) bldA-dependent expression of the *Streptomyces exfoliatus* M11 lipase gene (lipA) is mediated by the product of a contiguous gene, lipR, encoding a putative transcriptional activator. *J Bacteriol* 179:7816–7826
48. Shirazi SH, Rahman SR, Rahman MM (1998) Production of extracellular lipases by *Saccharomyces cerevisiae*. *World J Microbiol Biotechnol* 14:595–597.
49. Swift S, Throup JP, Williams P, Salmond GP, Stewart GS (1996) Quorum sensing: a population-density component in the determination of bacterial phenotype. *Trend Biochem Sci* 21:214–219
50. Sztajer H, Maliszewka I, Wieczorek J (1988) Production of exogenous lipases by bacteria, fungi and actinomycetes. *Enzym. Microb. Technol* 10:492–497

- 631 51. Tan T, Zhang M, Xu J, Zhang J (2004) Optimization of culture conditions and
632 properties of lipase from *Penicillium camembertii* Thom PG-3. Process Biochem
633 39:1495–1502
- 634 52. Tan TW, Zhang M, Wang BW, Ying CH, Deng L (2003) Screening of high lipase
635 producing *Candida* sp. and production of lipase by fermentation. Process
636 Biochem 39:459–465
- 637 53. Thomas A, Mathew M, Valsa AK, Mohan S, Manjula R (2003) Optimisation of
638 growth conditions for the production of extracellular lipase by *Bacillus mycoides*
639 Ind J Microbiol 43:67–69
- 640 54. Wescott CR, Klivanov AM (1994) The solvent dependence of enzyme specificity.
641 Biochim Biophys Acta 1206:1–9
- 642 55. Winkler UK, Stuckman M (1979) Glycogen, hyaluronate and some other
643 polysaccharides greatly enhance the formation of exolipase by *Serratia*
644 *marcescens*. J Bacteriol 138:663–679
- 645 56. Yang FC, Huang HC, Yang MJ (2003) The influence of environmental conditions
646 on the mycelial growth of *Antrodia cinnamomea* in submerged cultures. Enzyme
647 Microb Technol 33:395–402
- 648 57. Yu L, Xu Y, Yu X (2009) Purification and properties of a highly enantioselective
649 L-menthyl acetate hydrolase from *Burkholderia cepacia*. J Mol Catal B Enzym
650 57:27–33
- 651 58. Yu MR, Qin SW, Tan TW (2007) Purification and characterization of the
652 extracellular lipase Lip2 from *Yarrowia lipolytica*. Process Biochem 42:384–391

59. Zaks A, Klivanov AM (1985) Enzyme-catalyzed processes in organic solvents.
Proc Natl Acad Sci U S A 82:3192–3196
60. Zhang Y, Meg K, Wang Y, Luo H, Yang P, Shi P, Wu N, Fan Y, Li J, Yao B
(2008) A novel proteolysis-resistant lipase from keratinolytic *Streptomyces*
fradiae var. k11 Enzym Microb Technol 42:346–352
61. Zhao L, Xu J, Zhao J, Pan J, Wang Z (2008) Biochemical properties and potential
applications of an organic solvent tolerant lipase isolated from *Serratia*
marcescens ECU1010. Process Biochem 43:626–633
62. Zhu K, Jutila A, Patkar EKJ, Svendsen A, Kinnunen PK (2001) Impact of the
tryptophan residues of *Humicola lanuginosa* lipase on its thermal stability. J
Biochim Biophys Acta 1547:329–338

Table 1. Lipase screening of actinomycete strains in basal medium:

Strain	Rhodamine B agar [*]	Lipase activity ^{**} (IU/ml)
<i>Amycolatopsis coloradensis</i> DSM 44225	—	0.046 ± 0.001 ^{abc}
<i>A. amakusaensis</i> NRRL B-3351	+	0.059 ± 0.004 ^{bc}
<i>A. fastidiosa</i> DSM 43855	+	0.040 ± 0.002 ^{abc}
<i>A. mediterranei</i> DSM 43304	+	0.086 ± 0.012 ^d
<i>A. rubida</i> DSM 44637	+	0.149 ± 0.017 ^f
<i>A. sulphurea</i> DSM 46092	+	0.062 ± 0.008 ^{cd}
<i>Nocardia araoensis</i> DSM 44729	+	0.031 ± 0.002 ^a
<i>N. higoensis</i> DSM 44732	+	0.037 ± 0.001 ^{abc}
<i>N. kruckzakiae</i> DSM 44877	+	0.033 ± 0.003 ^a
<i>Streptomyces amquistii</i> NRRL B-1685	+	0.046 ± 0.001 ^{abc}
<i>S. griseus</i> subsp. <i>griseus</i> DSM 40236	+	0.116 ± 0.010 ^e
<i>S. coelicolor</i> A3(2)	+	0.033 ± 0.001 ^a
<i>S. annulatus</i> NRRL B-2000	+	0.051 ± 0.002 ^{abc}
<i>S. arabicus</i> NRRL B-1733	+	0.030 ± 0.001 ^a
<i>S. aurantiogriseus</i> NRRL B-5416	+	0.038 ± 0.006 ^{abc}
<i>S. rochei</i> DSM 40231	+	0.141 ± 0.004 ^{ef}
<i>S. aureoverticillatus</i> NRRL B-3326	—	0.034 ± 0.004 ^{ab}
<i>S. althioticus</i> NRRL B-3981	+	0.037 ± 0.004 ^{ab}

Data are means ± standard deviations of three determinations. Column data followed by the same superscript

letter were not significantly different ($p \leq 0.05$; by Tukey test)

^{*}The RBA plates, after 6 days incubation at 28 °C, were exposed to UV (350 nm) to detect fluorescence:

orange fluorescence (+); no orange fluorescence (—).

^{**}Lipase activity in cell-free supernatant after 96 h growth in basal medium with 1% (v/v) olive oil as inducer.

Table 2. Effect of inoculum level on lipase production:

Inoculum volume (% v/v)	Lipase (IU/ml)	Biomass (mg/ml)	Final pH
2	0.055 ± 0.005 ^a	0.7 ± 0.4 ^a	8.66 ± 0.07
4	0.064 ± 0.002 ^{ab}	1.0 ± 0.3 ^a	8.82 ± 0.06
6	0.079 ± 0.010 ^{abc}	1.5 ± 0.1 ^a	8.60 ± 0.08
8	0.085 ± 0.009 ^{bcd}	4.8 ± 0.4 ^b	8.76 ± 0.04
10	0.105 ± 0.003 ^d	5.8 ± 0.6 ^b	8.77 ± 0.03
12	0.102 ± 0.005 ^{cd}	5.6 ± 0.7 ^b	8.73 ± 0.04

*The shake flask experiments were performed in basal medium for 96 h at 28 °C, 130 rpm.

**Data are means ± standard deviations of three determinations. Column data followed by the same superscript letter were not significantly different ($p \leq 0.05$; by Tukey test).

Table 3. Effect of inducers on lipase production:

Inducer	Lipase (IU/ml)	Biomass (mg/ml)	Final pH
Natural oils (1.0%, v/v)			
Control*	0.081 ± 0.004 ^f	6.0 ± 2.8 ^{ab}	7.90 ± 0.03
Joboba	0.071 ± 0.003 ^{ef}	4.0 ± 1.4 ^a	8.84 ± 0.07
Corn	0.052 ± 0.002 ^{cd}	4.5 ± 3.5 ^{ab}	8.71 ± 0.04
Cottonseed	0.060 ± 0.007 ^{de}	4.5 ± 0.7 ^{ab}	8.70 ± 0.28
Grapeseed	0.046 ± 0.001 ^{bc}	2.5 ± 0.7 ^a	8.67 ± 0.03
Groundnut	0.037 ± 0.001 ^b	4.5 ± 0.7 ^{ab}	8.64 ± 0.06
Linseed	0.128 ± 0.004 ^h	16.0 ± 1.4 ^b	7.01 ± 0.01
Rapeseed	0.013 ± 0.004 ^a	2.5 ± 0.7 ^a	8.66 ± 0.18
Soybean	0.014 ± 0.001 ^a	1.5 ± 0.7 ^a	8.70 ± 0.42
Sunflower	0.104 ± 0.004 ^g	11.5 ± 2.1 ^{bc}	7.81 ± 0.07
Surfactants (0.5 %, w/v)			
Control*	0.125 ± 0.006 ^c	5.5 ± 0.1 ^d	7.26 ± 0.03
Span 40	0.031 ± 0.003 ^{ab}	1.6 ± 0.8 ^{ab}	8.71 ± 0.03
Span 65	0.034 ± 0.001 ^b	1.6 ± 0.1 ^{ab}	8.68 ± 0.04
Span 80	0.063 ± 0.002 ^d	4.7 ± 0.8 ^{cd}	8.10 ± 0.21
Tween 20	0.056 ± 0.003 ^{cd}	1.5 ± 0.7 ^{ab}	8.75 ± 0.03
Tween 21	0.040 ± 0.004 ^{bc}	0.4 ± 0.1 ^a	6.99 ± 0.01
Tween 40	0.053 ± 0.003 ^{cd}	1.2 ± 0.1 ^{ab}	8.74 ± 0.03
Tween 80	0.067 ± 0.007 ^d	3.2 ± 0.1 ^{bc}	8.54 ± 0.03
Triton X-100	0.016 ± 0.002 ^a	0.2 ± 0.1 ^a	7.26 ± 0.06

The shake flask experiments were performed for 96 h at 28 °C, 130 rpm. Data are means ± standard

deviations of three determinations. Column data followed by the same superscript letter were not significantly different ($p \leq 0.05$; by Tukey test).

* Control refers to the basal medium with 1% (v/v) olive oil as lipase inducer.

Table 4. Effect of sugar additives and metal ions on lipase production:

Sugar additive/metal ion	Lipase activity (IU/ml)	Biomass (mg/ml)	Final pH
Sugar additive (1.0%, w/v)			
Contol*	0.132 ± 0.007 ^{ab}	7.3 ± 0.5 ^{abc}	8.66 ± 0.01
Arabinose	0.158 ± 0.013 ^{abc}	4.6 ± 0.3 ^a	8.90 ± 0.08
Dextrin	0.194 ± 0.042 ^{bcd}	4.7 ± 0.2 ^a	8.57 ± 0.07
Fructose	0.467 ± 0.007 ^j	11.2 ± 0.9 ^{gh}	8.65 ± 0.04
Galactose	0.349 ± 0.007 ^{fghi}	12.7 ± 0.2 ^{hi}	8.81 ± 0.03
Glucose	0.317 ± 0.019 ^{efgh}	10.7 ± 0.1 ^{defgh}	8.69 ± 0.03
Lactose	0.399 ± 0.021 ^{hij}	14.9 ± 0.4 ⁱ	8.72 ± 0.03
Mannitol	0.343 ± 0.035 ^{fghi}	10.9 ± 0.6 ^{fgh}	8.64 ± 0.03
Maltose	0.424 ± 0.071 ^{ij}	8.2 ± 0.4 ^{bcdef}	8.79 ± 0.01
Maltotetraose	0.237 ± 0.007 ^{cde}	8.7 ± 0.6 ^{cdef}	8.49 ± 0.03
Mannose	0.266 ± 0.003 ^{defg}	8.2 ± 0.2 ^{bcde}	8.64 ± 0.03
Raffinose	0.350 ± 0.007 ^{fghi}	8.1 ± 0.5 ^{bcd}	8.71 ± 0.03
Rhamnose	0.211 ± 0.014 ^{bcd}	10.6 ± 0.2 ^{efgh}	8.73 ± 0.06
Sorbitol	0.410 ± 0.006 ^{hij}	9.7 ± 1.3 ^{cdefg}	8.66 ± 0.04
Sucrose	0.253 ± 0.014 ^{cdef}	10.2 ± 1.8 ^{defgh}	8.88 ± 0.03
Starch	0.078 ± 0.007 ^a	5.7 ± 0.1 ^{ab}	8.76 ± 0.03
Xylose	0.417 ± 0.008 ^{ij}	9.8 ± 0.5 ^{cdefg}	8.78 ± 0.04
Metal ions (%, w/v)			
Control**	0.096 ± 0.009 ^a	6.1 ± 0.4 ^a	8.70 ± 0.10
Fe ³⁺ (0.05%)	0.109 ± 0.005 ^a	6.9 ± 0.5 ^a	8.62 ± 0.07
Ca ²⁺ (0.05%)	0.102 ± 0.001 ^a	5.8 ± 0.4 ^a	8.75 ± 0.06
Mg ²⁺ (0.025%) + Fe ³⁺ (0.025%)	0.097 ± 0.003 ^a	4.8 ± 0.6 ^a	8.60 ± 0.07
Fe ³⁺ (0.025%) + Ca ²⁺ (0.025%)	0.104 ± 0.001 ^a	6.1 ± 1.8 ^a	8.82 ± 0.04
Mg ²⁺ (0.025%) + Ca ²⁺ (0.025%)	0.092 ± 0.004 ^a	5.8 ± 0.3 ^a	8.76 ± 0.03

The shake flask experiments were performed for 96 h at 28 °C, 130 rpm. Data are means ± standard

deviations of three determinations. Column data followed by the same superscript letter were not significantly different ($p \leq 0.05$; by Tukey test).

* Control refers to the basal medium in the absence of any sugar additive.

** Control with 0.05% (w/v) of Mg²⁺ in the basal medium.

Table 5. Effect of nitrogen sources on lipase production:

Nitrogen source	Lipase activity (IU/ml)	Biomass (mg/ml)	Final pH
Organic nitrogen source (0.5%, w/v)			
Control*	0.071 ± 0.002 ^d	6.6 ± 1.9 ^{cd}	8.84 ± 0.03
Phytone peptone	0.131 ± 0.021 ^e	8.8 ± 0.4 ^d	8.72 ± 0.01
Yeast extract	0.104 ± 0.005 ^e	9.1 ± 0.1 ^d	8.69 ± 0.03
Corn steep liquor	0.061 ± 0.006 ^{cd}	7.6 ± 0.6 ^d	8.77 ± 0.03
Beef extract	0.046 ± 0.004 ^{bcd}	8.4 ± 0.7 ^d	8.90 ± 0.17
Skim milk	0.042 ± 0.001 ^{abc}	4.7 ± 0.1 ^{bc}	8.75 ± 0.17
Wheat peptone	0.030 ± 0.002 ^{ab}	6.6 ± 0.4 ^{cd}	8.94 ± 0.03
Fish peptone	0.026 ± 0.006 ^{ab}	1.0 ± 0.1 ^a	8.63 ± 0.03
Tryptone	0.024 ± 0.002 ^{ab}	8.1 ± 0.9 ^d	8.64 ± 0.03
Casein hydrolysate	0.023 ± 0.002 ^{ab}	3.6 ± 0.3 ^{ab}	8.59 ± 0.01
Casein	0.022 ± 0.001 ^{ab}	1.0 ± 0.1 ^a	8.82 ± 0.03
Wheat gluten	0.017 ± 0.001 ^a	2.0 ± 0.4 ^{ab}	8.78 ± 0.04
Different % combination of PP and YE			
0.5% PP + 0.1% YE	0.096 ± 0.005 ^a	6.7 ± 0.4 ^a	8.67 ± 0.03
0.4% PP + 0.2% YE	0.102 ± 0.001 ^a	7.4 ± 0.3 ^a	8.77 ± 0.04
0.3% PP + 0.3% YE	0.100 ± 0.001 ^a	6.6 ± 0.6 ^a	8.87 ± 0.07
0.2% PP + 0.4% YE	0.104 ± 0.002 ^a	8.9 ± 0.5 ^a	8.74 ± 0.01
0.1% PP + 0.5% YE	0.096 ± 0.002 ^a	6.8 ± 0.8 ^a	8.78 ± 0.10
0.6% PP	0.092 ± 0.007 ^a	6.9 ± 0.7 ^a	8.73 ± 0.01
0.6% YE	0.097 ± 0.001 ^a	6.8 ± 0.6 ^a	8.81 ± 0.03
Inorganic nitrogen source (0.05%, w/v)			
Control***	0.098 ± 0.001 ^{de}	5.1 ± 0.1 ^b	8.96 ± 0.03
NH ₄ Cl	0.071 ± 0.001 ^{ab}	2.5 ± 0.7 ^{ab}	8.82 ± 0.04
(NH ₄) ₂ SO ₄	0.078 ± 0.002 ^{abc}	3.0 ± 1.4 ^{ab}	9.04 ± 0.06
(NH ₄) ₂ HPO ₄	0.104 ± 0.003 ^c	5.7 ± 0.5 ^b	8.78 ± 0.03
NH ₄ H ₂ PO ₄	0.090 ± 0.002 ^{cde}	4.0 ± 1.4 ^{ab}	9.00 ± 0.03
CH ₃ COONH ₄	0.086 ± 0.001 ^{bcd}	3.6 ± 0.8 ^{ab}	9.14 ± 0.06
NH ₄ NO ₃	0.090 ± 0.005 ^{cde}	4.9 ± 0.5 ^{ab}	9.02 ± 0.04
Urea	0.062 ± 0.011 ^a	1.5 ± 0.7 ^a	9.13 ± 0.18

The shake flask experiments were performed for 96 h at 28 °C, 130 rpm. Data are means ± standard deviations of three

determinations. Column data followed by the same superscript letter were not significantly different ($p \leq 0.05$; by Tukey test).

* Control refers to the basal medium with 0.5% (w/v) Bacto-peptone as organic nitrogen source.

** Refers to control containing 0.5% (w/v) of phytone peptone (PP) and 0.1% (w/v) of yeast extract (YE) in the basal medium.

*** Refers to control containing 0.05 % (w/v) of NaNO₃ as inorganic nitrogen source in basal medium.

Table 6. Stability of *A. mediterranei* DSM 43304 lipase in organic solvents:

Organic solvent	log <i>P</i>	Relative activity (%)	
		After 1 h	After 24 h
Control		100.0 ± 1.2	100.0 ± 0.6
DMSO	− 1.3	109.4 ± 2.3	89.6 ± 3.7 ^a
DMF	− 1.04	131.3 ± 6.3 ^a	137.6 ± 0.9 ^a
Methanol	− 0.76	129.5 ± 2.8 ^a	136.5 ± 0.6 ^a
2-propanol	− 0.28	129.5 ± 4.3 ^a	112.6 ± 2.5 ^a
Ethanol	− 0.24	109.6 ± 2.0	115.9 ± 3.4 ^a
Acetone	− 0.23	131.7 ± 5.2 ^a	134.6 ± 2.8 ^a
<i>t</i> -Butanol	0.35	96.3 ± 5.8	75.4 ± 3.4 ^a
Pyridine	0.64	72.4 ± 3.5 ^a	49.8 ± 2.8 ^a
Diethyl ether	0.85	116.8 ± 2.6 ^a	136.9 ± 1.8 ^a
Benzene	2.0	169.7 ± 2.0 ^a	169.1 ± 1.2 ^a
Chloroform	2.0	113.8 ± 6.3	134.1 ± 2.2 ^a
Toluene	2.5	153.3 ± 3.6 ^a	157.2 ± 2.8 ^a
Carbon tetrachloride	2.64	137.2 ± 3.7 ^a	117.6 ± 0.9 ^a
Cyclohexene	2.86	138.6 ± 2.3 ^a	109.3 ± 2.2
<i>p</i> -Xylene	3.1	161.9 ± 2.0 ^a	133.7 ± 2.2 ^a
<i>n</i> -Hexane	3.5	156.7 ± 3.2 ^a	111.7 ± 4.3 ^a
Dodecane	6.6	115.5 ± 3.3 ^a	68.0 ± 3.9 ^a
Petroleum ether	≈ 3.0	140.5 ± 2.6 ^a	123.7 ± 2.2 ^a

Lipase preparation was incubated in each organic solvent (25%) at 30 °C for 1 and 24 h.

Values represent the mean of three replicates.

^a Significantly different ($p \leq 0.05$; by Tukey test) with respect to the control.

Table 7. Effect of metal ions/effectors on stability of *A. mediterranei* DSM 4334 lipase:

Metal ion/effector (1 mM)	Relative activity (%)
Control	100.0 ± 1.0
Ag ⁺	88.2 ± 5.5
Ba ²⁺	99.1 ± 1.0
Ca ²⁺	91.1 ± 4.5
Co ²⁺	91.1 ± 0.6
Cu ²⁺	100.2 ± 5.2
Fe ³⁺	115.2 ± 7.1 ^a
Hg ²⁺	20.0 ± 3.2 ^a
Li ⁺	79.2 ± 2.6 ^a
Mg ²⁺	81.3 ± 3.5 ^a
Mn ²⁺	112.9 ± 5.2
Ni ²⁺	95.2 ± 1.9
Pb ²⁺	97.0 ± 0.6
Zn ²⁺	95.9 ± 1.6
NH ₄ ⁺	103.8 ± 1.8
EDTA	103.6 ± 1.6
SDS	100.0 ± 2.9
Urea	88.1 ± 2.9
PMSF	85.3 ± 3.2 ^a

The lipase preparation was incubated in the presence of various compounds at 60 °C for 10 min.

Values represent the mean of three replicates.

^a Significantly different ($p \leq 0.05$; by Tukey test) with respect to the control.

Figure captions:

Figure 1.

Screening for lipolytic actinomycete strains on rhodamine B agar.

a. *Amycolatopsis coloradensis* DSM 44225; **b.** *A. amakusaensis* NRRL B-3351; **c.** *A. fastidiosa* DSM 43855; **d.** *A. mediterranei* DSM 43304; **e.** *A. rubida* DSM 44637; **f.** *A. sulphurea* DSM 46092; **g.** *Nocardia araoensis* DSM 44729; **h.** *N. higoensis* DSM 44732; **i.** *N. kruckzakiae* DSM 44877; **j.** *Streptomyces amquistii* NRRL b-1685, **k.** *S. griseus* subsp. *griseus* DSM 40236; **l.** *S. coelicolor* A3(2); **m.** *S. annulatus* NRRL B-2000; **n.** *S. arabicus* NRRL B-1733; **o.** *S. aurantiogriseus* NRRL B-5416; **p.** *S. rochei* DSM 40231; **q.** *S. aureoverticillatus* NRRL B-3326; **r.** *S. althioticus* NRRL B-3981

Figure 2.

Time course of lipase production by *A. mediterranei* DSM 43304. Basal medium was seeded with 10% (v/v) inoculum and incubated at 28 °C for 192 h at 130 rpm. Samples were withdrawn at 24 h intervals to monitor biomass (■), pH (○) and lipase activity (□) in cell-free supernatants.

Figure 3.

Effect of incubation temperature on lipase production (□) and biomass (■) during growth of *A. mediterranei* DSM 43304 in shake flask cultures after 96 h of incubation at temperatures varying from 20–45 °C.

Figure 4.

Effect of initial pH on lipase production (□) and biomass (■) in shake flask cultures after 96 h of cultivation with initial medium pH varying from 5.0 to 9.0.

Figure 5.

Effect of temperature on *A. mediterranei* DSM 43304 lipase activity (■) and stability (□).
Assay conditions: 30–90 °C, 50 mM Tris–HCl buffer pH 8.0. Thermal stability of the
enzyme was studied by incubating the enzyme at various temperatures (30, 40, 50, 60, 70,
80 and 90 °C) for 3 h. Residual activity (%) at each temperature was calculated relative to
that at 0 h as 100%.

Figure 6.

Effect of pH on *A. mediterranei* DSM 43304 lipase activity (■) and stability (□).
For stability studies, residual activities were measured after 24 h incubation at 4 °C in the
presence of different buffers: HCl–KCl (pH 2.0), sodium citrate (pH 3.0), succinate–NaOH
(pH 4.0 and 5.0), sodium phosphate (pH 6.0 and 7.0), Tris–HCl (pH 8.0 and 9.0) and
glycine–NaOH (pH 10.0), all buffers at 50 mM concentration. Assay conditions: 60 °C,
Tris–HCl buffer, pH 8.0, 50 mM. The activities were compared to the activity determined
in 50 mM Tris–HCl buffer, pH 8.0 without pre-incubation.

Figure 1.

1022 **Figure 2.**

1023
1024
1025
1026
1027
1028
1029
1030
1031
1032
1033
1034
1035

1036 **Figure 3.**

1037
1038
1039
1040
1041
1042
1043
1044
1045
1046
1047
1048
1049

1050 **Figure 4.**

1051
1052
1053
1054
1055
1056
1057
1058
1059
1060
1061
1062
1063

Figure 5.

1078 **Figure 6.**

1079

1080

1081

1082