



2013

The Relationship between Socioeconomic Status and Nutritional knowledge in Women during Pregnancy

Michelle Lane

Technological University Dublin, Michelle.Lane@student.dit.ie

Ellen Barrett

Dublin Institute of Technology, Ellen.Barrett@student.dit.ie

Amy O'Higgins

University College Dublin, amyohiggins@rcsi.ie

Laura Mullaney

Dublin Institute of Technology, lauraemullaney@gmail.com

Michael Turner

University College Dublin, Michael.Turner@ucd.ie

See next page for additional authors

Follow this and additional works at: <https://arrow.dit.ie/scschbioart>



Part of the [Dietetics and Clinical Nutrition Commons](#)

Recommended Citation

Lane, M., Barrett, E. M., O'Higgins, A. C., Mullaney, L. E., Turner, M. J., McCartney, D. M. (2013). The relationship between socioeconomic status and nutritional knowledge in women during pregnancy, *Proceedings of the Nutrition Society*, 72 (OCE3), E162. doi:10.1017/S0029665113001857

This Article is brought to you for free and open access by the School of Biological Sciences at ARROW@TU Dublin. It has been accepted for inclusion in Articles by an authorized administrator of ARROW@TU Dublin. For more information, please contact yvonne.desmond@dit.ie, arrow.admin@dit.ie, brian.widdis@dit.ie.



Authors

Michelle Lane, Ellen Barrett, Amy O'Higgins, Laura Mullaney, Michael Turner, and Daniel McCartney

The relationship between socioeconomic status and nutritional knowledge in women during pregnancy

M. Lane¹, E. M. Barrett¹, A. O’Higgins², L. Mullaney¹, M. J. Turner² and D. McCartney¹

¹Dublin Institute of Technology, Dublin 8, Republic of Ireland and ²The Coombe Women’s and Infants University Hospital, Cork Street, Dublin 8, Republic of Ireland

Epidemiological data have shown that socioeconomic status affects nutritional knowledge and dietary quality⁽¹⁾. Irish national food surveys have also suggested that socioeconomic differences in food group, nutrient intakes, dietary attitudes and health behaviours exist^(2,3). It is known that pregnancy can enhance a women’s nutritional awareness⁽⁴⁾. The aim of the present study was to assess whether nutritional knowledge in pregnant women in Ireland is affected by socioeconomic status.

One hundred and sixteen pregnant women aged 16–41 years (mean age of 29.5 years) were recruited at their initial antenatal booking visit (10–15 weeks gestation) at a large Dublin maternity hospital. Formal education and material deprivation were used to assess the socioeconomic status of respondents. The women were classified into three educational categories: no formal education to lower secondary education, upper secondary education and third level education. Deprivation status was assessed by determining the number of basic necessities⁽⁵⁾ respondents had had to forego in the previous year due to lack of money, with respondents categorised into three groupings: low(none), medium(1–2) and high (3+).

The women’s level of nutritional knowledge was assessed using a self-administered questionnaire of 12 multiple choice questions. The questions were formulated from the Best Practice for Infant Feeding in Ireland guidelines published by the Food Safety Authority of Ireland in 2012⁽⁶⁾. The questionnaire assessed respondents’ knowledge regarding essential food groups and nutrients in pregnancy as well as lifestyle factors such as smoking and exercise. Each respondent received a score out of 12 on their nutritional knowledge questionnaire (max. = 11, min. = 1), with participants then categorised as having high (score ≥7) or low (score ≤6) nutritional knowledge based on these scores.

	No/Primary/Lower secondary education	Upper secondary education	Third level education	Significance
Low nutritional knowledge	8	21	15	<i>p</i> = 0.01
High nutritional knowledge	5	16	51	

Pearson’s chi-square test.

The results of a Pearson’s chi-square test confirmed that formal education was inversely related to nutritional knowledge (*p*<0.05). When adjusted for the confounding effect of material disadvantage (as assessed by deprivation status), low formal education remained predictive of a lower nutritional knowledge score (*p* = 0.02), while deprivation status did not predict poorer knowledge scores (*p* = 0.278). This indicates that socio-ecological rather than material disadvantage is the primary predictor of poor nutritional knowledge in this group.

1. Darmon N & Drewnowski A (2008) *Am J Clin Nutr* **87**, 1107–1117.
2. Hearty AP, McCarthy SN, Kearney JM *et al.* (2007) Relationship between attitudes towards healthy eating and dietary behaviour, lifestyle and demographic factors in a representative sample of Irish adults. *Appetite* **48**, 1–11.
3. Kelleher C, Friel S, Nolan G *et al.* (2002) *Proc Nutr Soc* **61**, 527–536.
4. Szwajcer E, Hiddink GJ, Maas L *et al.* (2012) *Fam Pract* **29**, i82–i88.
5. Government of Ireland (2007) *National Action Plan for Social Inclusion 2007–2016*. Dublin: The Stationery Office. www.socialinclusion.ie/documents/NAPinclusionReportPDF.pdf – accessed 10th March 2013.
6. Food Safety of Ireland (2012) *Best Practice for Infant Feeding in Ireland, From pre-conception through the first year of an infant’s life*. Dublin: Lower Abbey Street. www.fsai.ie/bestpracticeforinfantfeedinginireland.html – accessed 12th March 2013.