



2011-03-01

Maternal Health Behaviours During Pregnancy in an Irish Obstetric Population and Their Associations with Socio-Demographic and Infant Characteristics

Roslyn Tarrant

Dublin Institute of Technology, roslyn.tarrant@gmail.com

Katherine Younger

Dublin Institute of Technology, katherine.younger@dit.ie

Margaret Sheridan-Pereira

Trinity College Dublin, drmargaretsheridan@gmail.com

John Kearney

Dublin Institute of Technology, john.kearney@dit.ie

Follow this and additional works at: <http://arrow.dit.ie/scschbioart>



Part of the [Clinical Epidemiology Commons](#), [Community Health and Preventive Medicine Commons](#), and the [Dietetics and Clinical Nutrition Commons](#)

Recommended Citation

Tarrant RC, Younger KM, Sheridan-Pereira M & Kearney JM (2011) Maternal health behaviours during pregnancy in an Irish obstetric population and associations with socio-demographic and infant characteristics. *European Journal of Clinical Nutrition* March 2 (on-line ahead of publication). doi:10.1038/ejcn.2011.16.

This Article is brought to you for free and open access by the School of Biological Sciences at ARROW@DIT. It has been accepted for inclusion in Articles by an authorized administrator of ARROW@DIT. For more information, please contact yvonne.desmond@dit.ie, arrow.admin@dit.ie, brian.widdis@dit.ie.



ORIGINAL ARTICLE

Maternal health behaviours during pregnancy in an Irish obstetric population and their associations with socio-demographic and infant characteristics

R Tarrant¹, K Younger², M Sheridan-Pereira³ and JM Kearney²

¹The Department of Clinical Nutrition and Dietetics, Our Lady's Children's Hospital, Dublin, Republic of Ireland; ²School of Biological Sciences, Dublin Institute of Technology, Dublin, Republic of Ireland and ³The Department of Paediatrics, The Coombe Women and Infants University Hospital, Dublin, Republic of Ireland

Q1

Background/Objectives: To examine the prevalence and combined occurrence of peri-conceptual folic acid (FA) supplement use, smoking and alcohol consumption during pregnancy in a sample of women in Dublin, and determine the factors associated with these health behaviours.

Subjects/Methods: A prospective observational study (2004–2006) involving the recruitment of 491 pregnant women from antenatal clinics in a Dublin maternity hospital, with postpartum follow-up of 450 eligible mothers. Data on FA use, maternal smoking and alcohol consumption patterns during pregnancy were collected from the antenatal patient-administered questionnaire, which was completed by participants, and returned to the investigator on the day of recruitment.

Results: The median gestational age of women at recruitment was 36 weeks. A combined 24.2% of mothers commenced FA at the recommended time, avoided alcohol consumption and smoking during pregnancy. In all, 35.3% of mothers reported to consuming alcohol, 20.9% smoked during pregnancy and 44.4% commenced FA at the recommended time. Mothers <25 years were more likely to have not taken FA at the recommended time (adjusted odds ratio (aOR): 4.0, 95% confidence interval (CI): 1.64–9.77) and were more likely to have smoked during pregnancy (aOR: 3.56, 95% CI: 1.32–9.57). Irish nationality positively predicted both alcohol consumption (aOR: 4.37, 95% CI: 1.88–10.15) and smoking (aOR: 10.92, 95% CI: 1.35–87.98) during pregnancy.

Conclusions: Educational efforts are still necessary to convince women of Irish nationality, in particular, of the adverse effects of smoking and alcohol consumption on fetal outcome. Women <25 years should be specifically targeted in smoking cessation and FA promotional campaigns.

European Journal of Clinical Nutrition (2011) **0**, 000–000. doi:10.1038/ejcn.2011.16

Keywords: maternal health; pregnancy; cigarette smoking; alcohol consumption; folic acid; prenatal care

Introduction

Optimal pregnancy and infant health outcomes are associated with non-smoking (Kramer, 2003) and absolute maternal avoidance of alcohol consumption (World Health Organisation, 2006) during pregnancy, in addition to compliance with peri-conceptual folic acid (FA) supplementation (Kirke *et al.*, 1992). Maternal cigarette smoking during pregnancy is associated with increased risks for

placenta previa (Monica and Lilja, 1995) and abruption (Pollack *et al.*, 2000), intrauterine growth retardation (Horta *et al.*, 1997), preterm delivery (Kyrklund-Blomberg and Cnattingius, 1998) and sudden infant death syndrome (Wisborg *et al.*, 2000). Prenatal alcohol exposure is a known preventable cause of birth defects (Krulwich, 2005), and Fetal Alcohol syndrome, the leading cause of mental retardation worldwide and the foremost preventable cause of neurobehavioral and developmental abnormalities (Banakar *et al.*, 2009). To avoid these adverse risks, policy in Ireland recommends non-smoking (Health Service Executive, 2008) and absolute abstinence from alcohol both during pregnancy and among women wishing to conceive (Department of Health and Children, 2008).

Q2

Correspondence: Dr JM Kearney, School of Biological Sciences, Dublin Institute of Technology, Kevin Street, Dublin 8, Republic of Ireland.

E-mail: john.kearney@dit.ie

Received 7 July 2010; revised 13 January 2011; accepted 13 January 2011

Furthermore, it is universally accepted that maternal compliance with peri-conceptual FA supplementation can substantially decrease the risk of congenital malformations (Wilcox *et al.*, 2007), as well as the incidence (Czeizel and Dudas, 1992) and recurrence (MRC, 1991) of neural tube defects (NTDs), including spina bifida, anencephaly and encephalocele. However, the potential for preventing NTDs in Europe by increasing folate status is far from being achieved (Busby *et al.*, 2005). Department of Health, Republic of Ireland (1992) recommends that women who could possibly become pregnant should consume a daily 400 µg FA supplement 3 months before conception and during the first 12 weeks of pregnancy, in addition to folate-rich foods. Compliance with this recommendation is particularly important for women in Ireland, as Irish NTD rates are among the highest in Europe (Food Safety Authority of Ireland, 2006).

In spite of these concerns, only limited research has quantified the prevalence and predictive factors of these health behaviours among women in Ireland. Such research is critical to enable policymakers and clinicians design targeted interventions and effective preventative strategies, especially in high-risk populations. The present study, thus, aimed to examine the prevalence and combined occurrence of peri-conceptual FA supplement use, smoking and alcohol consumption during pregnancy in a sample of women in Dublin, and to determine the factors associated with these health behaviours.

Methods

Study population and design

Details of the study methods and representativeness are reported in Tarrant *et al.* (2010a). Briefly, from June 2004 to October 2006 this prospective observational study involved the recruitment of 539 pregnant women ≥ 24 weeks gestational age from separate public and private/semi-private antenatal clinics in the CWIUH (Coombe Women and Infants University Hospital, a maternity hospital which reported the highest number of mothers who delivered infants weighing > 500 g from the three Dublin maternity hospitals in 2006 ($n = 8084$). From the 539 women recruited who were eligible to partake, 491 women (91%) agreed to participate and gave signed consent. Following application of exclusion criteria, 450 mother–infant pairs were subsequently followed up at 6 weeks and 6 months postpartum. This final sample ($n = 450$) represents 5.7, 5.8 and 5.6% of the total number of mothers who delivered infants weighing > 500 g in the CWIUH in west Dublin during the years 2004, 2005 and 2006, respectively (Coombe Women and Infants University Hospital, 2006). Ethical approval was granted by the Ethical Committees of the CWIUH and Dublin Institute of Technology.

Data collection

Following consent to study participation, mothers completed the first patient-administered questionnaire while waiting in antenatal clinics, and on the same day returned the questionnaire to the investigator. Data were collected on maternal education level, occupation, age, marital status, parity and whether the pregnancy was planned or unplanned (Tarrant, 2008). Mothers reported on their height and pre-pregnancy weight measurement, from which body mass index was computed (weight (kg)/height (m^2)), and classified according to World Health Organisation (2003) definitions. All patient-administered questionnaires were checked by the investigator with mothers present, to ensure completeness of the data and to clarify any ambiguities.

Women were asked to state whether they (1) discontinued cigarette smoking upon recognition of pregnancy, were (2) non-smokers pre- and during pregnancy or (3) current smokers during pregnancy, in which case data were collected on the number of cigarettes smoked/day both pre- and during pregnancy. 'Non-smokers during pregnancy' refers to the women who were non-smokers pre- and during pregnancy, as well as those who discontinued smoking upon recognition of pregnancy. Data on mothers' alcohol consumption status during pregnancy were also recorded, including the number of alcohol units consumed/week both pre- and during pregnancy and the specific type of alcohol drink (wine, beer/stout, alcopop drinks and measure of spirits) consumed/unit.

Women were questioned on peri-conceptual FA supplement use, and data were elicited on the brand/type of FA supplement taken per day, as well as the timing of commencement. As neural tube closure occurs between 22 and 28 days postconception (Ray *et al.*, 2004), maternal uptake of FA 'at the recommended time peri-conceptionally' in the present study was defined as commencing FA supplementation at recommended doses ≥ 400 µg/day or at lower doses ≥ 200 µg/day, either preconceptionally or during the first month postconception. Women who reported taking a FA supplement and recorded the timing of commencement, but did not report the daily dose or brand consumed, were included in all analyses. Lack of compliance with peri-conceptual FA supplementation at the recommended time includes the women who took FA supplements > 1 month postconception, as well as those who did not take any supplemental FA.

While the study participants were completing the antenatal patient-administered questionnaire in clinic, information on maternal/paternal country of birth and race, as well as a record of whether FA was taken pre or postconceptionally was collected by the investigator from the mother's medical notes. Any ambiguities between maternal recordings of the timing of FA uptake vs the data collected from the medical notes were clarified at the time. The infants' medical notes provided data on gender, type of delivery, gestational age and health status of infants at birth, as well as birth

weight (kg), length (cm) and head circumference (cm) measurements. Infant feeding factors related to breastfeeding and weaning onto solid foods, as previously reported (Tarrant *et al.*, 2010a, b), were collected from the 6-week and 6-month interviewer-administered questionnaires.

Statistical analyses

The analyses were performed using the Statistical Package for the Social Sciences (version 17; SPSS Inc., Chicago, IL, USA). Categorical data were compared using the χ^2 statistics test or Fisher's exact test, and significance levels were set at $P < 0.05$. For continuous normally distributed data, independent *t*-tests and one-way analysis of variance were performed. Wilcoxon signed-rank tests were used for non-normally distributed data. Continuous variables are summarised using mean values \pm s.d. or medians with interquartile ranges, as appropriate.

Three separate logistic regression analyses were performed to determine the factors associated with the dependent variables: (1) alcohol consumption, (2) maternal cigarette smoking during pregnancy, as well as (3) non-compliance with peri-conceptual FA supplementation at the recommended time/no FA. First, within each category a univariate analysis was performed to identify the significant ($P < 0.05$) independent variables for inclusion in the subsequent multiple forward stepwise logistic regression analyses. The significant variables identified in the univariate analysis were then entered singly into the regression model, and were retained if statistical significance at $P < 0.05$ criterion was achieved. Final models are presented using adjusted odds ratios (aORs) with their 95% confidence intervals (CIs).

Results

Sample characteristics

Complete data were available for 450 mother-term infant pairs. The median age of participating women at antenatal recruitment, and hence, the timing of completion and return of patient-administered questionnaires to the investigator, was 36 weeks (interquartile range 32–39 weeks). Table 1 summarises the characteristics of the total sample, including those of the Irish national ($n = 401$) and non-Irish national ($n = 49$) mothers.

Irish national, compared with non-Irish national mothers, were significantly more likely to smoke (23.2 vs 2%, $P = 0.000$) and consume alcohol (37.9 vs 14.3%, $P = 0.002$) during pregnancy, however, equal proportions of mothers in both populations complied with peri-conceptual FA supplementation at the recommended time (44.6 vs 42.9%, $P = 0.933$). Although differences in maternal social class were found between the two populations ($P = 0.003$), no significant differences were observed in maternal age, education and pre-pregnancy body mass index, marital status, planned/unplanned pregnancy, infant birth weights, gender or parity.

Compliance with public health recommendations

As shown in Table 2, a combined 109 women (24.2%) complied with all three public health recommendations to avoid smoking and alcohol consumption during pregnancy, and commence supplemental FA at the recommended time. Complete non-compliance with all of these recommendations was found in 28 women (6.2%) in our sample.

A strong social class gradient was found between smoking during pregnancy as well as lack of compliance with peri-conceptual FA supplementation at the recommended time/no FA (Table 3), with a significantly higher prevalence of these health behaviours observed among mothers from lower social classes. No association was found between alcohol consumption during pregnancy and maternal social class with a similar proportion of mothers in social class I found to consume alcohol during pregnancy (40.7%) as those in social class III (40.8%).

Smoking during pregnancy

During pregnancy, 94 (20.9%) women smoked, 295 (65.5%) women were non-smokers and 61 (13.6%) women discontinued smoking upon recognition of pregnancy. In 77/94 smokers (82%) during pregnancy, the number of cigarettes smoked/day decreased significantly from a mean of 19.8 (s.d. 8.6) cigarettes smoked/day pre-pregnancy to 11.7 (s.d. 7.5) cigarettes smoked/day during pregnancy ($P = 0.000$). No change in maternal cigarette smoking behaviour was observed in 10 of the smokers (10.6%) during pregnancy, while 7 women (7.4%) increased the number of cigarettes smoked/day from pre- to during pregnancy.

Logistic regression analysis showed that the risk of maternal cigarette smoking during pregnancy was associated with single, multiparous mothers, aged < 25 years who delivered infants with a birth weight < 3 kg, and those who did not initiate breastfeeding postpartum (Table 4). Irish nationality was positively associated with smoking during pregnancy (OR: 10.92, 95% CI: 1.35–87.98).

In addition, mothers who smoked > 10 cigarettes/day ($n = 32$) delivered infants with significantly lower mean infant birth weights (3.11 kg (s.d. 0.52) vs 3.66 kg (s.d. 0.5), $P = 0.000$) and birth head circumference measurements (34.25 cm (s.d. 1.6) vs 35.33 cm (s.d. 1.37), $P = 0.001$) than mothers who never smoked at all pre/during pregnancy ($n = 295$).

Alcohol consumption during pregnancy

In 144/159 (90.5%) of the alcohol consumers during pregnancy, a decreased weekly alcohol consumption from pre-pregnancy (mean 7.13 (s.d. 5.58) alcohol units/week) to during pregnancy (mean 2.45 (s.d. 2.08) alcohol units/week) ($P = 0.000$) was observed. An increase in weekly alcohol consumption from pre- to during pregnancy was found in two women (1.2%) in this sample, while no change in alcohol consumption intake was found in 13 women (8%).

Table 1 Selected maternal socio-demographic, infant and postnatal infant feeding characteristics of the total sample, and by nationality

Characteristic	Total sample (n = 450) n (%)	Irish national mothers (n = 401) n (%)	Non-Irish national mothers (n = 49) n (%)	P-value
<i>Maternal social class (based on occupation)^a</i>				
Social class I	140 (31.1)	124 (30.9)	16 (32.7)	0.003
Social class II	128 (28.4)	123 (30.7)	5 (10.2)	
Social class III	49 (10.9)	46 (11.5)	3 (6.1)	
Unknown category (unemployed/students)	57 (12.7)	46 (11.5)	11 (22.4)	
Stay-at-home mothers' /home-makers'	76 (16.7)	62 (15.5)	14 (28.6)	
<i>Maternal age (years)</i>				
> 34	85 (18.9)	73 (18.2)	12 (24.5)	0.436
25–34	267 (59.3)	238 (59.4)	29 (59.2)	
<25	98 (21.8)	90 (22.4)	8 (16.3)	
<i>Highest maternal education qualification</i>				
Primary/secondary	177 (39.3)	160 (39.9)	17 (34.7)	0.076
Vocational/training course	124 (27.6)	115 (28.7)	9 (18.4)	
Third level degree/postgraduate level	149 (33.1)	126 (31.4)	23 (46.9)	
<i>Pre-pregnancy BMI (kg/m²)^b</i>				
<20	47 (10.4)	41 (10.2)	6 (12.2)	0.860
20–24.9	253 (56.2)	225 (56.1)	28 (57.1)	
≥25	150 (33.3)	135 (33.7)	15 (30.6)	
<i>Marital status</i>				
Married/cohabitating	407 (90.4)	359 (89.5)	48 (98)	0.069
Single	43 (9.6)	42 (10.5)	1 (2)	
<i>Parity</i>				
Primiparous	213 (47.3)	196 (48.9)	17 (34.7)	0.084
Multiparous	237 (52.7)	205 (51.1)	32 (65.3)	
<i>Planned current pregnancy</i>				
Yes	282 (62.7)	250 (62.3)	32 (65.3)	0.804
No	168 (37.3)	151 (37.7)	17 (34.7)	
<i>Smoking during pregnancy</i>				
Yes	94 (20.9)	93 (23.2)	1 (2)	0.000
No	356 (79.1)	308 (76.8)	48 (98)	
<i>Alcohol consumption during pregnancy</i>				
Yes	159 (35.3)	152 (37.9)	7 (14.3)	0.002
No	291 (64.7)	249 (62.1)	42 (85.7)	
<i>Folic acid supplement use at the recommended time</i>				
Yes ^c	200 (44.4)	179 (44.6)	21 (42.9)	0.933
No ^d	250 (55.5)	222 (55.4)	28 (57.1)	
<i>Birth weight (kg)</i>				
<3	65 (14.4)	58 (14.5)	7 (14.3)	0.781
3–4	309 (68.7)	277 (69.1)	32 (65.3)	
>4	76 (16.9)	66 (16.5)	10 (20.4)	
<i>Gender of infant</i>				
Male	250 (55.6)	223 (55.6)	27 (55.1)	1.000
Female	200 (44.4)	178 (44.4)	22 (44.9)	
<i>Initiated breastfeeding postpartum^e</i>				
Yes	228 (50.7)	189 (47.1)	39 (79.6)	0.000
No	222 (49.3)	212 (52.9)	10 (20.4)	

Abbreviation: BMI, body mass index.

Results are expressed as number of subjects and percentage.

Significant differences (*P*-values) between Irish national and non-Irish national mothers using cross tabulations and χ^2 statistics test or Fisher's exact test in cases where one or more of the cells has a frequency of 5 or less; the Yates's continuity correction value was used for all 2 × 2 tables.

^aMaternal occupations were categorised according to the social class categories from the UK (Office of Population Census and Surveys, 1991) and Irish (Central Statistics Office Census, 1996) classification system; a separate category was created for those who reported that they were 'stay-at-home mothers'.

^bBMI categories are in accordance with World Health Organisation (2003) definitions (<20 kg/m², underweight; 20–24.9 kg/m², normal weight; ≥25 kg/m², overweight). BMI values were derived from mothers' self-reported pre-pregnancy weight and height.

^cIncludes women who commenced folic acid supplementation at the recommended time, i.e., either preconception (*n* = 181) or during the first month postconception (*n* = 19), at doses 350–400 µg/day.

^dIncludes women who did not commence folic acid supplementation at the recommended time, i.e., >1 month post conception (*n* = 195); these women were categorised with the women who did not take any folic acid supplementation at all (*n* = 55).

^eBreastfeeding initiation was defined as all mothers who 'ever tried to breastfeed' postpartum.

Wine and beer/stout were the most frequently consumed alcoholic drinks by 94 and 59 mothers, respectively, followed by alcopop drinks ($n=15$) and spirits ($n=4$).

In both the univariate and logistic regression analyses, breastfeeding initiation and nationality were the only two variables that were significantly associated with alcohol consumption during pregnancy; notably, Irish nationality

was identified as an independent predictor of alcohol consumption during pregnancy (OR: 4.37, CI: 1.88–10.15) (Table 5).

Peri-conceptual FA supplementation

In all, 395 (87.7%) women reported taking a daily FA supplement during pregnancy, including *Clonfoliac* 400 µg FA/day ($n=315$), *Galfer FA* 350 µg FA/day ($n=21$) and *Pregnacare* 400 µg FA/day ($n=12$), while 47 women reported to not knowing the brand name or dose of the FA supplement consumed. Commencement of peri-conceptual FA supplementation at the 'recommended time' was reported in 200 (44.4%) women, including 181 preconceptionally ($n=39$, 1 month; $n=28$, 2 months; $n=49$, 3 months; $n=35$, 6 months; $n=30$, >6 months preconception) and 19 women during the first month postconception. Among these women who reported taking supplemental FA at the recommended time, the brand and daily dosage of the supplements consumed included: *Clonfoliac* 400 µg FA/day ($n=160$), *Pregnacare* 400 µg FA/day ($n=9$), *Galfer FA* 350 µg FA/day ($n=5$) while the dose and brand of supplemental FA consumed was unknown in 26 of cases. FA supplements were consumed by 160 women during the second month of pregnancy, and 35 women during the third month of pregnancy, while 55 (12.2%) women reported to not consuming any supplemental peri-conceptual FA.

After adjustment, factors positively associated with non-compliance with peri-conceptual FA supplementation at the recommended time/no FA, included maternal cigarette smoking during pregnancy, maternal age <25 years, pre-pregnancy body mass index <20 kg/m² and addition of

Table 2 Compliance with public health recommendations in a sample of pregnant women in Dublin ($n=450$)

Compliance rating	Did not smoke	Did not consume alcohol	Complied with folic acid supplementation at the recommended time ^a	n (%)
0 ^b	No	No	No	28 (6.2)
1 ^c	No	No	Yes	7 (1.5)
1 ^c	No	Yes	No	48 (10.6)
1 ^c	Yes	No	No	51 (11.3)
2 ^d	No	Yes	Yes	11 (2.4)
2 ^d	Yes	No	Yes	73 (16.2)
2 ^d	Yes	Yes	No	123 (27.3)
3 ^e	Yes	Yes	Yes	109 (24.2)

Results are expressed as number of subjects and percentage.

^aIncludes women who complied with folic acid supplementation at the recommended time, i.e., either preconception ($n=181$) or during the first month postconception ($n=19$), at doses 350–400 µg/day.

^bCompliance rating of 0 indicates complete non-compliance with all three public health recommendations.

^cCompliance rating of 1 indicates compliance with one public health recommendation and non-compliance with two recommendations.

^dCompliance rating of 2 indicates compliance with two public health recommendations and non-compliance with one recommendation.

^eA compliance rating of 3 indicates complete compliance with all three public health recommendations.

Table 3 Maternal health behaviours during pregnancy, by maternal social class^a ($n=450$)

Maternal health behaviour	Social class I n (%)	Social class II n (%)	Social class III n (%)	Unknown category (unemployed/students) n (%)	Stay-at-home mothers'/ home-makers' n (%)	P-value
Smoking during pregnancy						
Yes	13 (9.3)	19 (14.8)	12 (24.5)	20 (35.1)	30 (39.5)	0.000
No	127 (90.7)	109 (85.2)	37 (75.5)	37 (64.9)	46 (60.5)	
Alcohol consumption during pregnancy						
Yes	57 (40.7)	43 (33.6)	20 (40.8)	16 (28.1)	23 (30.3)	0.313
No	83 (59.3)	85 (66.4)	29 (59.2)	41 (71.9)	53 (69.7)	
Folic acid supplement use at the recommended time						
Yes ^b	85 (60.7)	69 (53.9)	13 (26.5)	7 (12.3)	26 (34.2)	0.000
No ^c	55 (39.3)	59 (46.1)	36 (73.5)	50 (87.7)	50 (65.8)	

Differences between social class category and maternal health behaviours were examined using cross tabulations and χ^2 statistics test; P-values indicate the significant differences between these groups.

^aMaternal social class was categorised according to mothers' occupation. Social class categories from the UK (Office of Population Census and Surveys, 1991) and Irish (Central Statistics Office Census, 1996) classification system were used with a separate category created for mothers who were 'stay-at-home' mothers.

^bWomen who commenced folic acid supplementation at the recommended time, i.e., either preconception ($n=181$) or during the first month postconception ($n=19$), at doses 350–400 µg/day.

^cWomen who did not commence folic acid supplementation at the recommended time, i.e., >1 month postconception ($n=195$); these women were categorised with the women who did not take any folic acid supplementation at all ($n=55$).

Table 4 Factors associated with smoking during pregnancy in a sample of women in Dublin ($n = 450$)

Characteristic	Non-smokers, n (%)	Smokers, n (%)	P-value ^a	OR ^b (95% CI)	Adjusted OR ^c (95% CI)	P-value ^d
<i>Maternal age (years)</i>						
> 34	76 (89.4)	9 (10.6)	0.000	1.0 ^e	1.0	0.011
25–34	221 (82.8)	46 (17.2)		1.75 (0.82–3.76)	1.46 (0.61–3.48)	
< 25	59 (60.2)	39 (39.8)		5.58 (2.5–14.43)	3.56 (1.32–9.57)	
<i>Highest maternal education qualification</i>						
Third level degree/postgraduate	139 (93.3)	10 (6.7)	0.000	1.0	1.0	0.065
Vocational/training course	98 (79)	26 (21)		3.68 (1.7–7.99)	1.78 (0.74–4.26)	
Primary/secondary	119 (67.2)	58 (32.8)		6.77 (3.31–13.84)	2.58 (1.14–5.84)	
<i>Nationality</i>						
Non-Irish national	48 (98)	1 (2)	0.001	1.0	1.0	0.025
Irish national	308 (76.8)	93 (23.2)		14.49 (1.9–106.4)	10.92 (1.35–87.98)	
<i>Marital status</i>						
Married/cohabitating	338 (83)	69 (17)	0.000	1.0	1.0	0.005
Single	18 (41.9)	25 (58.1)		6.8 (3.5–13.1)	2.93 (1.37–6.27)	
<i>Parity</i>						
Primiparous	178 (83.6)	35 (16.4)	0.037	1.0	1.0	0.008
Multiparous	178 (75.1)	59 (24.9)		1.68 (1.05–2.68)	2.31 (1.25–4.26)	
<i>Birth weight (kg)</i>						
> 4	71 (93.4)	5 (6.6)	0.000	1.0	1.0	0.000
3–4	250 (80.9)	59 (19.1)		3.35 (1.29–8.66)	2.36 (0.85–6.4)	
< 3	35 (53.8)	30 (46.2)		12.17 (4.34–34)	7.83 (2.57–23.77)	
<i>Initiated breastfeeding postpartum^f</i>						
Yes	212 (93)	16 (7)	0.000	1.0	1.0	0.001
No	144 (64.9)	78 (35.1)		7.17 (4.02–12.79)	3.04 (1.59–5.82)	

Abbreviations: CI, confidence interval; OR, odds ratio.

Results are expressed as number of subjects and percentage.

^aStatistical analysis by χ^2 statistics test. The Yates's continuity correction value was used for all 2×2 tables.

^bUnadjusted OR from logistic regression analysis.

^cOR adjusted for all other variables in the table using logistic regression analysis.

^dP-value from adjusted logistic regression model.

^e1.0 denotes the reference category.

^fBreastfeeding initiation was defined as all mothers who 'ever tried to breastfeed' postpartum.

non-recommended condiments (for example, gravy and sauces) to infants' weaning foods (Table 6). In particular, logistic regression analysis indicated that mothers who reported an unplanned pregnancy were significantly more likely to have not taken supplemental FA at the recommended time/at all (OR: 13.24, 95% CI: 7–24.7).

Discussion

Only 24.2% of mothers complied with the public health recommendations to commence peri-conceptual FA supplementation at the recommended time, avoid smoking and completely abstain from alcohol consumption during pregnancy. The fact that 6.2% of mothers did not comply with any of these recommendations highlights the potential health disparities and increased likelihood for a compromised pregnancy outcome within our sample of mothers (Cnattingius, 2004; Busby *et al.*, 2005).

Essentially, over half of the mothers in this study (55.5%) did not achieve optimal protection against NTDs. This finding is of major clinical importance owing to evidence that increased peri-conceptual intakes FA can significantly reduce the risk of first occurrence (Czeizel and Dudas, 1992) and recurrence (MRC, 1991) of NTDs in up to 72% of cases. The fact that Irish NTD rates are among the highest in Europe, affecting 1–1.5 per 1000 total births nationally, and that NTDs account for more national perinatal deaths than any other group of birth defects (Food Safety Authority of Ireland, 2006) further increases the importance of strict compliance with daily peri-conceptual FA supplementation among women who could possibly conceive in Ireland. Although voluntary fortification of foodstuffs with FA has led to a higher folate status in the Irish population, current or future mandatory fortification is not recommended (Food Safety Authority of Ireland, 2008); thus, there is increased maternal reliance on supplemental FA to achieve optimal protection against NTDs. However, the rate of compliance

Table 5 Factors associated with alcohol consumption during pregnancy in a sample of women in Dublin ($n=450$)

Characteristic	Non-alcohol consumers, n (%)	Alcohol consumers, n (%)	P-value ^a	OR (95% CI) ^b	Adjusted OR ^c (95% CI)	P-value ^d
Maternal age (years)						
> 34	48 (56.5)	37 (43.5)		1.0 ^e	1.0	
25–34	175 (65.5)	92 (34.5)		0.68 (0.41–1.12)	0.7 (0.41–1.17)	
<25	68 (69.4)	30 (30.6)	0.170	0.57 (0.31–1.05)	0.59 (0.31–1.15)	0.268
Highest maternal education qualification						
Third level degree/postgraduate	99 (66.4)	50 (33.6)		1.0	1.0	
Vocational/training course	71 (57.3)	53 (42.7)		1.478 (0.9–2.41)	1.63 (0.96–2.75)	
Primary/secondary	121 (68.4)	56 (31.6)	0.120	0.91 (0.57–1.45)	1.17 (0.69–1.99)	0.166
Nationality						
Non-Irish national	42 (85.7)	7 (14.3)		1.0	1.0	
Irish national	249 (62.1)	152 (37.9)	0.002	3.66 (1.6–8.35)	4.37(1.88–10.15)	0.001
Initiated breastfeeding postpartum^f						
No	155 (69.8)	67 (30.2)		1.0	1.0	
Yes	136 (59.6)	92 (40.4)	0.031	1.56 (1.06–2.31)	1.78 (1.15–2.76)	0.010

Abbreviations: CI, confidence interval; OR, odds ratio.

Results are expressed as number of subjects and percentage.

^aStatistical analysis by χ^2 statistics test. The Yates's continuity correction value was used for all 2×2 tables.

^bUnadjusted OR from logistic regression analysis.

^cOR adjusted for all other variables in the table using logistic regression analysis.

^dP-value from adjusted logistic regression model.

^e1.0 denotes the reference category.

^fBreastfeeding initiation was defined as all mothers who 'ever tried to breastfeed' postpartum.

with peri-conceptual FA supplementation at the recommended time in the present study (44.4%) is higher than what has been found in other Dublin-based studies, including 24% (Ward *et al.*, 2004) and 28% (McGuire *et al.*, 2010), suggesting an improved awareness of the importance of peri-conceptual FA supplementation in our sample of mothers. Suboptimal peri-conceptual FA supplementation rates are not just a concern in Ireland, but globally, with rates of 10.3 and 52% reported in Korea (Kim *et al.*, 2009) and the Netherlands (Bekkers and Eskes, 1999), respectively.

In agreement with the literature (Brough *et al.*, 2009; Kim *et al.*, 2009; McGuire *et al.*, 2010), mothers from lower social classes and those who reported an unplanned pregnancy were the least likely to comply with peri-conceptual FA supplementation at the recommended time/or take no FA. To increase peri-conceptual FA uptake, our data indicate that promotional campaigns should target younger women <25 years, smokers, those from lower socio-economic groups and, in particular, women not planning pregnancies. Public health strategies that raise awareness of the peri-conceptual timeframe for optimal NTD protection, among all women of childbearing age, should also be considered.

Previous Irish studies report considerably higher alcohol consumption during pregnancy rates of 78% (Daly *et al.*, 1992) and 89% (McMillan *et al.*, 2006) than those found in the present study (35.3%). Although we found that the vast majority of alcohol consumers decreased their weekly alcohol intake during pregnancy (90.5%), it appears that

more effective public health approaches are required to persuade pregnant women to completely abstain from alcohol consumption. Pre-pregnancy and antenatal advice to women should include the fact that alcohol is a potent teratogen, and consumption during pregnancy can cause a spectrum of malformations and neurological, cognitive and behavioural deficits that are fully preventable (Landgren *et al.*, 2010).

No association between alcohol consumption during pregnancy and social class was observed in our study, with mothers in social class I (40.7%) being as likely to consume alcohol as those in social class III (40.8%). The fact that maternal age and education level were not associated with alcohol consumption during pregnancy, and that the alcohol consumers compared with the non-alcohol consumers were more likely to initiate breastfeeding postpartum further indicates a lack of any socio-demographic disparity as regard alcohol consumption during pregnancy in our sample of mothers. The finding that Irish national, compared with non-Irish national mothers were significantly more likely to consume alcohol during pregnancy is consistent with a recent Irish study (Barry *et al.*, 2007), which similarly found a higher prevalence of alcohol consumption among Irish national (79%) compared with non-European Union mothers (26%).

Just over one fifth of mothers (20.9%) smoked during pregnancy in this study, indicating some improvement from rates of 29% (Walsh *et al.*, 2007) and 27.9% (Donnelly *et al.*,

Table 6 Factors associated with non-compliance with peri-conceptual folic acid supplementation at the recommended time/no folic acid in a sample of women in Dublin ($n=450$)

Characteristic	Peri-conceptual FA at RT, n (%)	Non-compliance with FA at RT, n (%)	P-value ^a	OR ^b (95% CI)	Adjusted OR ^c (95% CI)	P-value ^d
Maternal age (years)						
> 34	54 (63.5)	31 (36.5)		1.0 ^e	1.0	
25–34	134 (50.2)	133 (49.8)		1.72 (1.04–2.85)	1.31 (0.72–2.39)	
<25	12 (12.2)	86 (87.8)	0.000	12.48 (5.9–26.38)	4.0 (1.64–9.77)	0.007
Highest maternal education qualification						
Third level degree/postgraduate	85 (57)	64 (43)		1.0	1.0	
Vocational/training course	63 (50.8)	61 (49.2)		1.28 (0.79–2.07)	0.78 (0.43–1.44)	
Primary/secondary	52 (29.4)	125 (70.6)	0.000	3.19 (2.01–5.04)	1.3 (0.72–2.35)	0.266
Planned current pregnancy						
Yes	186 (66)	96 (34)		1.0	1.0	
No	14 (8.3)	154 (91.7)	0.000	21.3(11.69–38.8)	13.24 (7–24.7)	0.000
Smoking during pregnancy						
No	182 (51.1)	174 (48.9)		1.0	1.0	
Yes	18 (19.1)	76 (80.9)	0.000	4.41 (2.53–7.68)	2.26 (1.13–4.52)	0.020
Pre-pregnancy BMI (kg/m²)^f						
≥25	61 (40.7)	89 (59.3)		1.0	1.0	
20–24.9	130 (51.4)	123 (48.6)		0.64 (0.43–0.97)	0.74 (0.44–1.23)	
<20	9 (19.1)	38 (80.9)	0.000	2.89 (1.3–6.41)	2.35 (0.87–6.28)	0.044
Added non-recommended condiments to infants' foods^g						
No	159 (52.6)	143 (47.4)		1.0	1.0	
Yes	40 (27.2)	107 (72.8)	0.000	2.97 (1.93–4.56)	1.82 (1.05–3.13)	0.030

Abbreviations: BMI, body mass index; CI, confidence interval; FA, folic acid; OR, odds ratio; RT, recommended time.

Results are expressed as number of subjects and percentage.

^aStatistical analysis by χ^2 statistics test. The Yates's continuity correction value was used for all 2×2 tables.

^bUnadjusted OR from logistic regression analysis.

^cOR adjusted for all other variables in the table using logistic regression analysis.

^dP-value from adjusted logistic regression model.

^e1.0 denotes the reference category.

^fBMI categories are in accordance with World Health Organisation (2003) definitions (<20 kg/m², underweight; 20–24.9 kg/m², normal weight; ≥25 kg/m², overweight). BMI values were derived from mothers' self-reported pre-pregnancy weight and height.

^gIncludes mothers who added non-recommended condiments to their infants' weaning foods ($n=449$), including ordinary gravy, butter, sauces, vegetable stock, salt and sugar/honey.

2008) reported in previous Irish studies. Maternal cigarette smoking during pregnancy is a well-documented international concern with rates from 10.1% in Switzerland (Lemola and Grob, 2007) to 29.5% in USA (McFarlane *et al.*, 1996) reported. Consistent with our findings (Abrahamsson and Ejertsson, 2000; Hensley Alford *et al.*, 2009), multiparous, younger women <25 years from lower social classes were identified as high-risk groups for smoking continuation during pregnancy. Lower maternal education level has been shown to be an important determinant of smoking during pregnancy in other studies (Ebrahim *et al.*, 2000; Gilman *et al.*, 2008); however, a stronger association between smoking in pregnancy and younger maternal age (<25 years) was indicated in the present study. To our knowledge, smoking behaviours by nationality has not been previously studied in Ireland. While Irish nationality was identified as a positive predictor of alcohol consumption during pregnancy, we also identified Irish nationality as being an independent

predictor of maternal cigarette smoking during pregnancy. As smoking during pregnancy is associated with approximately one third of low birth weight, one quarter of small for gestational age and one sixth of preterm births (Delpisheh *et al.*, 2007) and that 1% of all children are given a suboptimal introduction to life owing to their mother's smoking habits during pregnancy (Kallen, 2001), interventions and smoking cessation programmes that target these high-risk groups would be a worthwhile public health investment for Ireland.

A significant decrease in the number of cigarettes smoked per day and in the number of alcohol units consumed per week was found among the smokers and alcohol consumers, respectively, from pre- to during pregnancy in our study. In addition, a higher compliance rate with daily supplemental FA use was reported during pregnancy ($n=395$; 87.7%) than preconceptionally ($n=181$; 40.2%). Such trends towards improved compliance with public health recommendations

in pregnancy have also been reported by other investigators (Crozier *et al.*, 2009). Taken together, these data provide encouraging evidence that change in behaviour is possible for women, with a major behavioural change motivator likely to be a desire for the best possible pregnancy and infant health outcome. However, low compliance rates with public health recommendations concerning alcohol, smoking and FA use have also been reported in non-pregnant women during the months before becoming pregnant. Inskip *et al.* (2009) reports that only 2.9% of women complied fully with the recommendations on alcohol (≤ 4 units of alcohol/week) and FA intakes (≥ 400 $\mu\text{g}/\text{day}$) in 3 months before becoming pregnant, while only 44% of women reported taking any supplemental FA. The possibility that pre-pregnancy recommendations are not publicised widely enough must be considered. Given the high rate of unplanned pregnancies reported in national and international literature, greater public health efforts are needed to improve compliance with nutrition and lifestyle recommendations in women of childbearing age across the socio-demographic spectrum.

This study has a number of limitations. First, as this was a single-sited prospective study capturing the health behaviours of a convenience sample of women in west Dublin, the prevalence of health behaviours under study and the factors affecting mothers in our sample may differ across the island of Ireland. Second, owing to social desirability, it is known that some women may overestimate their compliance with public health recommendations during pregnancy (Ford *et al.*, 1997), which may have led, somewhat, to a potential inaccurate estimation of the actual prevalence rates reported. Such over/underestimations are, however, unlikely to differ from other similarly designed studies. Third, mothers' retrospective recollection of the timing of periconceptional FA supplementation might be at risk of recall bias. However, collection of the timing of FA uptake from the mothers' medical notes and clarification of any ambiguities that may have arisen may have increased the accuracy of these data. Finally, several factors that have been reported as predictors of maternal health behaviours, including the smoking behaviours of the partner and maternal grandmother (Lemola and Grob, 2008), were not collected in this study, which may somewhat limit interpretation of the multivariate analyses.

In conclusion, our data indicate that a large proportion of women in Ireland, in particular women of Irish nationality, continue to smoke and consume alcohol during pregnancy. Suboptimal compliance with peri-conceptional FA supplementation was a particular concern in this study with less than half of the sample potentially achieving optimal protection against NTDs. To decrease health inequalities in Ireland, our data suggest that lower socio-economic groups and younger women <25 years should be specifically targeted in smoking cessation and FA promotion campaigns. In addition, more effective public health approaches are required to convince, in particular, Irish national women

from all socio-economic groups, of the adverse fetal and maternal health risks associated with alcohol consumption during pregnancy.

Conflict of interest

The authors declare no conflict of interest.

Acknowledgements

We are greatly indebted to the 491 mothers who participated in this study. They are also extremely grateful to the clinical and administrative staff in the CWIUH for their invaluable support and practical help, particularly, during the study recruitment and follow-up phases. We confirm that this study was funded exclusively by the Dublin Institute of Technology, Dublin, Republic of Ireland.

References

- Abrahamsson A, Ejlertsson G (2000). Smoking patterns during pregnancy: differences in socioeconomic and health-related variables. *Eur J Public Health* **10**, 208–213.
- Banakar MK, Kudlur NS, George S (2009). Fetal alcohol spectrum disorder (FASD). *Indian J Pediatr* **76**, 1173–1175.
- Barry S, Kearney A, Daly S, Lawler E, McNamee E, Barry J (2007). *Coombe Women's Hospital Study of Alcohol, Smoking and Illicit Drug Use 1987–2005*. Coombe Women's Hospital and Department of Health & Children: Dublin http://www.drugsandalcohol.ie/6131/1/Coombe_Women%27s_Hospital_Alcohol_etc_Report_1987-2005.pdf (accessed January 2011).
- Bekkers RL, Eskes TK (1999). Periconceptional folic acid intake in Nijmegen, Netherlands. *Lancet* **353**, 292.
- Brough L, Rees GA, Crawford MA, Dorman EK (2009). Social and ethnic differences in folic acid use preconception and during early pregnancy in the UK: effect on maternal folate status. *J Hum Nutr Diet* **22**, 100–107.
- Busby A, Abramsky L, Dolk H, Armstrong B, Eurocrat Folic Acid Working Group (2005). Preventing neural tube defects in Europe: population based study. *BMJ* **330**, 574–575.
- Central Statistics Office Census (1996). *Volume 7, Occupations*. Central Statistics Office: Dublin.
- Cnattingius S (2004). The epidemiology of smoking during pregnancy: smoking prevalence, maternal characteristics, and pregnancy outcomes. *Nicotine Tob Res* **6**, S125–S140.
- Coombe Women and Infants University Hospital (2006). *Annual Clinical Reports: 2004, 2005 and 2006*. Coombe Women and Infants University Hospital: Dublin.
- Crozier SR, Robinson SM, Borland SE, Godfrey KM, Cooper C, Inskip HM, The SWS Study Group (2009). Do women change their health behaviours in pregnancy? Findings from the Southampton Women's Survey. *Paediatr Perinat Epidemiol* **23**, 446–453.
- Czeizel AE, Dudas I (1992). Prevention of the first occurrence of neural-tube defects by periconceptional vitamin supplementation. *N Engl J Med* **327**, 1832–1835.
- Daly SF, Kiely J, Clarke TA, Matthews TG (1992). Alcohol and cigarette use in a pregnant Irish population. *Ir Med J* **85**, 156–157.
- Delpisheh A, Kelly Y, Rizwan S, Attia E, Drammond S, Brabin BJ (2007). Population attributable risk for adverse pregnancy outcomes related to smoking in adolescents and adults. *Public Health* **121**, 861–868.

- Department of Health and Children (2008). Minister Wallace renews advice not to drink alcohol in pregnancy. <http://www.dohc.ie/press/releases/2008/20080908.html> (accessed January 2011).
- Department of Health, Republic of Ireland (1992). *Folic Acid and the Prevention of Neural Tube Defects. Report from an Expert Advisory Group*. Department of Health Publications Unit: Dublin.
- Donnelly JC, Cooley SM, Walsh TA, Sarkar R, Durnea U, Geary MP (2008). Illegal drug use, smoking and alcohol consumption in a low-risk Irish primigravid population. *J Perinat Med* **36**, 70–72.
- Ebrahim SH, Floyd RL, Merritt RK, Decoufle P, Holzman D (2000). Trends in pregnancy-related smoking rates in the United States, 1987–1996. *JAMA* **19**, 361–366.
- Food Safety Authority of Ireland (2006). *Report of the National Committee on Folic Acid Food Fortification*. Food Safety Authority of Ireland: Dublin http://www.fsai.ie/uploadedFiles/folic_acid.pdf (accessed January 2011).
- Food Safety Authority of Ireland (2008). *Report of the Implementation Group on Folic Acid Food Fortification to the Department of Health and Children*. Food Safety Authority of Ireland: Dublin <http://www.fsai.ie/details.aspx?id=7706> (accessed January 2011).
- Ford RP, Tappin DM, Schluter PJ, Wild CJ (1997). Smoking during pregnancy: how reliable are maternal self reports in New Zealand? *J Epidemiol Community Health* **51**, 246–251.
- Gilman SE, Breslau J, Subramanian SV, Hitsman B, Koenen KC (2008). Social factors, psychopathology, and maternal smoking during pregnancy. *Am J Public Health* **98**, 448–453.
- Health Service Executive (2008). *Give your Baby a Breather: Help and Advice on Giving Up Smoking during Pregnancy*. Health Service Executive: Dublin.
- Hensley Alford SM, Lappin RE, Peterson L, Johnson CC (2009). Pregnancy associated smoking behavior and six year postpartum recall. *Matern Child Health J* **13**, 865–872.
- Horta BL, Victora CG, Menezes AM, Halpern R, Barros FC (1997). Low birthweight, preterm births and intrauterine growth retardation in relation to maternal smoking. *Paediatr Perinat Epidemiol* **11**, 140–151.
- Inskip HM, Crozier SR, Godfrey KM, Borland SE, Cooper C, Robinson SM, The Southampton Women's Survey Study Group (2009). Women's compliance with nutrition and lifestyle recommendations before pregnancy: general population cohort study. *BMJ* **338**, b481.
- Kallen K (2001). The impact of maternal smoking during pregnancy on delivery outcome. *Eur J Public Health* **11**, 329–333.
- Kim MH, Han JY, Cho YJ, Ahn HK, Kim JO, Ryu HM et al. (2009). Factors associated with a positive intake of folic acid in the periconceptional period among Korean women. *Public Health Nutr* **12**, 468–471.
- Kirke PN, Daly LE, Elwood JH (1992). A randomised trial of low dose folic acid to prevent neural tube defects. The Irish Vitamin Study Group. *Arch Dis Child* **67**, 1442–1446.
- Kramer MS (2003). The epidemiology of adverse pregnancy outcomes: an overview. *J Nutr* **133**, 1592S–1596S.
- Krulwich CJ (2005). Alcohol consumption during pregnancy. *Annu Rev Nurs Res* **23**, 101–134.
- Kyrklund-Blomberg NB, Cnattingius S (1998). Preterm birth and maternal smoking: risks related to gestational age and onset of delivery. *Am J Obstet Gynecol* **179**, 1051–1055.
- Landgren M, Svensson L, Stromland K, Andersson Gronlund M (2010). Prenatal alcohol exposure and neurodevelopmental disorders in children adopted from eastern Europe. *Pediatrics* **125**, e1178–e1185.
- Lemola S, Grob A (2007). Drinking and smoking in pregnancy: what questions do Swiss physicians ask? *Swiss Med Wkly* **137**, 66–69.
- Lemola S, Grob A (2008). Smoking cessation during pregnancy and relapse after childbirth: the impact of the grandmother's smoking status. *Matern Child Health J* **12**, 525–533.
- McFarlane J, Parker B, Soeken K (1996). Physical abuse, smoking, and substance use during pregnancy: prevalence, interrelationships, and effects on birth weight. *J Obstet Gynecol Neonatal Nurs* **25**, 313–320.
- McGuire M, Cleary B, Sahn L, Murphy DJ (2010). Prevalence and predictors of periconceptional folic acid uptake- prospective cohort study in an Irish urban obstetric population. *Hum Reprod* **25**, 535–543.
- McMillan H, Smaarani S, Walsh T, Khawaja N, Collins C, Byrne P et al. (2006). Smoking and alcohol in pregnancy. Survey in the immediate post-partum period. *Ir Med J* **99**, 283.
- Monica G, Lilja C (1995). Placenta previa, maternal smoking and recurrence risk. *Acta Obstet Gynecol Scand* **74**, 341–345.
- MRC Vitamin Study Research Group (1991). Prevention of neural tube defects. Results of the Medical Research Council Vitamin Study. MRC Vitamin Study Research Group. *Lancet* **338**, 131–137.
- Office of Population Census and Surveys (1991). *Standard Occupational Classification, Volume 3, Social Classifications and Coding Methodology*. Her Majesty's Stationery Office: London.
- Pollack H, Lantz PM, Frohna JG (2000). Maternal smoking and adverse birth outcomes among singletons and twins. *Am J Public Health* **90**, 395–400.
- Ray JG, Singh G, Burrows RF (2004). Evidence for suboptimal use of periconceptional folic acid supplements globally. *BJOG* **111**, 399–408.
- Tarrant RC (2008). Patient-administered questionnaire. In: *An investigation of the diets of infants born in Ireland during the first 6 months of life*. PhD thesis, pp. 434. Dublin Institute of Technology: Dublin. Available at: <http://arrow.dit.ie/cgi/viewcontent.cgi?article=1048&context=sciendoc> (accessed January 2011).
- Tarrant RC, Younger KM, Sheridan-Pereira M, White MJ, Kearney JM (2010a). The prevalence and determinants of breast-feeding initiation and duration in a sample of women in Ireland. *Public Health Nutr* **13**, 760–770.
- Tarrant RC, Younger KM, Sheridan-Pereira M, White MJ, Kearney JM (2010b). Factors associated with weaning practices in term infants: a prospective observational study in Ireland. *Br J Nutr* **104**, 1544–1554.
- Walsh T, O'Broin S, Cooley S, Donnelly J, Collins C, McMillan H et al. (2007). Maternal folate status and neural tube defects in Ireland: the need for a national food fortification program. *Ir Med J* **100**, 469–472.
- Ward M, Hutton J, McDonnell R, Bachir N, Scallan E, O'Leary M et al. (2004). Folic acid supplements to prevent neural tube defects: trends in East of Ireland 1996–2002. *Ir Med J* **97**, 274–276.
- Wilcox AJ, Lie RT, Solvoll K, Taylor J, McConaughy DR, Abyholm F et al. (2007). Folic acid supplements and risk of facial clefts: national population based case-control study. *BMJ* **334**, 464.
- Wisborg K, Kesmodel U, Henriksen TB, Olsen SF, Secher NJ (2000). A prospective study of smoking during pregnancy and SIDS. *Arch Dis Child* **83**, 203–206.
- World Health Organisation (2003). *Process for a WHO Global Strategy on Diet, Physical Activity and Health*. World Health Organisation: Geneva. WHA55.23.
- World Health Organisation (2006). Framework for alcohol policy in the WHO European Region. <http://www.euro.who.int/document/e88335.pdf> (accessed January 2011).