



2010-01-01

# Schrödinger Operators with $\delta'$ Interactions and the Krein–Stieltjes String

Aleksey Kostenko

*Dublin Institute of Technology, [aleksey.kostenko@dit.ie](mailto:aleksey.kostenko@dit.ie)*

M. M. Malamud

*Institute of Applied Mathematics and Mechanics NASU, Ukraine*

Follow this and additional works at: <http://arrow.dit.ie/scschmatart>

 Part of the [Mathematics Commons](#)

## Recommended Citation

Kostenko, A., Malamud, M.: Schrödinger Operators with  $\delta'$  Interactions and the Krein–Stieltjes String. *Doklady Mathematics*, Vol. 81, no. 3, 2010, pp.342-347.

This Article is brought to you for free and open access by the School of Mathematics at ARROW@DIT. It has been accepted for inclusion in Articles by an authorized administrator of ARROW@DIT. For more information, please contact [yvonne.desmond@dit.ie](mailto:yvonne.desmond@dit.ie), [arrow.admin@dit.ie](mailto:arrow.admin@dit.ie), [brian.widdis@dit.ie](mailto:brian.widdis@dit.ie).



This work is licensed under a [Creative Commons Attribution-NonCommercial-Share Alike 3.0 License](#)



# Schrödinger Operators with $\delta'$ -Interactions and the Krein–Stieltjes String<sup>1</sup>

A. S. Kostenko<sup>a, b</sup> and M. M. Malamud<sup>b</sup>

Presented by Academician V.P. Maslov October 21, 2009

Received November 9, 2009

**Abstract**—We investigate one dimensional symmetric Schrödinger operator  $H_{X, \beta}$  with  $\delta'$ -interactions of strength  $\beta = \{\beta_n\}_{n=1}^{\infty} \subset \mathbb{R}$  on a discrete set  $X = \{x_n\}_{n=1}^{\infty} \subset [0, b)$ ,  $b \leq +\infty$  ( $x_n \uparrow b$ ). We consider  $H_{X, \beta}$  as an extension of the minimal operator  $H_{\min} := -d^2/dx^2 \upharpoonright W_0^{2,2}(\mathbb{R} \setminus X)$  and study its spectral properties in the framework of the extension theory by using the technique of boundary triplets and the corresponding Weyl functions. The construction of a boundary triplet for  $H_{\min}^*$  is given in the case  $d_* := \inf_{n \in \mathbb{N}} |x_n - x_{n-1}| = 0$ . We show that spectral properties like self-adjointness, lower semiboundedness, nonnegativity, and discreteness of the spectrum of the operator  $H_{X, \beta}$  correlate with the corresponding properties of a certain Jacobi matrix. In the case  $\beta_n > 0$ ,  $n \in \mathbb{N}$ , these matrices form a subclass of Jacobi matrices generated by the Krein–Stieltjes strings. The connection discovered enables us to obtain simple conditions for the operator  $H_{X, \beta}$  to be self-adjoint, lower semibounded and discrete. These conditions depend significantly not only on  $\beta$  but also on  $X$ . Moreover, as distinct from the case  $d_* > 0$ , the spectral properties of Hamiltonians with  $\delta$ - and  $\delta'$ -interactions in the case  $d_* = 0$  substantially differ.

DOI: 10.1134/S1064562410030026

Let  $\mathcal{I} := [0, b)$ ,  $0 < b \leq +\infty$ , and let  $X = \{x_n\}_{n=0}^{\infty} \subset \mathcal{I}$  be a strictly increasing sequence such that  $\lim_{n \rightarrow \infty} x_n = b$  and

$x_0 := 0$ . Let also  $\beta := \{\beta_n\}_{n=1}^{\infty} \subset \mathbb{R}$ .

Consider a symmetric operator  $H_{X, \beta}^0$  in  $L^2(\mathcal{I})$  defined by the differential expression  $-\frac{d^2}{dx^2}$  on functions  $f \in W_{\text{comp}}^{2,2}(\mathcal{I} \setminus X)$  satisfying the following boundary conditions at the points  $x_n \in X$ :

$$\begin{aligned} f'(0) &= 0, \quad f'(x_n+) = f'(x_n-), \\ f(x_n+) - f(x_n-) &= \beta_n f(x_n), \quad n \in \mathbb{N}. \end{aligned} \quad (1)$$

Let us denote its closure by  $H_{X, \beta}$ . The operator  $H_{X, \beta}$  is interpreted as the Hamiltonian with  $\delta'$ -interactions of strength  $\beta_n$  at the centers  $x_n$  [2] and, as a rule, is associated with the formal differential expression

$$l_{X, \beta} := -\frac{d^2}{dx^2} + \sum_{x_n \in X} \beta_n(\cdot, \delta'_n) \delta'_n, \quad x \in \mathcal{I}, \quad (2)$$

where  $\delta_n := \delta(x - x_n)$  and  $\delta(\cdot)$  is the Dirac delta-function.

Spectral properties of the operator  $H_{X, \beta}$  are widely studied in the case  $d_* := \inf_{n \in \mathbb{N}} d_n > 0$ , where  $d_n := x_n - x_{n-1}$ . Thus, it is known that the operator  $H_{X, \beta}$  is self-adjoint [2] and its spectrum is not discrete in this case. Further results (investigation of spectrum, resolvent comparability etc.) as well as a comprehensive list of references can be found in [2, 14].

The cases  $d_* = 0$  and  $d_* > 0$  are significantly different. Recently, it has been found ([5, Theorem 4.7]) that  $H_{X, \beta}$  is self-adjoint for any  $\beta$  if  $\mathcal{I} = \mathbb{R}_+$  and  $d_* = 0$ . To the best of our knowledge, other spectral properties of the operator  $H_{X, \beta}$  in the case  $d_* = 0$  have not been studied yet. Let us also mention the recent papers [3, 15] dealing with spectral properties of Hamiltonians with  $\delta'$ -interactions on compact subsets of  $\mathbb{R}$  with Lebesgue measure zero.

The main aim of this note is the spectral analysis of the Hamiltonian  $H_{X, \beta}$  in the case  $d_* = 0$ . We investigate the operator  $H_{X, \beta}$  in the framework of extension theory of symmetric operators using the concept of boundary triplets and the corresponding Weyl functions (see [7, 9]). Let us note that this approach was first applied by Kochubei [12] in the study of operators

<sup>1</sup> The article was translated by the authors.

<sup>a</sup> Dublin Institute of Technology, Ireland

<sup>b</sup> Institute of Applied Mathematics and Mechanics NASU, Ukraine

e-mail: duzer80@gmail.com, mmm@telenet.dn.ua

with point interactions. He investigated Hamiltonians with  $\delta$ -interactions in the case  $d_* > 0$ . Let us stress that the main difficulty of this approach is the construction of an adequate boundary triplet.

In this note, we present a corresponding construction in the case  $d_* = 0$  and show that the spectral properties like self-adjointness, lower semiboundedness, and discreteness of the spectrum of the operator  $H_{X,\beta}$  correlate with the corresponding spectral properties of the Jacobi matrix  $B_{X,\beta}$  defined by (15) (see below). It turns out that this class of matrices is closely connected with the class of Krein–Stieltjes string operators. Namely, if  $\beta_n > 0, n \in \mathbb{N}$ , then this class of matrices is a subclass of Krein–Stieltjes operators (see Section 5). Discovered connection enables us to obtain simple criteria for the operator  $H_{X,\beta}$  to be self-adjoint, lower semi-bounded, and discrete. These conditions depend substantially not only on  $\beta$  but also on  $X$ . Moreover, as distinct from the case  $d_* > 0$ , the spectral properties of Hamiltonians with  $\delta$ - and  $\delta'$ -interactions in the case  $d_* = 0$  differ completely.

Detailed treatment of the results discussed in this note are given in [11].

### 1. BOUNDARY TRIPLETS FOR DIRECT SUMS AND WEYL FUNCTIONS

Let  $A$  be a closed symmetric operator with dense domain  $\mathfrak{D}(A)$  in a Hilbert space  $\mathfrak{H}$  and equal deficiency indices  $n_+(A) = n_-(A)$  ( $n_\pm(A) := \dim(\mathfrak{N}_{\pm i}), \mathfrak{N}_z := \ker(A^* - zI)$ ).

**Definition 1** ([7]). A collection  $\Pi = \{\mathcal{H}, \Gamma_0, \Gamma_1\}$ , where  $\mathcal{H}$  is an auxiliary Hilbert space,  $\Gamma_0$  and  $\Gamma_1$  are linear mappings from  $\mathfrak{D}(A^*)$  to  $\mathcal{H}$ , is called a boundary triplet for the operator  $A^*$  if the abstract Green identity holds

$$(A^*f, g) - (f, A^*g) = (\Gamma_1 f, \Gamma_0 g)_{\mathcal{H}} - (\Gamma_0 f, \Gamma_1 g)_{\mathcal{H}} \quad (3)$$

for all  $f, g \in \mathfrak{D}(A^*)$ ,

and the mapping  $\Gamma: f \rightarrow \{\Gamma_0 f, \Gamma_1 f\}$  from  $\mathfrak{D}(A^*)$  to  $\mathcal{H} \oplus \mathcal{H}$  is surjective.

A boundary triplet is not unique and, moreover, it enables one to parameterize the set  $\text{Ext}_A$  of all proper extensions  $\tilde{A}$  ( $A \subset \tilde{A} \subset A^*$ ) of the operator  $A$  in terms of abstract boundary conditions. Let  $A_j := A^*|_{\ker(\Gamma_j)}$ ,  $j \in \{0, 1\}$ . It is known that  $A_j = A_j^*$ ,  $j \in \{0, 1\}$ .

**Proposition 1** ([7]). Let  $\Pi = \{\mathcal{H}, \Gamma_0, \Gamma_1\}$  be a boundary triplet for  $A^*$ . Then the mapping

$$(\text{Ext}_A \ni) \tilde{A} \rightarrow \Gamma \mathfrak{D}(\tilde{A})$$

$$= \{ \{ \Gamma_0 f, \Gamma_1 f \} : f \in \mathfrak{D}(\tilde{A}) \} =: \Theta \in \mathfrak{C}(\mathcal{H}) \quad (4)$$

establishes a bijective correspondence between  $\text{Ext}_A$  and the set  $\mathfrak{C}(\mathcal{H})$  of closed linear relations in  $\mathcal{H}$ . Denote  $A_\Theta := \tilde{A}$ , where  $\Theta$  is determined by Eq. (4). Then the following statements hold

- (i)  $A_\Theta = A_\Theta^* \Leftrightarrow \Theta = \Theta^*$ ;
- (ii)  $A_\Theta$  is symmetric  $\Leftrightarrow \Theta$  is symmetric. Moreover,  $n_\pm(A_\Theta) = n_\pm(\Theta)$ ;
- (iii)  $\mathfrak{D}(A_\Theta) \cap \mathfrak{D}(A_0) = \mathfrak{D}(A) \Leftrightarrow \Theta \in \mathfrak{C}(\mathcal{H})$ . In this case  $\Theta$  is called a boundary operator and relation (4) becomes

$$A_\Theta = A^* \upharpoonright \ker(\Gamma_1 - \Theta \Gamma_0). \quad (5)$$

**Definition 2** ([9]). Let  $\Pi = \{\mathcal{H}, \Gamma_0, \Gamma_1\}$  be a boundary triplet for  $A^*$ . Operator valued function  $M(\cdot): \rho(A_0) \rightarrow [\mathcal{H}]$  defined by

$$\Gamma_1 f_z = M(z) \Gamma_0 f_z, \quad f_z \in \ker(A^* - z), \quad z \in \rho(A_0), \quad (6)$$

is called the Weyl function corresponding to the boundary triplet  $\Pi$ .

Further, let  $S_n$  be a closed densely defined symmetric operator in  $\mathfrak{H}_n$  such that  $n_+(S_n) = n_-(S_n) \leq \infty, n \in \mathbb{N}$ . In the Hilbert space  $\mathfrak{H} := \bigoplus_{n=1}^\infty \mathfrak{H}_n$ , consider the operator  $A := \bigoplus_{n=1}^\infty S_n$ . Let also  $\Pi_n = \{\mathcal{H}_n, \Gamma_0^{(n)}, \Gamma_1^{(n)}\}$  be a boundary triplet for the operator  $S_n^*$ . Define a direct sum  $\Pi := \bigoplus_{n=1}^\infty \Pi_n$  of boundary triplets  $\Pi_n$  by setting

$$\mathcal{H} = \bigoplus_{n=1}^\infty \mathcal{H}_n, \quad \Gamma_j := \bigoplus_{j=1}^\infty \Gamma_j^{(n)}, \quad \Gamma_j: \mathfrak{D}(A^*) \rightarrow \mathcal{H}, \quad j \in \{0, 1\}. \quad (7)$$

Green’s formulae (3) for operators  $S_n^*, n \in \mathbb{N}$ , yield validity of Green’s formula for the operator  $A^* = \bigoplus_{n=1}^\infty S_n$  and also vectors  $f = \bigoplus_{n=1}^\infty f_n, g = \bigoplus_{n=1}^\infty g_n \in \mathfrak{D}(\Gamma_0) \cap \mathfrak{D}(\Gamma_1) \subset \mathfrak{D}(A^*)$ , where

$$\mathfrak{D}(\Gamma_j) := \left\{ \varphi = \bigoplus_{n=1}^\infty \varphi_n : \sum_{n \in \mathbb{N}} \|\Gamma_j^{(n)} \varphi_n\|_{\mathcal{H}_n}^2 < \infty \right\}.$$

It is shown in [12] that a boundary triplet  $\Pi$  for the operator  $A^* = \bigoplus_{n=1}^\infty S_n^*$  can be chosen in the form (7) if

there exist  $\varepsilon > 0$  and a sequence  $\{\rho_n\}_{n=1}^\infty \subset [\varepsilon, \varepsilon^{-1}]$  such that all pairs  $\{\rho_n S_n, \rho_n S_{n0}\}, n \in \mathbb{N}$ , are unitary equivalent to  $\{S_1, S_{10}\}$ . However, it is not difficult to give an example of the operator  $A = \bigoplus_{n=1}^\infty S_n$  for which a triplet (7) is not a boundary triplet for  $A^*$ . The reason is the following. The mapping  $\Gamma = \{\Gamma_0, \Gamma_1\}: \mathfrak{D}(A^*) \rightarrow \mathcal{H} \oplus \mathcal{H}$  may not fulfill the following necessary conditions:

$$\mathfrak{D}(\Gamma_0) \cap \mathfrak{D}(\Gamma_1) = \mathfrak{D}(A^*), \quad \text{ran}(\Gamma) = \mathcal{H} \oplus \mathcal{H}.$$

Next theorem provides criteria for the collection  $\Pi = \{\mathcal{H}, \Gamma_0, \Gamma_1\}$  of the form (7) to be a boundary triplet for the operator  $A^* = \bigoplus_{n=1}^{\infty} S_n^*$ .

**Theorem 1.** Let  $A = \bigoplus_{n=1}^{\infty} S_n$ ,  $\Pi_n = \{\mathcal{H}_n, \Gamma_0^{(n)}, \Gamma_1^{(n)}\}$

be a boundary triplet for the operator  $S_n^*$ ,  $n \in \mathbb{N}$ , and let  $\Pi = \{\mathcal{H}, \Gamma_0, \Gamma_1\}$  be defined by (7). Then:

(i) the triplet  $\Pi = \{\mathcal{H}, \Gamma_0, \Gamma_1\}$  is a boundary relation (in the sense of [8]) for the operator  $A^*$ ;

(ii) the collection  $\Pi = \{\mathcal{H}, \Gamma_0, \Gamma_1\}$  is a boundary triplet for the operator  $A^*$  precisely when the mappings  $\Gamma_j$  are bounded, i.e.,

$$\sup_{n \in \mathbb{N}} \|\Gamma_0^{(n)}\| < \infty, \quad \sup_{n \in \mathbb{N}} \|\Gamma_1^{(n)}\| < \infty; \tag{8}$$

(iii) the collection  $\Pi = \{\mathcal{H}, \Gamma_0, \Gamma_1\}$  is a boundary triplet for the operator  $A^*$  if and only if the following conditions are satisfied

$$\sup_n \|M_n(i)\| < \infty, \quad \sup_n \|(\operatorname{Im} M_n(i))^{-1}\| < \infty. \tag{9}$$

Here  $M_n(\cdot)$  is the Weyl function corresponding to the triplet  $\Pi_n$ .

(iv) If  $a_0 = \bar{a}_0 (\in \mathbb{R})$  is the point of a regular type of the operator  $A$ , then the collection  $\Pi = \{\mathcal{H}, \Gamma_0, \Gamma_1\}$  is a boundary triplet for the operator  $A^*$  precisely when

$$\sup_{n \in \mathbb{N}} \|M_n(a_0)\| < \infty, \quad \sup_{n \in \mathbb{N}} \|(M_n'(a_0))^{-1}\| < \infty. \tag{10}$$

Theorem 1 enables one to regularize an arbitrary collection of boundary triplets  $\tilde{\Pi}_n = \{\tilde{\mathcal{H}}_n, \tilde{\Gamma}_0^{(n)}, \tilde{\Gamma}_1^{(n)}\}$  in such a way that the direct sum of the regularized triplets is a boundary triplet for the operator  $A^*$ . Namely, assume that  $Q_n = Q_n^* \in [\tilde{\mathcal{H}}_n]$  and  $R_n \in [\tilde{\mathcal{H}}_n]$  with  $R_n^{-1} \in [\tilde{\mathcal{H}}_n]$ . We set

$$\Gamma_0^{(n)} := R_n \tilde{\Gamma}_0^{(n)}, \quad \Gamma_1^{(n)} := (R_n^{-1})^* (\tilde{\Gamma}_1^{(n)} - Q_n \tilde{\Gamma}_0^{(n)}). \tag{11}$$

**Corollary 1.** Let  $\tilde{\Pi}_n = \{\tilde{\mathcal{H}}_n, \tilde{\Gamma}_0^{(n)}, \tilde{\Gamma}_1^{(n)}\}$  be a boundary triplet for  $S_n^*$ , and let  $\tilde{M}_n(\cdot)$  be the corresponding Weyl function,  $n \in \mathbb{N}$ . If

$$\begin{aligned} \sup_{n \in \mathbb{N}} \{ \|(R_n^{-1})^* (\tilde{M}_n(i) - Q_n) R_n^{-1}\| < \infty, \\ \sup_{n \in \mathbb{N}} \|R_n (\operatorname{Im} (\tilde{M}_n(i)))^{-1} R_n^*\| < \infty, \end{aligned} \tag{12}$$

then the direct sum  $\Pi = \bigoplus_{n=1}^{\infty} \Pi_n$  (see (7)) of triplets  $\Pi_n =$

$\{\mathcal{H}_n, \Gamma_0^{(n)}, \Gamma_1^{(n)}\}$  defined by (11) is a boundary triplet for

the operator  $A^* = \bigoplus_{n=1}^{\infty} S_n^*$ .

**Remark 1.** The existence of the regularization (11) was first established in [13, Theorem 5.3]. Namely, in

relations (11) it is taken  $Q_n := \operatorname{Re} \tilde{M}_n(i)$  and  $R_n := \sqrt{\operatorname{Im} \tilde{M}_n(i)}$ ,  $n \in \mathbb{N}$  (cf. [13]). However, let us stress that the latter regularization is not suitable for our aims since it does not lead to a parametrization of the Hamiltonians  $H_{X, \beta}$  by Jacobi matrices.

## 2. BOUNDARY TRIPLETS FOR SCHRODINGER OPERATORS

Consider the minimal (closed) symmetric operator

$$H := H_{\min} := -\frac{d^2}{dx^2}, \quad \mathfrak{D}(H_{\min}) = W_0^{2,2}(\mathcal{J} \setminus X). \tag{13}$$

It is clear that  $n_{\pm}(H_{\min}) = \infty$  and the operator  $H_{\min}^*$  is defined by the same differential expression on the domain  $\mathfrak{D}(H_{\min}^*) = W^{2,2}(\mathcal{J} \setminus X)$ . It is easy to see that  $H_{\min} =$

$$\bigoplus_{n \in \mathbb{N}} H_n, \quad \text{where } H_n := -\frac{d^2}{dx^2}, \quad \mathfrak{D}(H_n) = W_0^{2,2}[x_{n-1}, x_n].$$

Define the mappings

$$\begin{aligned} \Gamma_0^{(n)} f &:= \begin{pmatrix} d_n^{1/2} f(x_{n-1}+) \\ -d_n^{1/2} f(x_n-) \end{pmatrix}, \\ \Gamma_1^{(n)} f &:= \begin{pmatrix} \frac{d_n f'(x_{n-1}+) + (f(x_{n-1}+) - f(x_n-))}{d_n^{3/2}} \\ \frac{d_n f'(x_n-) + (f(x_{n-1}+) - f(x_n-))}{d_n^{3/2}} \end{pmatrix}. \end{aligned} \tag{14}$$

A straightforward calculation shows that the triplet  $\Pi_n = \{\mathbb{C}^2, \Gamma_0^{(n)}, \Gamma_1^{(n)}\}$  is a boundary triplet for the operator  $H_n^*$ . Moreover, we have the following

**Theorem 2.** Let  $d^* := \sup_{n \in \mathbb{N}} d_n < \infty$  and let  $\Gamma_0^{(n)}, \Gamma_1^{(n)}$

be defined by (14). Then the triplet  $\Pi = \bigoplus_{n=1}^{\infty} \Pi_n = \{\mathcal{H}, \Gamma_0, \Gamma_1\}$  defined by (7) is a boundary triplet for the operator  $H_{\min}^*$ .

**Proof.** The condition  $d^* < \infty$  yields that  $z = 0$  is a point of a regular type for the operator  $H_{\min}$ . To complete the proof of Theorem 2 it suffices to check (10) at the point  $z = 0$ .

## 3. PARAMETRIZATION OF HAMILTONIANS WITH $\delta'$ -INTERACTIONS

For the remainder of this note, we can assume without loss of generality that  $\beta_n \neq 0$  for all  $n \in \mathbb{N}$ . Further, consider the semi-infinite Jacobi matrix

$$B_{X,\beta} := \begin{pmatrix} d_1^{-2} & d_1^{-2} & 0 & 0 & 0 & \dots \\ d_1^{-2} & \frac{d_1^{-1}}{\beta_1} + d_1^{-2} & \frac{d_1^{-1/2}d_2^{-1/2}}{\beta_1} & 0 & 0 & \dots \\ 0 & \frac{d_1^{-1/2}d_2^{-1/2}}{\beta_1} & \frac{d_2^{-1}}{\beta_1} + d_2^{-2} & d_2^{-2} & 0 & \dots \\ 0 & 0 & d_2^{-2} & \frac{d_2^{-1}}{\beta_2} + d_2^{-2} & \frac{d_2^{-1/2}d_3^{-1/2}}{\beta_2} & \dots \\ 0 & 0 & 0 & \frac{d_2^{-1/2}d_3^{-1/2}}{\beta_2} & \frac{d_3^{-1}}{\beta_2} + d_3^{-2} & \dots \\ \dots & \dots & \dots & \dots & \dots & \dots \end{pmatrix}. \tag{15}$$

In  $\ell^2(\mathbb{N})$ , the minimal symmetric operator is naturally associated with the matrix  $B_{X,\beta}$  (see [4]). We denote it also by  $B_{X,\beta}$ .

The following Lemma shows that the boundary operator  $\Theta$  parameterizing  $H_{X,\beta}$  via (5) is a Jacobi matrix of the form (15).

**Lemma 1.** *Let  $\Pi = \{\mathcal{H}, \Gamma_0, \Gamma_1\}$  be a boundary triplet for  $H_{\min}^*$  defined in Theorem 2. Then  $H_{X,\beta} = H_{B_{X,\beta}}$ , that is*

$$H_{X,\beta} = H_{\min}^* \upharpoonright \mathfrak{D}(H_{X,\beta}),$$

$$\mathfrak{D}(H_{X,\beta}) = \{f \in W^{2,2}(\mathbb{R} \setminus X) : \Gamma_1 f = B_{X,\beta} \Gamma_0 f\}.$$

Theorem 2, Lemma 1 and some results from [7, 9] enable us to establish the following connection between spectral properties of the Hamiltonian  $H_{X,\beta}$  and the Jacobi matrix (boundary operator)  $B_{X,\beta}$ .

**Theorem 3.** *Assume that  $d^* < \infty$ . Then:*

- (i) *the following relation holds  $n_{\pm}(H_{X,\beta}) = n_{\pm}(B_{X,\beta})$ ;*
- (ii) *the operator  $H_{X,\beta}$  is lower semibounded precisely when the operator  $B_{X,\beta}$  is lower semibounded;*
- (iii) *if  $n_{\pm}(H_{X,\beta}) = 0$ , i.e.,  $H_{X,\beta} = H_{X,\beta}^*$ , then the spectrum of the operator  $H_{X,\beta}$  is discrete precisely when  $\lim_{n \rightarrow \infty} d_n = 0$  and the spectrum of the operator  $B_{X,\beta}$  is discrete.*

#### 4. A CONNECTION WITH THE KREIN–STIELTJES STRING

It is clear that the parameterizing matrix of the form (15) admits the following factorization

$$B_{X,\beta} = R_X^{-1}(I + U)D_{X,\beta}^{-1}(I + U^*)R_X^{-1}, \tag{16}$$

where  $U$  is the unilateral shift operator in  $\ell^2(\mathbb{N})$  and

$$D_{X,\beta} := \bigoplus_{n=1}^{\infty} \begin{pmatrix} d_n & 0 \\ 0 & \beta_n \end{pmatrix}, \quad R_X := \bigoplus_{n=1}^{\infty} \begin{pmatrix} \sqrt{d_n} & 0 \\ 0 & \sqrt{d_n} \end{pmatrix}. \tag{17}$$

We put  $l_{2n-1} := d_n, l_{2n} := \beta_n, m_{2n-1} = m_{2n} := d_n, n \in \mathbb{N}$ . In the case  $\beta_n > 0, n \in \mathbb{N}$ , the difference equation associated with the matrix  $B_{X,\beta}$  describes the motion of an inhomogeneous string (Krein–Stieltjes string) with mass distribution  $\mathcal{M}(y) = \sum_{y_n < y} m_n$ , where  $y_n - y_{n-1} := l_n$

and  $y_0 := 0$ . This class of matrices is studied sufficiently well. In particular, criterion of self-adjointness (Hamburger’s Theorem [1, Theorem 0.5]) and criterion of discreteness of spectrum (the Kac–Krein Theorem [10]) are known and are formulated in terms of sequences  $l = \{l_n\}_{n=1}^{\infty}$  and  $m = \{m_n\}_{n=1}^{\infty}$ . We hardly exploit these criteria as well as Theorem 3 in the following sections.

#### 5. SELF-ADJOINTNESS

Combining Theorem 3 (i) with the representation (16), (17), we obtain

**Theorem 4.** *Deficiency indices of the operator  $H_{X,\beta}$  are equal and are not greater than one,  $n_{+}(H_{X,\beta}) = n_{-}(H_{X,\beta}) \leq 1$ . Furthermore,  $H_{X,\beta}$  is self-adjoint if and only if at least one of the following conditions hold:*

- (1)  $\sum_{n=1}^{\infty} d_n = \infty$ , i.e.  $\mathcal{F} = \mathbb{R}_+$ ;
- (2)  $\sum_{n=1}^{\infty} \left[ d_{n+1} \left| \sum_{i=1}^n (\beta_i + d_i) \right|^2 \right] = \infty$ .

**Remark 3.** In the case of positive strengths  $\beta_n$ , Theorem 4 follows from Hamburger’s Theorem [1]. In the case of arbitrary strengths, we use the formulae for  $P_n(0)$  and  $Q_n(0)$  [1, p. 291] and the self-adjointness criterion [4, Lemma VII.1.5].

Theorem 4 immediately yields the following result of Buschmann–Stolz–Weidmann [5, Theorem 4.7].

**Corollary 2** ([5]). *If  $\mathcal{F} = \mathbb{R}_+$ , then the operator  $H_{X,\beta}$  is self-adjoint.*

6. LOWER SEMIBOUNDED HAMILTONIANS

Using Theorem 3(ii) and the form of the matrix  $B_{X,\beta}$ , we arrive at the following result.

**Proposition 2.** *For the operator  $H_{X,\beta}$  to be lower semibounded it is necessary that*

$$\beta_n^{-1} \geq -C_1 d_n - \min\{d_n^{-1}, d_{n+1}^{-1}\}, \quad n \in \mathbb{N} \quad (18)$$

and sufficient that

$$\beta_n^{-1} \geq -C_2 \min\{d_n, d_{n+1}\}, \quad n \in \mathbb{N}. \quad (19)$$

Here  $C_1$  and  $C_2$  are positive constants independent of  $n \in \mathbb{N}$ .

**Corollary 3.** *If  $0 < d_* \leq d^* < \infty$ , then the operator  $H_{X,\beta}$  is lower semi-bounded precisely when  $\inf_{n \in \mathbb{N}} \beta_n^{-1} > -\infty$ .*

7. DISCRETENESS OF SPECTRUM

We set  $\beta_n^- := \beta_n$  if  $\beta_n < 0$  and  $\beta_n^- := -\infty$  if  $\beta_n > 0$ . Applying the discreteness criterion [10] to the matrix  $B_{X,\beta}$  and combining this with Theorem 3 (iii), we arrive at the following necessary conditions for discreteness of the spectrum of the operator  $H_{X,\beta}$ .

**Proposition 3.** *Let  $\mathcal{F} = \mathbb{R}_+$  and  $d_n \rightarrow 0$ . The spectrum of the operator  $H_{X,\beta}$  is not discrete if at least one of the following conditions hold:*

- (i)  $\lim_{n \rightarrow \infty} x_n \sum_{j=n}^{\infty} d_j^3 > 0$ ;
- (ii)  $\beta_n \geq -C d_n^3, n \in \mathbb{N}, C > 0$ ;
- (iii)  $\beta_n^- \leq -C(d_n^{-1} + d_{n+1}^{-1}), n \in \mathbb{N}, C > 0$ .

**Corollary 4.** *If either  $\beta_n > 0$  for all  $n \in \mathbb{N}$  or  $\{d_n\}_{n=1}^{\infty} \notin \beta(\mathbb{N})$ , then the spectrum of the operator  $H_{X,\beta}$  is not discrete.*

Proposition 3 shows that the spectrum of the operator  $H_{X,\beta}$  is discrete in the very exceptional cases. For instance, the spectrum is not discrete if either  $\beta \subset \mathbb{R}_+$  or if there exists a subsequence  $\{\beta_{n_k}\} \subset \mathbb{R}_-$ , which tends to  $-\infty$  sufficiently fast. Furthermore,  $d_n$  must tend to zero sufficiently fast for the spectrum to be discrete. Sufficient conditions for discreteness of the spectrum of the Hamiltonian  $H_{X,\beta}$  are given in the following

**Theorem 5.** *Assume that  $\beta_n + d_n \geq 0$  for all  $n \in \mathbb{N}$ .*

(i) *If  $b < +\infty$ , and  $X$  and  $\beta$  are such that the operator  $H_{X,\beta}$  is self-adjoint, then the spectrum of  $H_{X,\beta}$  is discrete if and only if*

$$\lim_{n \rightarrow \infty} (b - x_n) \sum_{j=1}^{\infty} (\beta_j + d_j) = 0. \quad (20)$$

(ii)  *$\mathcal{F} = \mathbb{R}_+$ , then the spectrum of  $H_{X,\beta}$  is discrete precisely when*

$$\lim_{n \rightarrow \infty} x_n \sum_{j=n}^{\infty} d_j^3 = 0, \quad \lim_{n \rightarrow \infty} x_n \sum_{j=n}^{\infty} (\beta_j + d_j) = 0. \quad (21)$$

**Remark 4.** Let us note that we use another parametrization of the Hamiltonian  $H_{X,\beta}$  for proving Propositions 3 (i), (ii) and Theorem 5 (for the details, see [11, Sect. 6]).

8. ON THE NEGATIVE SPECTRUM OF THE OPERATOR  $H_{X,\beta}$

Let  $T$  be a self-adjoint operator in  $\mathfrak{S}$  and let  $E_T(\cdot)$  be its spectral function. Dimension of the negative subspace  $E_T(-\infty, 0)\mathfrak{S}$  of the operator  $T$  is denoted by  $\kappa_-(T)$ ,  $\kappa_-(T) := \dim(E_T(-\infty, 0)\mathfrak{S})$ .

**Proposition 4.** *Let  $d^* < \infty$  and let  $\kappa_-(\beta)$  be the number of negative elements in the sequence  $\beta = \{\beta_n\}_{n=1}^{\infty}$ . If  $H_{X,\beta} = H_{X,\beta}^*$ , then*

$$\kappa_-(H_{X,\beta}) = \kappa_-(\beta). \quad (22)$$

In particular,  $H_{X,\beta} > 0$  if  $\beta_n > 0, n \in \mathbb{N}$ .

**Proof.** Let  $M(\cdot)$  be the Weyl function of the operator  $H_{\min}$ , which corresponds to the boundary triplet  $\Pi$  defined in Theorem 2. Using the form of the Weyl function, we get the following equality  $M(0) := s\text{-}\lim_{x \rightarrow -0} M(x) = 0$ . By [9, Theorem 4],  $\kappa_-(H_{X,\beta}) = \kappa_-(B_{X,\beta} - M(0)) = \kappa_-(B_{X,\beta})$ . On the other hand, by (16), (17),  $\kappa_-(B_{X,\beta}) = \kappa_-(D_{X,\beta}) = \kappa_-(\beta)$ .

**Remark 5.** Using rather different approaches, Proposition 4 was obtained in [15] under assumptions that  $\kappa_-(\beta) = |\mathcal{X}| < \infty$ , and also in [6] assuming that  $|\mathcal{X}| = \infty$  and  $d_* > 0$ .

9. HAMILTONIANS WITH  $\delta$ -INTERACTIONS

Let  $\alpha = \{\alpha_n\}_{n=1}^{\infty} \subset \mathbb{R}$ . Consider the differential expression

$$l_{X,\alpha} = -\frac{d^2}{dx^2} + \sum_{n=1}^{\infty} \alpha_n \delta(x - x_n), \quad x \in \mathcal{F}. \quad (23)$$

In  $L^2(\mathcal{F})$ , one associates with (23) a closed symmetric operator  $H_{X,\alpha}^0$ , defined by the differential expression

$-\frac{d^2}{dx^2}$  on functions  $f \in W_{\text{comp}}^{2,2}(\mathcal{F} \setminus X)$  satisfying the following boundary conditions at the points  $x_n \in X$ :

$$f'(0) = 0, \quad f(x_n+) = f(x_n-), \quad (24)$$

$$f'(x_n+) - f'(x_n-) = \alpha_n f(x_n), \quad n \in \mathbb{N}.$$

Let  $H_{X,\alpha}$  be its closure. The operator  $H_{X,\alpha}$  is known as the Hamiltonian with  $\delta$ -interactions of strengths  $\alpha_n$  at the centers  $x_n$  [2].

The analogs of Lemma 1 and Theorem 3 hold true for the operator  $H_{X,\alpha}$  (see [11, Sect. 5]) and the role of the matrix (15) is played by the Jacobi matrix

$$B_{X,\alpha} = \begin{pmatrix} \frac{1}{r_1^2} \left( \alpha_1 + \frac{1}{d_1} + \frac{1}{d_2} \right) & -\frac{1}{r_1 r_2 d_2} & 0 & \dots \\ -\frac{1}{r_1 r_2 d_2} & \frac{1}{r_2^2} \left( \alpha_2 + \frac{1}{d_2} + \frac{1}{d_3} \right) & -\frac{1}{r_2 r_3 d_3} & \dots \\ 0 & -\frac{1}{r_2 r_3 d_3} & \frac{1}{r_3^2} \left( \alpha_3 + \frac{1}{d_3} + \frac{1}{d_4} \right) & \dots \\ \dots & \dots & \dots & \dots \end{pmatrix}, \tag{25}$$

where  $r_n := \sqrt{d_n + d_{n+1}}$  and  $d_n = x_n - x_{n-1}$ ,  $n \in \mathbb{N}$ . It is easy to see that

$$B_{X,\alpha} = R_X^{-1} ((I - U) D_X^{-1} (I - U^*) + A_\alpha) R_X^{-1}, \tag{26}$$

where  $A_\alpha = \text{diag}(\alpha_n)$ ,  $D_X = \text{diag}(d_n)$ , and  $R_X = \text{diag}(r_n)$ . By (16) and (26), the structure of matrices (15) and (25) are completely different in the case  $d_* = 0$ . Therefore, the spectral properties of Hamiltonians with  $\delta'$  and  $\delta$ -interactions on  $X$  are essentially different in this case (see [11]).

### ACKNOWLEDGMENTS

The first author acknowledges the support from the ESI Junior Research Fellowship Programme and IRCSET Post-Doctoral Fellowship Program.

### REFERENCES

1. N. I. Akhiezer, *The Classical Moment Problem and Some Related Questions in Analysis* (Fizmatgiz, Moscow, 1961; Oliver and Boyd Ltd, Edinburgh, London, 1965).
2. S. Albeverio, F. Gesztesy, R. Hoegh-Krohn, and H. Holden, *Solvable Models in Quantum Mechanics*, Sec. Edition (AMS Chelsea Publ., 2005).
3. S. Albeverio and L. Nizhnik, *Math. Nachr.* **279**, 467 (2006).
4. Ju. M. Berezanskii, *Expansions in Eigenfunctions of Self-Adjoint Operators* (Naukova Dumka, Kiev, 1965;

Translations of Mathematical Monographs, AMS, 1968, Vol. 17).

5. D. Buschmann, G. Stolz, and J. Weidmann, *J. Reine Angew. Math.* **467**, 169 (1995).
6. N. Goloschapova and L. Oridoroga, *Integr. Equations Oper. Theory* **67** (1), 1–14 (2010); DOI 10.1007/s00020-010-1759-x.
7. V. I. Gorbachuk and M. L. Gorbachuk, *Boundary Value Problems for Operator-Differential Equations* (Naukova Dumka, Kiev, 1984; Mathematics and Its Applications (Soviet Series), Kluwer Academic Publishers Group, Dordrecht, 1991, Vol. 48).
8. V. A. Derkach, S. Hassi, M. M. Malamud, and H. S. V. de Snoo, *Trans. Am. Math. Soc.* **358** (12), 5351 (2006).
9. V. A. Derkach and M. M. Malamud, *J. Fund. Anal.* **95**, 1 (1991).
10. I. S. Kac and M. G. Krein, *Izv. VUZov, Ser. Matem.* **2** (3), 136 (1958).
11. A. S. Kostenko and M. Malamud, *J. Differ. Equations* (2010), DOI: 10.1016/j.jde.2010.02.011.
12. A. N. Kochubei, *Ukr. Matem. Zhurn.* **41** (10), 1391 (1989) [*Ukr. Math. J.* **41**, 1391 (1989)].
13. M. M. Malamud and H. Neidhardt, arXiv:0907.0650v1.
14. V. A. Mikhajlets, *Dokl. Akad. Nauk* **348** (6), 727 (1996).
15. L. P. Nizhnik, *Funktsion. Analiz i Ego Prilozh.* **37** (1), 85 (2003) [*Funct. Anal. Appl.* **37** (1), 72 (2003)].